



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:21 pm GMT

PDB ID : 5A86
Title : Structure of pregnane X receptor in complex with a Sphingosine 1- Phosphate Receptor 1 Antagonist
Authors : Xue, Y.; Oster, L.
Deposited on : 2015-07-13
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1062 (2.26-2.26)
Clashscore	112137	1178 (2.26-2.26)
Ramachandran outliers	110173	1145 (2.26-2.26)
Sidechain outliers	110143	1146 (2.26-2.26)
RSRZ outliers	101464	1066 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	
1	B	314	
2	C	17	
2	D	17	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5081 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NUCLEAR RECEPTOR SUBFAMILY 1 GROUP I MEMBER 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	271	Total	C	N	O	S	0	0	0
			2213	1423	375	397	18			
1	B	274	Total	C	N	O	S	0	0	0
			2239	1439	382	400	18			


There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	119	MET	-	EXPRESSION TAG	UNP O75469
A	120	LYS	-	EXPRESSION TAG	UNP O75469
A	121	LYS	-	EXPRESSION TAG	UNP O75469
A	122	GLY	-	EXPRESSION TAG	UNP O75469
A	123	HIS	-	EXPRESSION TAG	UNP O75469
A	124	HIS	-	EXPRESSION TAG	UNP O75469
A	125	HIS	-	EXPRESSION TAG	UNP O75469
A	126	HIS	-	EXPRESSION TAG	UNP O75469
A	127	HIS	-	EXPRESSION TAG	UNP O75469
A	128	HIS	-	EXPRESSION TAG	UNP O75469
A	129	GLY	-	EXPRESSION TAG	UNP O75469
B	119	MET	-	EXPRESSION TAG	UNP O75469
B	120	LYS	-	EXPRESSION TAG	UNP O75469
B	121	LYS	-	EXPRESSION TAG	UNP O75469
B	122	GLY	-	EXPRESSION TAG	UNP O75469
B	123	HIS	-	EXPRESSION TAG	UNP O75469
B	124	HIS	-	EXPRESSION TAG	UNP O75469
B	125	HIS	-	EXPRESSION TAG	UNP O75469
B	126	HIS	-	EXPRESSION TAG	UNP O75469
B	127	HIS	-	EXPRESSION TAG	UNP O75469
B	128	HIS	-	EXPRESSION TAG	UNP O75469
B	129	GLY	-	EXPRESSION TAG	UNP O75469

- Molecule 2 is a protein called NUCLEAR RECEPTOR COACTIVATOR 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	15	Total	C	N	O	0	0	0
			131	82	27	22			
2	D	17	Total	C	N	O	0	0	0
			141	87	29	25			

- Molecule 3 is 4-CHLORO-N-[(1R)-1-[1-ETHYL-6-(TRIFLUOROMETHYL)BENZIMIDAZOL-2-YL]ETHYL]BENZENESULFONAMIDE (three-letter code: D7E) (formula: $C_{18}H_{17}ClF_3N_3O_2S$).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
3	A	1	Total	C	Cl	F	N	O	S	0	0
			28	18	1	3	3	2	1		
3	B	1	Total	C	Cl	F	N	O	S	0	0
			28	18	1	3	3	2	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	125	Total	O	0	0
			125	125		
4	B	133	Total	O	0	0
			133	133		
4	C	5	Total	O	0	0
			5	5		
4	B	1	Total	O	0	0
			1	1		
4	D	2	Total	O	0	0
			2	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total 1	O 1	0	0
4	D	6	Total 6	O 6	0	0
4	A	3	Total 3	O 3	0	0
4	B	1	Total 1	O 1	0	0
4	A	1	Total 1	O 1	0	0
4	B	1	Total 1	O 1	0	0
4	A	1	Total 1	O 1	0	0
4	B	3	Total 3	O 3	0	0
4	A	1	Total 1	O 1	0	0
4	B	5	Total 5	O 5	0	0
4	A	1	Total 1	O 1	0	0
4	B	1	Total 1	O 1	0	0
4	D	1	Total 1	O 1	0	0
4	A	6	Total 6	O 6	0	0
4	B	3	Total 3	O 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: NUCLEAR RECEPTOR SUBFAMILY 1 GROUP I MEMBER 2



• Molecule 1: NUCLEAR RECEPTOR SUBFAMILY 1 GROUP I MEMBER 2



• Molecule 2: NUCLEAR RECEPTOR COACTIVATOR 1



• Molecule 2: NUCLEAR RECEPTOR COACTIVATOR 1





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	83.67Å 89.20Å 106.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.88 – 2.25 14.82 – 2.25	Depositor EDS
% Data completeness (in resolution range)	97.6 (14.88-2.25) 97.6 (14.82-2.25)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.22 (at 2.24Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.231 , 0.256 0.222 , 0.242	Depositor DCC
R_{free} test set	1877 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	43.9	Xtriage
Anisotropy	0.093	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 73.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5081	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: D7E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/2261	0.55	0/3045
1	B	0.40	0/2288	0.54	0/3078
2	C	0.27	0/132	0.50	0/175
2	D	0.36	0/142	0.56	0/188
All	All	0.40	0/4823	0.54	0/6486

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2213	0	2222	38	0
1	B	2239	0	2244	27	0
2	C	131	0	139	1	0
2	D	141	0	147	0	0
3	A	28	0	17	2	0
3	B	28	0	17	3	0
4	A	138	0	0	0	0
4	B	149	0	0	1	0
4	C	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	9	0	0	0	0
All	All	5081	0	4786	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:383:GLN:HB3	1:A:384:PRO:CD	1.91	1.00
1:A:383:GLN:HB3	1:A:384:PRO:HD2	1.59	0.84
1:A:198:LYS:HB3	1:A:417:ILE:HD11	1.60	0.83
1:A:381:ARG:HH11	1:A:381:ARG:HG2	1.46	0.81
1:A:308:LEU:HD13	1:A:324:LEU:HD21	1.64	0.80
1:A:214:GLN:HE21	1:A:216:ARG:HH11	1.33	0.75
1:A:177:VAL:HG21	1:B:215:LEU:HD13	1.74	0.69
2:C:683:LEU:HD12	2:C:686:ARG:HH21	1.60	0.67
1:A:352:ASP:CG	1:A:401:ARG:HH12	1.98	0.67
1:A:383:GLN:HB3	1:A:384:PRO:HD3	1.77	0.66
1:B:199:TRP:HZ3	1:B:233:GLY:O	1.82	0.63
1:A:177:VAL:CG2	1:B:215:LEU:HD13	2.29	0.62
1:B:251:PHE:CD1	1:B:284:CYS:SG	2.94	0.60
1:A:198:LYS:CB	1:A:417:ILE:HD11	2.29	0.60
1:A:235:GLU:HB3	1:B:219:ASP:HB2	1.84	0.59
1:A:211:VAL:HG21	1:A:306:TYR:HB3	1.83	0.59
1:B:254:ILE:HD11	1:B:287:ARG:HD2	1.85	0.59
1:A:254:ILE:HD11	1:A:287:ARG:HD2	1.84	0.58
1:A:235:GLU:CB	1:B:219:ASP:HB2	2.35	0.57
1:B:144:THR:OG1	1:B:147:GLN:HG3	2.07	0.54
1:B:199:TRP:O	1:B:203:ARG:HG3	2.09	0.53
1:A:282:GLU:HG2	1:A:400:LEU:HG	1.90	0.53
1:A:144:THR:OG1	1:A:147:GLN:HG3	2.09	0.53
1:A:214:GLN:HE21	1:A:216:ARG:NH1	2.02	0.52
1:B:282:GLU:HG2	1:B:400:LEU:HG	1.92	0.52
1:A:283:LEU:HD13	1:A:344:GLN:HB3	1.91	0.51
1:A:198:LYS:HB3	1:A:417:ILE:CD1	2.37	0.51
1:A:381:ARG:NH1	1:A:381:ARG:HG2	2.17	0.51
1:A:215:LEU:HD13	1:B:177:VAL:CG2	2.42	0.50
1:B:383:GLN:HB2	1:B:386:HIS:CD2	2.46	0.49
1:B:318:LEU:HB3	1:B:324:LEU:HD12	1.95	0.49
1:A:289:ASN:ND2	1:A:340:TYR:OH	2.46	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:382:PRO:HG2	1:B:383:GLN:OE1	2.12	0.49
1:A:221:SER:HB3	1:B:228:PRO:HD3	1.94	0.48
1:B:383:GLN:CD	1:B:383:GLN:H	2.16	0.48
1:A:318:LEU:HB3	1:A:324:LEU:HD22	1.96	0.47
1:B:207:CYS:HA	1:B:208:SER:HA	1.66	0.47
1:A:209:LEU:HA	3:A:1433:D7E:H11C	1.97	0.46
1:A:308:LEU:CD1	1:A:324:LEU:HD21	2.38	0.46
1:B:202:VAL:HG11	1:B:417:ILE:HD11	1.98	0.46
1:A:251:PHE:CD1	1:A:284:CYS:SG	3.10	0.45
1:B:243:MET:HG3	3:B:1432:D7E:C6	2.46	0.45
1:A:317:GLN:O	1:A:320:LEU:HB2	2.17	0.45
1:A:206:LEU:HD23	1:A:209:LEU:HD12	1.99	0.44
1:B:306:TYR:CD2	3:B:1432:D7E:CL25	3.07	0.44
1:A:219:ASP:HB2	1:B:235:GLU:HB3	1.99	0.44
1:A:306:TYR:CD2	3:A:1433:D7E:CL25	3.08	0.43
1:B:251:PHE:HD1	1:B:284:CYS:SG	2.40	0.43
1:A:308:LEU:HD22	1:A:324:LEU:HD11	2.01	0.42
1:A:202:VAL:O	1:A:206:LEU:HD12	2.19	0.42
1:A:211:VAL:CG2	1:A:306:TYR:HB3	2.49	0.42
1:A:415:GLN:HG2	1:A:431:ILE:HD13	2.01	0.42
1:A:250:MET:HG3	1:A:288:PHE:CZ	2.55	0.42
1:A:381:ARG:HA	1:A:382:PRO:HD2	1.68	0.41
1:B:209:LEU:HB2	3:B:1432:D7E:H11C	2.03	0.41
1:B:254:ILE:HD12	1:B:283:LEU:HB3	2.02	0.41
1:B:250:MET:HG3	1:B:288:PHE:CZ	2.55	0.41
1:B:334:GLN:HA	4:B:2089:HOH:O	2.21	0.41
1:B:206:LEU:HD23	1:B:236:ILE:HB	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/314 (84%)	256 (97%)	7 (3%)	2 (1%)	22	20
1	B	268/314 (85%)	257 (96%)	11 (4%)	0	100	100
2	C	13/17 (76%)	12 (92%)	1 (8%)	0	100	100
2	D	15/17 (88%)	14 (93%)	1 (7%)	0	100	100
All	All	561/662 (85%)	539 (96%)	20 (4%)	2 (0%)	38	42

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	382	PRO
1	A	383	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/278 (88%)	233 (96%)	11 (4%)	32	36
1	B	246/278 (88%)	228 (93%)	18 (7%)	16	15
2	C	15/16 (94%)	14 (93%)	1 (7%)	19	18
2	D	16/16 (100%)	16 (100%)	0	100	100
All	All	521/588 (89%)	491 (94%)	30 (6%)	23	24

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	143	LEU
1	A	234	LYS
1	A	270	GLU
1	A	283	LEU
1	A	303	ARG
1	A	319	LEU
1	A	324	LEU
1	A	332	LYS
1	A	360	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	381	ARG
1	A	383	GLN
1	B	145	GLU
1	B	149	MET
1	B	193	ARG
1	B	194	GLU
1	B	206	LEU
1	B	209	LEU
1	B	210	LYS
1	B	235	GLU
1	B	283	LEU
1	B	308	LEU
1	B	309	GLU
1	B	310	ASP
1	B	315	PHE
1	B	317	GLN
1	B	319	LEU
1	B	324	LEU
1	B	364	GLN
1	B	381	ARG
2	C	691	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	214	GLN
1	A	224	ASN
1	A	272	GLN
1	A	289	ASN
1	A	364	GLN
1	B	383	GLN
1	B	386	HIS
1	B	407	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	D7E	A	1433	-	26,30,30	1.28	4 (15%)	35,46,46	0.91	1 (2%)
3	D7E	B	1432	-	26,30,30	1.27	3 (11%)	35,46,46	1.23	3 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	D7E	A	1433	-	-	0/21/23/23	0/3/3/3
3	D7E	B	1432	-	-	0/21/23/23	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1433	D7E	C11-N10	-2.44	1.31	1.34
3	B	1432	D7E	C11-N10	-2.23	1.31	1.34
3	A	1433	D7E	F27-C26	2.10	1.40	1.32
3	A	1433	D7E	F29-C26	2.16	1.40	1.32
3	B	1432	D7E	F29-C26	2.33	1.41	1.32
3	A	1433	D7E	S16-N15	3.20	1.66	1.61

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1432	D7E	S16-N15	3.80	1.67	1.61

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1432	D7E	C24-C19-S16	-2.70	116.78	119.77
3	B	1432	D7E	O18-S16-C19	-2.46	104.91	107.95
3	A	1433	D7E	F29-C26-C6	-2.05	108.12	112.95
3	B	1432	D7E	C20-C19-S16	2.11	122.11	119.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1433	D7E	2	0
3	B	1432	D7E	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	271/314 (86%)	0.57	31 (11%) 6 5	26, 52, 106, 137	0
1	B	274/314 (87%)	0.46	36 (13%) 4 3	26, 49, 114, 145	0
2	C	15/17 (88%)	1.11	2 (13%) 4 3	47, 60, 92, 95	0
2	D	17/17 (100%)	0.55	1 (5%) 23 26	39, 55, 95, 105	0
All	All	577/662 (87%)	0.53	70 (12%) 5 4	26, 52, 110, 145	0

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	312	ALA	8.6
1	B	233	GLY	7.4
1	B	204	LYS	6.8
2	C	682	SER	6.3
1	B	196	ALA	6.3
1	B	192	SER	6.0
1	A	207	CYS	5.9
1	A	231	SER	5.8
1	B	141	GLN	5.7
1	A	229	ALA	5.5
1	B	315	PHE	5.5
1	A	310	ASP	5.4
1	A	194	GLU	5.2
1	A	208	SER	5.2
1	B	310	ASP	5.1
1	A	311	THR	5.1
1	A	316	GLN	5.0
1	B	194	GLU	4.6
1	A	320	LEU	4.6
1	B	382	PRO	4.4
1	B	193	ARG	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	207	CYS	4.2
1	A	318	LEU	4.2
1	B	385	ALA	4.2
1	B	198	LYS	4.1
1	A	317	GLN	4.1
1	B	205	ASP	3.8
1	A	233	GLY	3.5
1	B	320	LEU	3.4
1	B	149	MET	3.4
1	A	307	CYS	3.4
1	A	319	LEU	3.3
1	A	209	LEU	3.3
1	A	235	GLU	3.3
1	B	231	SER	3.2
1	A	308	LEU	3.2
1	A	218	GLU	3.1
1	B	206	LEU	3.1
1	A	432	THR	3.1
1	B	218	GLU	3.1
1	A	324	LEU	3.1
1	B	430	GLY	3.0
1	B	142	GLY	3.0
1	A	385	ALA	2.9
1	B	384	PRO	2.9
1	B	199	TRP	2.9
1	B	209	LEU	2.9
1	B	317	GLN	2.8
1	A	195	GLU	2.8
1	B	232	GLY	2.7
1	A	429	PHE	2.7
1	A	382	PRO	2.7
1	B	383	GLN	2.6
1	B	316	GLN	2.5
1	B	414	ILE	2.5
1	B	171	ASN	2.4
1	A	211	VAL	2.4
2	D	682	SER	2.4
1	B	314	GLY	2.4
1	A	203	ARG	2.3
1	A	383	GLN	2.3
1	B	176	GLY	2.3
1	A	364	GLN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	234	LYS	2.2
1	A	196	ALA	2.2
1	B	319	LEU	2.1
1	B	168	HIS	2.1
2	C	685	GLU	2.1
1	B	146	GLU	2.1
1	B	197	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	D7E	B	1432	28/28	0.94	0.14	-0.31	59,71,73,74	0
3	D7E	A	1433	28/28	0.94	0.10	-0.94	49,61,73,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.