



Full wwPDB NMR Structure Validation Report ⓘ

Feb 12, 2017 – 08:58 pm GMT

PDB ID : 2A9L
Title : SOLUTION NMR STRUCTURE OF A SUBSTRATE FOR THE ARCHAEAL PRE-TRNA SPLICING ENDONUCLEASES: THE BULGE-HELIX-BULGE MOTIF, MINIMIZED AVERAGE STRUCTURE
Authors : Diener, J.L.; Moore, P.B.
Deposited on : 1998-04-07

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	trunk28760
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR


The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
RNA backbone	3398	623

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	38	 53% 16% 26% 5%

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 1231 atoms, of which 415 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called ABISS7 RNA.


Mol	Chain	Residues	Atoms						Trace
1	A	38	Total	C	H	N	O	P	0
			1231	365	415	155	259	37	

4 Residue-property plots

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: ABISS7 RNA

Chain A: 



5 Refinement protocol and experimental data overview

The models were refined using the following method: *TORSION ANGLE MOLECULAR DYNAMICS*.

Of the 65 calculated structures, 1 were deposited, based on the following criterion: *LEAST RESTRAINT VIOLATION*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	
X-PLOR	structure solution	3.851
CNS	structure solution	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality

6.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.80	0/914 (0.0%)	1.79	26/1425 (1.8%)
All	All	0.80	0/914 (0.0%)	1.79	26/1425 (1.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0	12
All	All	0	12

There are no bond-length outliers.

All angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	31	U	N1-C1'-C2'	8.09	124.52	114.00
1	A	12	G	N9-C1'-C2'	7.90	124.27	114.00
1	A	12	G	C5'-C4'-C3'	7.81	128.50	116.00
1	A	22	G	C1'-O4'-C4'	7.78	116.13	109.90
1	A	13	A	N9-C1'-C2'	7.42	123.64	114.00
1	A	13	A	C4'-C3'-C2'	-7.34	95.26	102.60
1	A	25	C	C6-N1-C2	-7.21	117.42	120.30
1	A	10	C	O4'-C1'-N1	6.73	113.58	108.20
1	A	21	A	N1-C2-N3	6.67	132.63	129.30
1	A	21	A	N1-C6-N6	-6.15	114.91	118.60
1	A	13	A	O4'-C1'-N9	-6.07	103.34	108.20
1	A	6	A	C4'-C3'-C2'	-6.04	96.56	102.60
1	A	21	A	C3'-C2'-C1'	5.98	106.28	101.50
1	A	6	A	O4'-C1'-N9	5.95	112.96	108.20
1	A	31	U	O4'-C1'-N1	-5.72	103.63	108.20
1	A	12	G	C5'-C4'-O4'	5.68	115.92	109.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	11	A	O4'-C1'-N9	-5.64	103.69	108.20
1	A	22	G	C4'-C3'-C2'	-5.64	96.96	102.60
1	A	11	A	C3'-C2'-C1'	-5.63	97.00	101.50
1	A	22	G	O4'-C1'-C2'	-5.54	100.26	105.80
1	A	21	A	N9-C4-C5	5.53	108.01	105.80
1	A	22	G	C5'-C4'-C3'	5.45	124.71	116.00
1	A	13	A	O4'-C1'-C2'	-5.16	100.64	105.80
1	A	7	C	C6-N1-C2	-5.09	118.27	120.30
1	A	14	G	N7-C8-N9	5.05	115.63	113.10
1	A	21	A	P-O3'-C3'	5.02	125.73	119.70

There are no chirality outliers.

All planar outliers are listed below.

Mol	Chain	Res	Type	Group
1	A	19	A	Sidechain
1	A	13	A	Sidechain
1	A	25	C	Sidechain
1	A	12	G	Sidechain
1	A	18	G	Sidechain
1	A	27	G	Sidechain
1	A	23	A	Sidechain
1	A	14	G	Sidechain
1	A	22	G	Sidechain
1	A	31	U	Sidechain
1	A	34	C	Sidechain
1	A	15	G	Sidechain

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	816	415	415	4
All	All	816	415	415	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:21:A:H2'	1:A:22:G:O4'	0.49	2.07
1:A:18:G:N2	1:A:21:A:C6	0.45	2.85
1:A:18:G:N2	1:A:21:A:C5	0.41	2.89
1:A:10:C:H2'	1:A:11:A:O4'	0.40	2.16

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	A	37/38 (97%)	5 (14%)	0 (0%)	0.42
All	All	37/38 (97%)	5 (14%)	0 (0%)	0.42

The overall RNA backbone suiteness is 0.42.

All RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	19	A
1	A	26	G
1	A	13	A
1	A	22	G
1	A	6	A

There are no RNA pucker outliers to report.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided