



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 11:13 am GMT

PDB ID : 5AAU
Title : Optimization of a novel binding motif to to (E)-3-(3,5-difluoro-4-((1R,3R)-2-(2-fluoro-2-methylpropyl)-3-methyl-2,3,4,9-tetrahydro-1H-pyrido(3,4-b)indol-1-yl)phenyl)acrylic acid (AZD9496), a potent and orally bioavailable selective estrogen receptor downregulator and antagonist
Authors : Norman, R.A.; Bradbury, R.H.; de Almeida, C.; Andrews, D.M.; Ballard, P.; Buttar, D.; Callis, R.J.; Currie, G.S.; Curwen, J.O.; Davies, C.D.; de Savi, C.; Donald, C.S.; Feron, L.J.L.; Glossop, S.C.; Hayter, B.R.; Karoutchi, G.; Lamont, S.G.; MacFaul, P.; Moss, T.; Pearson, S.E.; Rabow, A.A.; Tonge, M.; Walker, G.E.; Weir, H.M.; Wilson, Z.
Deposited on : 2015-07-28
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	5047 (1.90-1.90)
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	252	<div> <div>6%</div> <div> <div></div> <div>76%</div> <div>13%</div> <div>•</div> <div>9%</div> </div> </div>
1	B	252	<div> <div>10%</div> <div> <div></div> <div>75%</div> <div>12%</div> <div>•</div> <div>12%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3901 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ESTROGEN RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	229	Total	C	N	O	S	0	10	0
			1865	1198	321	329	17			
1	B	222	Total	C	N	O	S	0	6	0
			1759	1132	295	317	15			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	303	GLY	-	EXPRESSION TAG	UNP P03372
A	304	SER	-	EXPRESSION TAG	UNP P03372
A	305	HIS	-	EXPRESSION TAG	UNP P03372
A	306	MET	-	EXPRESSION TAG	UNP P03372
A	381	SER	CYS	ENGINEERED MUTATION	UNP P03372
A	417	SER	CYS	ENGINEERED MUTATION	UNP P03372
A	536	SER	LEU	ENGINEERED MUTATION	UNP P03372
B	303	GLY	-	EXPRESSION TAG	UNP P03372
B	304	SER	-	EXPRESSION TAG	UNP P03372
B	305	HIS	-	EXPRESSION TAG	UNP P03372
B	306	MET	-	EXPRESSION TAG	UNP P03372
B	381	SER	CYS	ENGINEERED MUTATION	UNP P03372
B	417	SER	CYS	ENGINEERED MUTATION	UNP P03372
B	536	SER	LEU	ENGINEERED MUTATION	UNP P03372

- Molecule 2 is 3-(1-(4-CHLOROPHENYL)-3,4-DIHYDRO-1H-PYRIDO(3,4-B)INDOL-2(9H)-YL)PROPANOIC ACID (three-letter code: XBR) (formula: C₂₀H₁₉ClN₂O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	0
			25	20	1	2	2	
2	B	1	Total	C	Cl	N	O	0
			25	20	1	2	2	


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	137	Total	O	0	0
			137	137		
3	B	90	Total	O	0	0
			90	90		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ESTROGEN RECEPTOR



• Molecule 1: ESTROGEN RECEPTOR



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	106.41 Å 51.40 Å 85.18 Å 90.00° 93.66° 90.00°	Depositor
Resolution (Å)	25.70 – 1.90 25.70 – 1.90	Depositor EDS
% Data completeness (in resolution range)	90.9 (25.70-1.90) 91.0 (25.70-1.90)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 1.91 Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.173 , 0.226 0.205 , 0.250	Depositor DCC
R_{free} test set	1627 reflections (4.92%)	DCC
Wilson B-factor (Å ²)	27.8	Xtriage
Anisotropy	0.131	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 52.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3901	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: XBR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.21	4/1923 (0.2%)	1.16	12/2601 (0.5%)
1	B	1.18	3/1808 (0.2%)	1.09	12/2446 (0.5%)
All	All	1.19	7/3731 (0.2%)	1.12	24/5047 (0.5%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	385	GLU	CD-OE1	-7.59	1.17	1.25
1	A	352	ARG	CZ-NH2	6.58	1.41	1.33
1	B	383	TRP	CB-CG	-6.54	1.38	1.50
1	B	388	MET	C-O	6.01	1.34	1.23
1	A	353	GLU	CD-OE1	5.98	1.32	1.25
1	A	385	GLU	CD-OE2	-5.72	1.19	1.25
1	A	473	ASP	CB-CG	5.12	1.62	1.51

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	436	ARG	NE-CZ-NH1	9.91	125.26	120.30
1	A	436	ARG	NE-CZ-NH2	-7.65	116.48	120.30
1	A	313	ASP	CB-CG-OD1	6.86	124.47	118.30
1	A	515	ARG	CG-CD-NE	-6.66	97.81	111.80
1	B	515	ARG	NE-CZ-NH1	6.59	123.59	120.30
1	A	321[A]	ASP	CB-CG-OD1	6.37	124.04	118.30
1	A	321[B]	ASP	CB-CG-OD1	6.37	124.04	118.30
1	B	394	ARG	CG-CD-NE	6.20	124.81	111.80
1	A	515	ARG	NE-CZ-NH2	-6.19	117.20	120.30
1	A	484	ASP	CB-CG-OD2	-5.83	113.05	118.30
1	B	434	ARG	NE-CZ-NH1	5.77	123.18	120.30
1	B	543	MET	CG-SD-CE	-5.73	91.03	100.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	379	LEU	CB-CG-CD1	5.69	120.67	111.00
1	B	396	MET	CG-SD-CE	5.49	108.98	100.20
1	A	374	ASP	CB-CG-OD2	-5.45	113.39	118.30
1	B	538	ASP	CB-CG-OD1	5.39	123.15	118.30
1	B	515	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	A	434	ARG	NE-CZ-NH2	-5.34	117.63	120.30
1	A	394	ARG	CB-CA-C	-5.33	99.75	110.40
1	B	401	LYS	CD-CE-NZ	-5.22	99.70	111.70
1	A	374	ASP	CB-CG-OD1	5.08	122.87	118.30
1	B	394	ARG	CB-CA-C	-5.04	100.33	110.40
1	B	484	ASP	CB-CG-OD1	5.03	122.83	118.30
1	B	515	ARG	CG-CD-NE	-5.03	101.23	111.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1865	0	1884	26	1
1	B	1759	0	1770	17	1
2	A	25	0	18	1	0
2	B	25	0	18	2	0
3	A	137	0	0	2	0
3	B	90	0	0	1	0
All	All	3901	0	3690	34	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:376:VAL:HG22	1:A:544:LEU:HD12	1.51	0.93
1:A:454:LEU:O	1:A:458:VAL:HG13	1.91	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:497:LEU:HD11	1:B:497:LEU:HD22	1.74	0.70
1:A:454:LEU:O	1:A:458:VAL:CG1	2.41	0.69
1:A:459:TYR:CD2	1:B:513[B]:HIS:CE1	2.83	0.67
1:B:368:VAL:HG22	3:B:2024:HOH:O	2.00	0.62
1:B:397:GLU:OE1	1:B:436[B]:ARG:NH1	2.35	0.59
1:A:459:TYR:HD2	1:B:513[B]:HIS:ND1	2.04	0.55
1:A:459:TYR:OH	1:B:434:ARG:NE	2.40	0.54
1:A:376:VAL:HG22	1:A:544:LEU:CD1	2.33	0.54
1:A:526:TYR:CZ	1:A:534:VAL:HG21	2.43	0.54
1:A:459:TYR:HD2	1:B:513[B]:HIS:CE1	2.25	0.54
1:A:310:LEU:O	1:A:481:LYS:HE3	2.07	0.53
1:A:394:ARG:HG3	3:A:2062:HOH:O	2.09	0.52
3:A:2099:HOH:O	1:B:506[A]:GLN:NE2	2.42	0.51
1:A:459:TYR:HD2	1:B:513[B]:HIS:CG	2.28	0.51
1:A:497:LEU:CD1	1:B:497:LEU:HD22	2.42	0.49
1:B:353[B]:GLU:OE2	2:B:1547:XBR:H2	2.13	0.49
1:A:455:ASN:HA	1:A:458:VAL:HG13	1.96	0.48
1:B:316:VAL:HG21	1:B:489:LEU:HD21	1.96	0.48
1:B:435:PHE:CD1	1:B:440:LEU:HD22	2.49	0.48
1:A:373[B]:HIS:HD2	1:A:374:ASP:OD1	1.97	0.47
1:A:373[B]:HIS:CE1	1:A:377[B]:HIS:NE2	2.82	0.47
1:A:424:ILE:HD13	2:A:1553:XBR:O19	2.15	0.47
1:B:424:ILE:HD13	2:B:1547:XBR:O19	2.14	0.47
1:A:459:TYR:CD2	1:B:513[B]:HIS:ND1	2.83	0.46
1:A:396:MET:HE2	1:A:436:ARG:HA	1.98	0.46
1:B:372:LEU:O	1:B:376:VAL:HG23	2.16	0.46
1:A:525:LEU:HD13	1:A:534:VAL:HG12	1.98	0.45
1:A:310:LEU:O	1:A:481:LYS:CE	2.65	0.44
1:A:519:ASN:HD21	1:B:516:HIS:HA	1.84	0.43
1:A:373[B]:HIS:CD2	1:A:377[B]:HIS:CD2	3.07	0.42
1:A:373[B]:HIS:CD2	1:A:377[B]:HIS:HD2	2.38	0.42
1:A:433:SER:HB2	1:A:436:ARG:NH2	2.35	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:359[A]:ASN:ND2	1:B:328:TYR:OH[3_545]	1.77	0.43

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	231/252 (92%)	229 (99%)	2 (1%)	0	100	100
1	B	220/252 (87%)	219 (100%)	1 (0%)	0	100	100
All	All	451/504 (90%)	448 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	208/226 (92%)	200 (96%)	8 (4%)	38	27
1	B	193/226 (85%)	187 (97%)	6 (3%)	45	36
All	All	401/452 (89%)	387 (96%)	14 (4%)	40	30

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	304	SER
1	A	341	SER
1	A	423	GLU
1	A	433	SER
1	A	471	GLU
1	A	481	LYS
1	A	528	MET
1	A	549	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	330	GLU
1	B	341	SER
1	B	355	VAL
1	B	406	PRO
1	B	413	ASN
1	B	538	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	439	ASN
1	A	474	HIS
1	A	519	ASN
1	A	550	HIS
1	B	519	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	XBR	A	1553	-	24,28,28	1.81	6 (25%)	27,40,40	2.44	10 (37%)
2	XBR	B	1547	-	24,28,28	1.63	7 (29%)	27,40,40	1.78	8 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XBR	A	1553	-	-	0/7/22/22	0/4/4/4
2	XBR	B	1547	-	-	0/7/22/22	0/4/4/4

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1547	XBR	C7-C8	-3.14	1.33	1.38
2	A	1553	XBR	C7-C8	-2.83	1.33	1.38
2	A	1553	XBR	C8-C11	-2.51	1.49	1.51
2	B	1547	XBR	C3-C4	-2.31	1.37	1.41
2	A	1553	XBR	C2-C1	2.03	1.43	1.38
2	A	1553	XBR	C7-C5	2.11	1.45	1.41
2	B	1547	XBR	C22-C23	2.22	1.42	1.38
2	B	1547	XBR	C11-N12	2.28	1.51	1.47
2	B	1547	XBR	C7-C5	2.45	1.46	1.41
2	B	1547	XBR	C14-C7	2.66	1.57	1.51
2	A	1553	XBR	C1-C6	2.67	1.42	1.36
2	B	1547	XBR	C8-C11	2.77	1.54	1.51
2	A	1553	XBR	C25-C20	5.19	1.47	1.39

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1553	XBR	C22-C21-C20	-5.35	115.78	121.20
2	A	1553	XBR	C1-C6-C5	-4.42	114.57	120.88
2	B	1547	XBR	C13-C14-C7	-4.37	104.76	111.65
2	A	1553	XBR	C15-N12-C13	-3.89	102.81	111.07
2	A	1553	XBR	C14-C13-N12	-3.68	102.75	110.32
2	A	1553	XBR	C20-C11-C8	-3.18	107.52	112.93
2	B	1547	XBR	C22-C21-C20	-2.92	118.23	121.20
2	A	1553	XBR	C13-C14-C7	-2.73	107.34	111.65
2	A	1553	XBR	C25-C24-C23	-2.61	116.45	119.24
2	B	1547	XBR	C2-C3-C4	-2.14	116.84	120.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1547	XBR	C25-C20-C21	2.02	120.83	118.30
2	A	1553	XBR	C24-C23-C22	2.13	124.16	121.25
2	B	1547	XBR	C21-C22-C23	2.17	121.56	119.24
2	B	1547	XBR	C15-N12-C13	2.45	116.26	111.07
2	B	1547	XBR	C22-C23-CL	2.55	123.37	119.35
2	B	1547	XBR	C1-C2-C3	2.85	124.50	120.45
2	A	1553	XBR	C2-C1-C6	2.99	124.71	120.45
2	A	1553	XBR	C25-C20-C21	4.57	124.03	118.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1553	XBR	1	0
2	B	1547	XBR	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	229/252 (90%)	0.36	16 (6%) 17 19	15, 29, 53, 69	0
1	B	222/252 (88%)	0.55	26 (11%) 5 5	15, 24, 46, 62	0
All	All	451/504 (89%)	0.45	42 (9%) 9 10	15, 26, 49, 69	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	462	LEU	5.5
1	B	526	TYR	5.3
1	B	331	TYR	4.7
1	B	340	ALA	4.5
1	A	331	TYR	4.5
1	A	461	PHE	4.2
1	A	552	PRO	3.7
1	B	473	ASP	3.7
1	B	321	ASP	3.6
1	B	527	SER	3.4
1	B	337	PHE	3.4
1	B	463	SER	3.3
1	B	546	ALA	3.2
1	B	338	SER	3.1
1	A	386	ILE	3.1
1	B	339	GLU	3.1
1	A	528	MET	3.1
1	A	460	THR	3.0
1	B	341	SER	2.9
1	A	511	LEU	2.9
1	B	417	SER	2.8
1	B	413	ASN	2.8
1	A	459	TYR	2.7
1	B	494	GLY	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	452	ILE	2.7
1	A	391	LEU	2.7
1	A	514	ILE	2.6
1	B	418	VAL	2.6
1	B	320	LEU	2.5
1	B	317	SER	2.4
1	A	384	LEU	2.3
1	B	528	MET	2.3
1	A	392	VAL	2.3
1	B	311	THR	2.3
1	B	345	LEU	2.2
1	A	389	ILE	2.2
1	A	527	SER	2.2
1	B	491	ALA	2.2
1	A	388	MET	2.2
1	A	387	LEU	2.1
1	B	419	GLU	2.0
1	B	511	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	XBR	A	1553	25/25	0.93	0.12	-0.51	18,22,27,28	0
2	XBR	B	1547	25/25	0.95	0.10	-0.62	18,22,27,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.