



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 12:33 am GMT

PDB ID : 2AC3
Title : Structure of human Mnk2 Kinase Domain
Authors : Jauch, R.; Wahl, M.C.; Netter, C.; Jakel, S.; Schreiter, K.; Aicher, B.; Jackle, H.
Deposited on : 2005-07-18
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	<div> <div>18%</div> <div>73%</div> <div>13%</div> <div>•</div> <div>12%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2377 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MAP kinase-interacting serine/threonine kinase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	277	Total	C	N	O	S	27	0	0
			2215	1403	383	414	15			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	70	GLY	-	EXPRESSION TAG	UNP Q9HBH9
A	71	SER	-	EXPRESSION TAG	UNP Q9HBH9

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	161	Total	O	0	0
			161	161		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MAP kinase-interacting serine/threonine kinase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	104.50Å 104.50Å 72.35Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	15.00 – 2.10 19.96 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.6 (15.00-2.10) 97.0 (19.96-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.215 , 0.254 0.249 , 0.243	Depositor DCC
R_{free} test set	1268 reflections (4.88%)	DCC
Wilson B-factor (Å ²)	52.2	Xtriage
Anisotropy	0.110	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 73.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.032 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2377	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/2264	0.58	1/3056 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	90	LEU	CA-CB-CG	5.27	127.42	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2215	0	2141	28	0
2	A	1	0	0	0	0
3	A	161	0	0	3	0
All	All	2377	0	2141	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:103:ASN:HD21	1:A:105:ILE:HG13	1.41	0.84
1:A:79:PHE:O	1:A:82:VAL:O	2.02	0.78
1:A:91:GLY:HA3	1:A:98:VAL:H	1.53	0.72
1:A:139:HIS:HD2	1:A:141:ASN:H	1.44	0.65
1:A:139:HIS:CD2	1:A:141:ASN:H	2.17	0.62
1:A:103:ASN:ND2	1:A:106:THR:H	2.00	0.60
1:A:365:HIS:CD2	1:A:367:TRP:H	2.23	0.56
1:A:178:ASN:ND2	1:A:181:GLU:H	2.03	0.56
1:A:103:ASN:HD21	1:A:105:ILE:CG1	2.15	0.55
1:A:164:GLY:O	1:A:170:HIS:HE1	1.89	0.55
1:A:129:GLU:OE2	3:A:657:HOH:O	2.18	0.54
1:A:102:ILE:CD1	1:A:109:GLU:HG2	2.37	0.53
1:A:83:TYR:HA	1:A:102:ILE:O	2.09	0.52
1:A:365:HIS:HD2	1:A:367:TRP:H	1.56	0.52
1:A:105:ILE:HG13	1:A:106:THR:H	1.73	0.52
1:A:80:GLU:HB2	3:A:671:HOH:O	2.12	0.49
1:A:86:GLN:O	1:A:99:GLN:NE2	2.46	0.48
1:A:102:ILE:HD12	1:A:109:GLU:HG2	1.98	0.45
1:A:103:ASN:ND2	1:A:105:ILE:HG13	2.20	0.44
1:A:178:ASN:HD22	1:A:178:ASN:C	2.20	0.44
1:A:311:CYS:HA	1:A:312:PRO:HD3	1.85	0.43
1:A:324:GLU:HG2	3:A:691:HOH:O	2.19	0.43
1:A:151:GLU:O	1:A:152:GLU:C	2.57	0.43
1:A:119:PRO:HA	1:A:120:GLY:HA2	1.65	0.42
1:A:343:ASP:OD2	1:A:365:HIS:HE1	2.03	0.42
1:A:218:GLN:HG2	1:A:220:SER:O	2.20	0.41
1:A:257:MET:HE3	1:A:261:VAL:HG12	2.01	0.41
1:A:86:GLN:HE22	1:A:102:ILE:HG12	1.86	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	271/316 (86%)	256 (94%)	11 (4%)	4 (2%)	12 6

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	122	ILE
1	A	152	GLU
1	A	87	GLU
1	A	83	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	241/275 (88%)	229 (95%)	12 (5%)	28 26

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	85	LEU
1	A	88	ASP
1	A	90	LEU
1	A	92	GLU
1	A	152	GLU
1	A	175	ARG
1	A	178	ASN
1	A	180	LEU
1	A	212	LEU
1	A	263	GLU
1	A	326	LYS
1	A	357	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	103	ASN
1	A	121	HIS
1	A	135	GLN
1	A	139	HIS
1	A	170	HIS
1	A	178	ASN
1	A	210	ASN
1	A	336	HIS
1	A	355	GLN
1	A	365	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	274/316 (86%)	1.07	58 (21%) 1 1	45, 71, 105, 147	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	89	VAL	10.3
1	A	95	HIS	7.7
1	A	302	ASP	7.6
1	A	105	ILE	6.4
1	A	94	ALA	5.9
1	A	134	TYR	5.4
1	A	92	GLU	5.4
1	A	93	GLY	4.9
1	A	301	SER	4.9
1	A	366	PRO	4.6
1	A	281	LEU	4.5
1	A	149	PHE	4.3
1	A	148	PHE	4.1
1	A	285	LEU	4.1
1	A	189	VAL	4.1
1	A	231	SER	3.9
1	A	265	PHE	3.8
1	A	310	ALA	3.8
1	A	145	LEU	3.7
1	A	251	CYS	3.6
1	A	282	GLY	3.6
1	A	304	GLY	3.5
1	A	305	TRP	3.5
1	A	230	GLY	3.4
1	A	252	GLY	3.4
1	A	119	PRO	3.3
1	A	278	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	279	TRP	3.2
1	A	303	CYS	3.2
1	A	326	LYS	3.2
1	A	283	VAL	3.0
1	A	118	GLN	2.9
1	A	112	VAL	2.7
1	A	213	CYS	2.7
1	A	284	ILE	2.7
1	A	324	GLU	2.6
1	A	90	LEU	2.6
1	A	355	GLN	2.6
1	A	115	ILE	2.5
1	A	328	GLU	2.5
1	A	146	ILE	2.5
1	A	198	ASN	2.4
1	A	185	VAL	2.4
1	A	288	LEU	2.4
1	A	152	GLU	2.3
1	A	287	ILE	2.3
1	A	289	LEU	2.3
1	A	186	VAL	2.3
1	A	136	CYS	2.3
1	A	114	ILE	2.2
1	A	125	ARG	2.2
1	A	339	CYS	2.2
1	A	369	GLN	2.1
1	A	332	LYS	2.1
1	A	142	VAL	2.1
1	A	267	GLU	2.0
1	A	133	LEU	2.0
1	A	147	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ZN	A	531	1/1	0.96	0.12	-0.55	90,90,90,90	0

6.5 Other polymers [i](#)

There are no such residues in this entry.