



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 07:31 pm GMT

PDB ID : 5ACB  
Title : Crystal Structure of the Human Cdk12-Cyclink Complex  
Authors : Dixon Clarke, S.E.; Elkins, J.M.; Pike, A.C.W.; Mackenzie, A.; Goubin, S.; Strain-Damerell, C.; Mahajan, P.; Tallant, C.; Chalk, R.; Wiggers, H.; Kopec, J.; Fitzpatrick, F.; Burgess-Brown, N.; Carpenter, E.P.; von Delft, F.; Arrow-smith, C.H.; Edwards, A.M.; Bountra, C.; Bullock, A.  
Deposited on : 2015-08-14  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2259 (2.70-2.70)
Clashscore	112137	2590 (2.70-2.70)
Ramachandran outliers	110173	2550 (2.70-2.70)
Sidechain outliers	110143	2550 (2.70-2.70)
RSRZ outliers	101464	2275 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	259	<div> <div>3%</div> <div>87%</div> <div>5%</div> <div>8%</div> </div>
1	B	259	<div> <div>93%</div> <div>.</div> <div>.</div> </div>
2	C	340	<div> <div>3%</div> <div>91%</div> <div>5%</div> <div>.</div> </div>
2	D	340	<div> <div>4%</div> <div>89%</div> <div>5%</div> <div>6%</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYCLIN-K.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	237	Total	C	N	O	S	0	0	0
			1907	1245	307	344	11			
1	B	248	Total	C	N	O	S	0	0	0
			2038	1325	337	363	13			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	SER	-	EXPRESSION TAG	UNP G3V5E1
A	10	MET	-	EXPRESSION TAG	UNP G3V5E1
B	9	SER	-	EXPRESSION TAG	UNP G3V5E1
B	10	MET	-	EXPRESSION TAG	UNP G3V5E1


- Molecule 2 is a protein called CYCLIN-DEPENDENT KINASE 12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	325	Total	C	N	O	P S	0	0	0
			2571	1654	424	475	1 17			
2	D	320	Total	C	N	O	P S	0	0	0
			2477	1601	403	456	1 16			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	713	SER	-	EXPRESSION TAG	UNP Q9NYV4
C	714	MET	-	EXPRESSION TAG	UNP Q9NYV4
D	713	SER	-	EXPRESSION TAG	UNP Q9NYV4
D	714	MET	-	EXPRESSION TAG	UNP Q9NYV4

- Molecule 3 is N-[4-[(3R)-3-[[5-CHLORANYL-4-(1H-INDOL-3-YL)PYRIMIDIN-2-YL]AMINO]PIPERIDIN-1-YL]CARBONYLPHENYL]-4-(DIMETHYLAMINO)BUTANAMIDE (three-letter code: 5I1) (formula: C<sub>30</sub>H<sub>34</sub>ClN<sub>7</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	C	1	Total 40	C 30	Cl 1	N 7	O 2	0	0
3	D	1	Total 40	C 30	Cl 1	N 7	O 2	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CYCLIN-K




- Molecule 1: CYCLIN-K



- Molecule 2: CYCLIN-DEPENDENT KINASE 12



- Molecule 2: CYCLIN-DEPENDENT KINASE 12



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.78Å 148.69Å 91.58Å 90.00° 93.81° 90.00°	Depositor
Resolution (Å)	91.38 – 2.70 41.30 – 2.70	Depositor EDS
% Data completeness (in resolution range)	98.6 (91.38-2.70) 98.7 (41.30-2.70)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.97 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0123	Depositor
R, $R_{free}$	0.221 , 0.262 0.223 , 0.264	Depositor DCC
$R_{free}$ test set	1756 reflections (5.13%)	DCC
Wilson B-factor (Å <sup>2</sup> )	54.3	Xtriage
Anisotropy	0.590	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 29.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	9073	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, 5I1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.38	0/1959	0.54	0/2667
1	B	0.38	0/2095	0.58	0/2842
2	C	0.38	0/2619	0.58	0/3546
2	D	0.37	0/2524	0.57	0/3429
All	All	0.38	0/9197	0.57	0/12484

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1907	0	1803	5	0
1	B	2038	0	1981	1	0
2	C	2571	0	2494	5	0
2	D	2477	0	2336	7	0
3	C	40	0	0	1	0
3	D	40	0	0	1	0
All	All	9073	0	8614	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:52:GLU:OE1	1:B:92:ARG:NH1	2.38	0.57
1:A:52:GLU:OE1	1:A:92:ARG:NH1	2.37	0.56
2:C:865:ILE:CD1	2:C:925:ILE:HD13	2.39	0.53
2:D:865:ILE:CD1	2:D:925:ILE:HD13	2.40	0.52
2:D:986:PHE:O	2:D:989:ILE:HG22	2.12	0.49
2:C:986:PHE:O	2:C:989:ILE:HG22	2.12	0.49
2:C:777:ILE:HD13	2:C:881:ALA:HB3	1.97	0.47
2:D:777:ILE:HD13	2:D:881:ALA:HB3	1.97	0.46
3:D:2000:5I1:CL5	3:D:2000:5I1:CBE	3.03	0.43
1:A:145:ARG:NH2	2:D:760:LEU:HD22	2.33	0.43
2:C:865:ILE:HD11	2:C:925:ILE:HD13	2.00	0.43
1:A:233:ARG:HD2	1:A:235:TRP:CH2	2.54	0.43
2:D:865:ILE:HD11	2:D:925:ILE:HD13	2.01	0.43
3:C:2000:5I1:CBE	3:C:2000:5I1:CL5	3.04	0.42
2:C:903:PRO:HD2	2:C:906:LEU:HD12	2.02	0.42
2:D:903:PRO:HD2	2:D:906:LEU:HD12	2.02	0.41
1:A:142:VAL:HG11	2:D:803:LYS:HA	2.02	0.41
1:A:72:ALA:O	1:A:76:ILE:HG12	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	233/259 (90%)	229 (98%)	4 (2%)	0	100	100
1	B	246/259 (95%)	243 (99%)	3 (1%)	0	100	100
2	C	322/340 (95%)	300 (93%)	17 (5%)	5 (2%)	11	28
2	D	313/340 (92%)	293 (94%)	18 (6%)	2 (1%)	28	56
All	All	1114/1198 (93%)	1065 (96%)	42 (4%)	7 (1%)	28	56



All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	1036	TRP
2	D	795	ASP
2	C	795	ASP
2	C	799	ALA
2	C	897	ILE
2	D	897	ILE
2	C	1039	CYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/233 (84%)	191 (98%)	4 (2%)	59	85
1	B	216/233 (93%)	210 (97%)	6 (3%)	49	79
2	C	270/307 (88%)	266 (98%)	4 (2%)	70	90
2	D	249/307 (81%)	245 (98%)	4 (2%)	68	89
All	All	930/1080 (86%)	912 (98%)	18 (2%)	62	87

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	GLN
1	A	121	ARG
1	A	134	ASP
1	A	245	ASP
1	B	37	GLN
1	B	121	ARG
1	B	134	ASP
1	B	233	ARG
1	B	245	ASP
1	B	260	GLN
2	C	800	LEU
2	C	820	LEU
2	C	981	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	C	1028	MET
2	D	762	ASN
2	D	820	LEU
2	D	981	ARG
2	D	1043	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	GLN
1	B	89	GLN
1	B	156	GLN
2	C	783	HIS
2	C	977	GLN
2	D	762	ASN
2	D	783	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	TPO	C	893	2	9,10,11	0.79	0	10,14,16	1.10	1 (10%)
2	TPO	D	893	2	9,10,11	0.90	0	10,14,16	1.08	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TPO	C	893	2	-	0/8/11/13	0/0/0/0
2	TPO	D	893	2	-	0/8/11/13	0/0/0/0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	893	TPO	O-C-CA	-2.15	120.14	125.15
2	C	893	TPO	O-C-CA	-2.11	120.22	125.15

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	5I1	C	2000	2	41,44,44	1.94	10 (24%)	52,61,61	2.03	10 (19%)
3	5I1	D	2000	2	41,44,44	1.90	10 (24%)	52,61,61	1.82	11 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	5I1	C	2000	2	-	0/22/36/36	0/5/5/5
3	5I1	D	2000	2	-	0/22/36/36	0/5/5/5

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	2000	5I1	CAM-CAH	-7.01	1.39	1.50
3	D	2000	5I1	CAM-CAH	-6.44	1.39	1.50
3	D	2000	5I1	CAX-CAW	-4.23	1.34	1.41
3	C	2000	5I1	CAX-CAW	-4.14	1.34	1.41
3	C	2000	5I1	CAJ-NBI	-3.76	1.34	1.41
3	D	2000	5I1	CBD-CAV	-3.42	1.35	1.42
3	C	2000	5I1	CBD-CAV	-3.40	1.35	1.42
3	D	2000	5I1	CAJ-NBI	-3.07	1.35	1.41
3	D	2000	5I1	CAV-CAW	-2.66	1.35	1.42
3	C	2000	5I1	CAV-CAW	-2.65	1.35	1.42
3	D	2000	5I1	CAB-NAD	2.01	1.49	1.46
3	C	2000	5I1	C4-C5	2.15	1.41	1.38
3	D	2000	5I1	CBJ-CAP	2.18	1.55	1.51
3	C	2000	5I1	CBJ-CAP	2.30	1.55	1.51
3	D	2000	5I1	C6-N1	2.40	1.39	1.34
3	C	2000	5I1	C6-N1	2.43	1.39	1.34
3	C	2000	5I1	CAB-CBH	2.56	1.55	1.51
3	D	2000	5I1	C4-C5	2.83	1.42	1.38
3	C	2000	5I1	CAB-NAD	2.99	1.50	1.46
3	D	2000	5I1	CBG-NAD	3.40	1.53	1.47

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2000	5I1	N1-C2-N3	-6.90	120.04	126.68
3	D	2000	5I1	N1-C2-N3	-6.29	120.63	126.68
3	C	2000	5I1	CAT-CBA-CBG	-4.90	104.51	110.96
3	D	2000	5I1	C5-C4-N3	-4.69	116.11	121.48
3	C	2000	5I1	C5-C4-N3	-4.54	116.28	121.48
3	C	2000	5I1	OAE-CAH-CAM	-3.77	113.14	120.21
3	C	2000	5I1	CBL-CBJ-CAP	-2.98	104.97	113.32
3	D	2000	5I1	C6-C5-CL5	-2.93	114.32	118.47
3	D	2000	5I1	CAT-CBH-CAB	-2.42	106.42	109.63
3	D	2000	5I1	C5-C6-N1	-2.38	120.45	122.84
3	C	2000	5I1	C6-C5-CL5	-2.31	115.21	118.47

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	2000	5I1	CBL-CBJ-CAP	-2.20	107.13	113.32
3	D	2000	5I1	CAG-NAR-CBM	-2.12	102.41	110.75
3	C	2000	5I1	C5-C6-N1	-2.11	120.72	122.84
3	D	2000	5I1	CBN-NAR-CAG	2.24	115.73	109.72
3	C	2000	5I1	CAI-C4-N3	2.83	119.33	114.94
3	D	2000	5I1	CAI-C4-N3	2.99	119.58	114.94
3	C	2000	5I1	NBB-C2-N3	3.11	121.54	117.05
3	D	2000	5I1	NBB-C2-N3	3.54	122.17	117.05
3	D	2000	5I1	CBG-NAD-CAB	3.57	120.82	113.00
3	C	2000	5I1	CAM-CAH-NAD	5.53	125.93	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	2000	5I1	1	0
3	D	2000	5I1	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	237/259 (91%)	0.08	7 (2%) 51 50	37, 58, 87, 108	0
1	B	248/259 (95%)	-0.06	1 (0%) 92 93	31, 47, 78, 129	0
2	C	324/340 (95%)	0.19	10 (3%) 49 49	39, 60, 95, 141	0
2	D	319/340 (93%)	0.26	14 (4%) 35 33	45, 69, 106, 129	0
All	All	1128/1198 (94%)	0.13	32 (2%) 53 54	31, 60, 97, 141	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	826	SER	4.7
2	C	800	LEU	4.4
2	D	1044	SER	4.0
2	D	805	ASP	4.0
2	C	799	ALA	4.0
2	D	891	PRO	3.7
1	A	171	LEU	3.7
1	A	222	ILE	3.2
2	C	1038	ASP	3.2
1	A	227	SER	3.2
2	C	891	PRO	3.2
1	B	261	GLY	3.0
2	C	749	THR	2.9
2	C	748	ASP	2.7
1	A	238	PHE	2.6
2	D	846	GLY	2.6
1	A	237	GLN	2.5
2	D	975	LYS	2.4
2	D	732	ILE	2.4
2	D	733	ILE	2.4
2	D	1041	GLU	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	D	804	LYS	2.4
1	A	239	VAL	2.4
2	D	748	ASP	2.3
1	A	252	HIS	2.3
2	D	886	SER	2.3
2	D	1032	ASP	2.3
2	C	750	GLY	2.3
2	D	968	TYR	2.2
2	D	729	ILE	2.1
2	C	1040	HIS	2.1
2	C	839	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	TPO	C	893	11/12	0.92	0.14	-	74,78,82,85	0
2	TPO	D	893	11/12	0.95	0.13	-	76,78,79,82	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	5I1	D	2000	40/40	0.91	0.33	1.57	71,77,112,113	0
3	5I1	C	2000	40/40	0.89	0.26	1.52	68,76,120,121	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.