



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 09:32 pm GMT

PDB ID : 4AE6
Title : A Crystal Structure of the Sperm Specific Isoform of Protein Kinase A (PKA) Subunit CalphaS Reveals a Binding Pocket for Hydrophobic Moieties
Authors : Hereng, T.H.; Backe, P.H.; Kahmann, J.; Scheich, C.; Bjoras, M.; Skalhegg, B.S.; Rosendal, K.R.
Deposited on : 2012-01-09
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	<div> <div>2%</div> <div> <div></div> <div>79%</div> <div>13%</div> <div>6%</div> </div> </div>
1	B	343	<div> <div>2%</div> <div> <div></div> <div>67%</div> <div>24%</div> <div>6%</div> </div> </div>

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 5683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CAMP-DEPENDENT PROTEIN KINASE CATALYTIC SUB-UNIT ALPHA 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	321	Total	C	N	O	P	S	0	4	0
			2680	1747	449	474	2	8			
1	B	323	Total	C	N	O	P	S	0	9	0
			2744	1789	462	482	2	9			

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	140	Total	O	0	0
			140	140		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	115	Total	O	0	0
			115	115		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CAMP-DEPENDENT PROTEIN KINASE CATALYTIC SUBUNIT ALPHA 2



• Molecule 1: CAMP-DEPENDENT PROTEIN KINASE CATALYTIC SUBUNIT ALPHA 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	75.78Å 89.50Å 114.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	63.24 – 2.10 63.24 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.6 (63.24-2.10) 99.6 (63.24-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.81 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.215 , 0.283 0.215 , 0.283	Depositor DCC
R_{free} test set	2337 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	34.6	Xtriage
Anisotropy	0.088	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 52.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5683	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, ACT, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.06	1/2740 (0.0%)	0.97	5/3689 (0.1%)
1	B	1.05	1/2817 (0.0%)	0.89	3/3789 (0.1%)
All	All	1.05	2/5557 (0.0%)	0.93	8/7478 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	165	ARG	CD-NE	-5.41	1.37	1.46
1	B	110	PHE	CE2-CZ	5.19	1.47	1.37

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	165	ARG	NE-CZ-NH2	-14.24	113.18	120.30
1	A	165	ARG	NE-CZ-NH1	11.10	125.85	120.30
1	A	184	ASP	CB-CG-OD1	7.72	125.25	118.30
1	A	217	LYS	CD-CE-NZ	-6.81	96.03	111.70
1	B	128	MET	CG-SD-CE	-5.59	91.26	100.20
1	A	183	THR	N-CA-C	5.31	125.33	111.00
1	B	41[A]	ASP	CB-CG-OD1	5.16	122.94	118.30
1	B	41[B]	ASP	CB-CG-OD1	5.16	122.94	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2680	0	2690	35	0
1	B	2744	0	2765	76	0
2	B	4	0	3	0	0
3	A	140	0	0	11	1
3	B	115	0	0	10	1
All	All	5683	0	5458	109	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:190:ARG:HH11	1:B:190:ARG:CG	1.77	0.98
1:B:131:HIS:CE1	1:B:135:ILE:HG13	2.00	0.97
1:B:190:ARG:HH11	1:B:190:ARG:HG2	1.35	0.92
1:B:103:LEU:HD22	1:B:185:PHE:HZ	1.36	0.90
1:B:23:LYS:O	1:B:27[A]:LEU:HD13	1.72	0.89
1:A:270:ARG:NH2	1:B:241:ASP:HB3	1.92	0.84
1:A:221[A]:TRP:HZ3	3:A:2088:HOH:O	1.63	0.79
1:B:242:GLN:HB2	1:B:245:GLN:HG3	1.65	0.79
1:A:221[A]:TRP:CZ3	3:A:2088:HOH:O	2.34	0.79
1:A:50:GLY:HA3	3:A:2033:HOH:O	1.83	0.78
1:B:118:MET:SD	3:B:2032:HOH:O	2.41	0.78
1:A:164:TYR:O	1:A:165:ARG:HB2	1.84	0.77
1:B:144:ARG:HD3	1:B:300:THR:CG2	2.17	0.75
1:B:211:LEU:O	1:B:213:LYS:N	2.23	0.72
1:B:222[A]:TRP:HZ3	3:B:2082:HOH:O	1.73	0.72
1:B:103:LEU:HD22	1:B:185:PHE:CZ	2.23	0.71
1:B:61[B]:LYS:HE3	1:B:68:HIS:NE2	2.06	0.70
1:A:99:ASN:OD1	1:A:105[A]:LYS:NZ	2.25	0.70
1:B:144:ARG:HD3	1:B:300:THR:HG23	1.74	0.69
1:B:142:HIS:CE1	1:B:146:TYR:CE2	2.82	0.67
1:A:286:ASN:HB3	1:A:290:ASP:OD1	1.95	0.67
1:B:227:LEU:O	1:B:231[A]:MET:HG3	1.95	0.66
1:B:142:HIS:HE1	1:B:146:TYR:CE2	2.14	0.66
1:B:298:ALA:HA	3:B:2098:HOH:O	1.94	0.66
1:A:201:THR:HG23	3:A:2093:HOH:O	1.96	0.66
1:A:46:ILE:HD11	1:A:61:LYS:HB2	1.79	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:99:ASN:ND2	3:A:2061:HOH:O	2.28	0.65
1:A:51:THR:HG23	3:A:2034:HOH:O	1.98	0.64
1:B:131:HIS:CE1	1:B:135:ILE:CG1	2.78	0.63
1:B:274:GLN:HG3	1:B:276:ASP:H	1.66	0.61
1:B:20:ALA:HA	3:B:2002:HOH:O	2.01	0.60
1:B:23:LYS:HG2	1:B:27[A]:LEU:CD1	2.31	0.60
1:B:222[B]:TRP:O	1:B:222[B]:TRP:CD1	2.56	0.58
1:A:253:GLY:O	3:A:2117:HOH:O	2.17	0.58
1:B:207:PRO:HG2	1:B:275:VAL:HG13	1.85	0.58
1:B:23:LYS:HG2	1:B:27[A]:LEU:HD11	1.84	0.58
1:B:278:THR:O	1:B:284:LEU:CD2	2.51	0.58
1:B:190:ARG:HH11	1:B:190:ARG:HG3	1.67	0.58
1:B:190:ARG:NH1	1:B:190:ARG:CG	2.49	0.57
1:B:80:VAL:HG22	1:B:85:ILE:HD11	1.87	0.57
1:B:18:PHE:HZ	1:B:306:TYR:HH	1.53	0.56
1:B:131:HIS:ND1	1:B:134:ARG:NH2	2.54	0.56
1:B:142:HIS:HE1	1:B:146:TYR:CZ	2.23	0.55
1:B:210:ILE:HD12	1:B:247:TYR:CG	2.42	0.55
1:A:275:VAL:HG23	3:A:2120:HOH:O	2.07	0.54
1:A:227:LEU:O	1:A:231:MET:HG3	2.09	0.52
1:A:78:LYS:NZ	1:A:82:LEU:HD11	2.23	0.52
1:B:210:ILE:CD1	1:B:247:TYR:HB3	2.39	0.52
1:B:155:GLU:HG3	1:B:288:VAL:HG21	1.91	0.52
1:B:208:GLU:CD	1:B:280:ARG:HH22	2.13	0.52
1:B:61[B]:LYS:HE3	1:B:68:HIS:CE1	2.45	0.52
1:B:160:LEU:O	1:B:161:ASP:HB2	2.10	0.51
1:A:134:ARG:HA	1:A:134:ARG:NE	2.25	0.51
1:B:25:ASP:N	1:B:25:ASP:OD1	2.43	0.51
1:B:210:ILE:O	1:B:210:ILE:HG13	2.10	0.50
1:B:222[A]:TRP:O	1:B:225:GLY:N	2.45	0.50
1:A:127:GLU:HG2	1:A:129:PHE:H	1.76	0.50
1:B:222[A]:TRP:CH2	1:B:238:PHE:CE2	3.01	0.49
1:A:265:LEU:HD13	1:A:296:TRP:CE2	2.48	0.48
1:A:126:GLY:HA3	1:A:131:HIS:HE2	1.77	0.48
1:A:23:LYS:HB2	3:A:2007:HOH:O	2.13	0.48
1:A:127:GLU:CD	1:A:129:PHE:HB3	2.34	0.48
1:B:94:ILE:HD13	1:B:188:ALA:HB3	1.96	0.47
1:A:275:VAL:HG22	3:A:2115:HOH:O	2.14	0.47
1:A:342:LYS:HE3	3:A:2133:HOH:O	2.14	0.47
1:B:131:HIS:HB2	3:B:2058:HOH:O	2.14	0.47
1:B:224:LEU:O	1:B:228:ILE:HG13	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:105:LYS:HB2	1:B:105:LYS:HE3	1.31	0.47
1:B:36:ASN:OD1	3:B:2009:HOH:O	2.20	0.47
1:B:190:ARG:HD2	3:B:2004:HOH:O	2.16	0.46
1:B:278:THR:O	1:B:284:LEU:HD21	2.15	0.46
1:B:142:HIS:CE1	1:B:146:TYR:CZ	3.04	0.46
1:B:105:LYS:HG2	1:B:107:GLU:OE1	2.16	0.46
1:B:243:PRO:HD2	3:B:2092:HOH:O	2.15	0.46
1:B:229:TYR:C	1:B:229:TYR:CD1	2.89	0.45
1:B:305:ILE:HD13	1:B:310:VAL:HG21	1.98	0.45
1:B:190:ARG:NH1	1:B:190:ARG:HG2	2.15	0.45
1:A:270:ARG:HH21	1:B:241:ASP:HB3	1.78	0.45
1:A:307:GLN:OE1	1:A:309:LYS:CE	2.65	0.45
1:B:222[B]:TRP:C	1:B:222[B]:TRP:CD1	2.90	0.45
1:A:138:PHE:HD2	1:A:231:MET:HB3	1.82	0.44
1:B:222[B]:TRP:CG	1:B:222[B]:TRP:O	2.70	0.44
1:B:290:ASP:HB2	3:B:2095:HOH:O	2.18	0.44
1:A:138:PHE:CD2	1:A:231:MET:HB3	2.52	0.43
1:A:115:ASN:HB2	1:A:117:TYR:CZ	2.54	0.43
1:B:133:ARG:HG3	3:B:2061:HOH:O	2.17	0.43
1:A:307:GLN:OE1	1:A:309:LYS:HE2	2.19	0.43
1:B:156:TYR:CZ	1:B:160:LEU:HD11	2.53	0.43
1:B:228:ILE:HB	1:B:269:LEU:HD21	2.01	0.43
1:A:342:LYS:HD2	1:A:342:LYS:N	2.31	0.42
1:B:154:PHE:CE2	1:B:220:ASP:HB3	2.53	0.42
1:B:235:TYR:HB2	1:B:236:PRO:HD2	2.01	0.42
1:A:127:GLU:OE1	1:A:129:PHE:HB3	2.19	0.42
1:B:277:LEU:HD23	1:B:283:ASN:HD21	1.83	0.42
1:A:45:ARG:HG2	1:A:58:MET:HE2	2.01	0.42
1:B:190:ARG:NH1	1:B:190:ARG:HG3	2.29	0.42
1:B:61[A]:LYS:HB3	1:B:61[A]:LYS:HE3	1.74	0.42
1:A:138:PHE:HB3	1:A:142:HIS:HB3	2.00	0.42
1:A:142:HIS:CD2	1:A:146:TYR:CE2	3.08	0.42
1:B:277:LEU:HD23	1:B:283:ASN:ND2	2.35	0.42
1:B:274:GLN:HG2	1:B:279:LYS:HB2	2.02	0.41
1:B:131:HIS:CD2	1:B:138:PHE:CE1	3.08	0.41
1:B:276:ASP:OD1	1:B:278:THR:OG1	2.28	0.41
1:B:142:HIS:HE1	1:B:146:TYR:OH	2.04	0.41
1:B:23:LYS:HG2	1:B:27[A]:LEU:HD13	2.01	0.41
1:A:126:GLY:HA3	1:A:131:HIS:NE2	2.36	0.40
1:B:207:PRO:CG	1:B:275:VAL:HG13	2.50	0.40
1:B:276:ASP:OD1	1:B:278:THR:N	2.54	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:135:ILE:HG23	1:B:135:ILE:HD12	1.91	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:2061:HOH:O	3:B:2112:HOH:O[3_545]	1.92	0.28

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	319/343 (93%)	308 (97%)	10 (3%)	1 (0%)	44	44
1	B	326/343 (95%)	302 (93%)	22 (7%)	2 (1%)	28	24
All	All	645/686 (94%)	610 (95%)	32 (5%)	3 (0%)	32	28

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	212	SER
1	A	125	GLY
1	B	253	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	286/301 (95%)	272 (95%)	14 (5%)	29	26
1	B	293/301 (97%)	262 (89%)	31 (11%)	8	4
All	All	579/602 (96%)	534 (92%)	45 (8%)	15	11

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	SER
1	A	51	THR
1	A	56	ARG
1	A	127	GLU
1	A	133	ARG
1	A	183	THR
1	A	241	ASP
1	A	243	PRO
1	A	252	SER
1	A	255	VAL
1	A	263	SER
1	A	285	LYS
1	A	339	ILE
1	A	342	LYS
1	B	16	LYS
1	B	21	LYS
1	B	24	GLU
1	B	25	ASP
1	B	31	GLU
1	B	36	ASN
1	B	61[A]	LYS
1	B	61[B]	LYS
1	B	83	LYS
1	B	86	GLU
1	B	105	LYS
1	B	133	ARG
1	B	184	ASP
1	B	190	ARG
1	B	195	THR
1	B	210	ILE
1	B	212	SER
1	B	217	LYS
1	B	222[A]	TRP
1	B	222[B]	TRP
1	B	239	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	254	LYS
1	B	255	VAL
1	B	277	LEU
1	B	278	THR
1	B	286	ASN
1	B	295	LYS
1	B	299	THR
1	B	331	GLU
1	B	343	CYS
1	B	348	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	115	ASN
1	A	142	HIS
1	A	286	ASN
1	B	115	ASN
1	B	142	HIS
1	B	181	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPO	A	197	1	9,10,11	1.30	0	10,14,16	1.61	3 (30%)
1	SEP	A	338	1	9,9,10	1.72	3 (33%)	9,12,14	2.79	3 (33%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TPO	B	197	1	9,10,11	1.13	1 (11%)	10,14,16	1.67	2 (20%)
1	SEP	B	338	1	9,9,10	1.39	2 (22%)	9,12,14	1.44	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	197	1	-	0/8/11/13	0/0/0/0
1	SEP	A	338	1	-	0/5/8/10	0/0/0/0
1	TPO	B	197	1	-	0/8/11/13	0/0/0/0
1	SEP	B	338	1	-	0/5/8/10	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	338	SEP	P-O3P	2.24	1.64	1.54
1	B	338	SEP	P-OG	2.27	1.67	1.60
1	A	338	SEP	P-O1P	2.59	1.59	1.50
1	B	338	SEP	P-O1P	2.73	1.60	1.50
1	B	197	TPO	CA-C	2.78	1.53	1.50
1	A	338	SEP	CA-C	2.86	1.54	1.50

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	338	SEP	O2P-P-OG	-3.01	98.73	106.73
1	B	197	TPO	CG2-CB-CA	-2.76	108.09	113.22
1	B	338	SEP	OG-CB-CA	2.02	110.15	108.17
1	A	197	TPO	CG2-CB-CA	2.15	117.20	113.22
1	A	197	TPO	O3P-P-O2P	2.26	116.73	107.61
1	B	338	SEP	O3P-P-O2P	2.69	118.46	107.61
1	A	197	TPO	C-CA-N	3.01	115.94	109.86
1	B	197	TPO	C-CA-N	3.13	116.17	109.86
1	A	338	SEP	OG-P-O1P	3.34	115.85	106.47
1	A	338	SEP	OG-CB-CA	5.94	114.03	108.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	B	1351	-	1,3,3	2.80	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACT	B	1351	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1351	ACT	CH3-C	2.80	1.52	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	319/343 (93%)	0.24	6 (1%) 67 71	18, 35, 55, 77	0
1	B	321/343 (93%)	0.37	6 (1%) 67 71	20, 36, 60, 80	0
All	All	640/686 (93%)	0.30	12 (1%) 67 71	18, 36, 59, 80	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	318	PHE	8.3
1	A	283[A]	ASN	3.5
1	B	222[A]	TRP	2.7
1	B	18	PHE	2.6
1	A	125	GLY	2.6
1	B	281	PHE	2.5
1	A	339	ILE	2.5
1	B	22	ALA	2.4
1	B	254	LYS	2.2
1	A	131	HIS	2.1
1	B	331	GLU	2.0
1	A	53	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	SEP	B	338	10/11	0.97	0.12	-	28,30,33,33	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	TPO	B	197	11/12	0.94	0.11	-	44,47,50,51	0
1	SEP	A	338	10/11	0.93	0.11	-	36,42,44,45	0
1	TPO	A	197	11/12	0.98	0.12	-	25,28,30,30	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ACT	B	1351	4/4	0.96	0.11	-0.53	29,31,31,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.