



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 02:22 pm GMT

PDB ID : 3AI3
Title : The crystal structure of L-Sorbose reductase from *Gluconobacter frateurii* complexed with NADPH and L-sorbose
Authors : Kubota, K.; Nagata, K.; Okai, M.; Miyazono, K.; Tanokura, M.
Deposited on : 2010-05-07
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	263	<div> <div style="width: 2%; background-color: red;"></div> <div style="width: 94%; background-color: green;"></div> <div style="width: 5%; background-color: yellow;"></div> <div style="width: 1%; background-color: orange;"></div> <div style="width: 1%; background-color: grey;"></div> </div> <div> <div style="width: 2%; background-color: red;"></div> <div style="width: 92%; background-color: green;"></div> <div style="width: 7%; background-color: yellow;"></div> <div style="width: 1%; background-color: orange;"></div> <div style="width: 1%; background-color: grey;"></div> </div>
1	C	263	<div> <div style="width: 94%; background-color: green;"></div> <div style="width: 5%; background-color: yellow;"></div> </div> <div> <div style="width: 94%; background-color: green;"></div> <div style="width: 5%; background-color: yellow;"></div> </div>
1	E	263	<div> <div style="width: 2%; background-color: red;"></div> <div style="width: 92%; background-color: green;"></div> <div style="width: 7%; background-color: yellow;"></div> <div style="width: 1%; background-color: orange;"></div> <div style="width: 1%; background-color: grey;"></div> </div> <div> <div style="width: 2%; background-color: red;"></div> <div style="width: 93%; background-color: green;"></div> <div style="width: 6%; background-color: yellow;"></div> <div style="width: 1%; background-color: orange;"></div> <div style="width: 1%; background-color: grey;"></div> </div>
1	G	263	<div> <div style="width: 2%; background-color: red;"></div> <div style="width: 93%; background-color: green;"></div> <div style="width: 6%; background-color: yellow;"></div> <div style="width: 1%; background-color: orange;"></div> <div style="width: 1%; background-color: grey;"></div> </div> <div> <div style="width: 2%; background-color: red;"></div> <div style="width: 93%; background-color: green;"></div> <div style="width: 6%; background-color: yellow;"></div> <div style="width: 1%; background-color: orange;"></div> <div style="width: 1%; background-color: grey;"></div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SOL	A	0	-	-	-	X
3	SOL	C	0	-	-	-	X
3	SOL	E	0	-	-	-	X
3	SOL	E	265	-	-	-	X
3	SOL	G	0	-	-	-	X
4	SOE	G	266	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8624 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NADPH-sorbose reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	0	0	0
			1989	1266	340	372	11			
1	C	263	Total	C	N	O	S	0	0	0
			1989	1266	340	372	11			
1	E	263	Total	C	N	O	S	0	0	0
			1989	1266	340	372	11			
1	G	263	Total	C	N	O	S	0	0	0
			1989	1266	340	372	11			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	116	LEU	HIS	ENGINEERED MUTATION	UNP A4PB64
C	116	LEU	HIS	ENGINEERED MUTATION	UNP A4PB64
E	116	LEU	HIS	ENGINEERED MUTATION	UNP A4PB64
G	116	LEU	HIS	ENGINEERED MUTATION	UNP A4PB64

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C₂₁H₃₀N₇O₁₇P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	E	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	G	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is SUGAR (L-SORBOSE) (three-letter code: SOL) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			12	6	6		
3	C	1	Total	C	O	0	0
			12	6	6		
3	E	1	Total	C	O	0	0
			12	6	6		
3	E	1	Total	C	O	0	0
			12	6	6		
3	G	1	Total	C	O	0	0
			12	6	6		

- Molecule 4 is SUGAR (ALPHA-L-SORBOPYRANOSE) (three-letter code: SOE) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	E	1	Total	C	O	0	0
			12	6	6		
4	G	1	Total	C	O	0	0
			12	6	6		
4	G	1	Total	C	O	0	0
			12	6	6		
4	G	1	Total	C	O	0	0
			12	6	6		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	89	Total 89	O 89	0	0
5	C	79	Total 79	O 79	0	0
5	E	103	Total 103	O 103	0	0
5	G	97	Total 97	O 97	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NADPH-sorbose reductase





- Molecule 1: NADPH-sorbose reductase



- Molecule 1: NADPH-sorbose reductase



- Molecule 1: NADPH-sorbose reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.04Å 124.41Å 124.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.69 – 1.80 19.69 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (19.69-1.80) 99.9 (19.69-1.80)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.46 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.212 , 0.249 0.174 , 0.204	Depositor DCC
R_{free} test set	4432 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	17.2	Xtriage
Anisotropy	0.037	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 50.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8624	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 60.74 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.4551e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, SOE, SOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	0/2026	0.66	0/2743
1	C	0.63	0/2026	0.65	0/2743
1	E	0.61	0/2026	0.65	0/2743
1	G	0.63	0/2026	0.67	0/2743
All	All	0.63	0/8104	0.66	0/10972

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1989	0	2004	12	0
1	C	1989	0	2004	10	0
1	E	1989	0	2004	17	0
1	G	1989	0	2004	15	0
2	A	48	0	26	1	0
2	C	48	0	26	0	0
2	E	48	0	26	0	0
2	G	48	0	26	0	0
3	A	12	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	12	0	0	1	0
3	E	24	0	0	2	0
3	G	12	0	0	0	0
4	E	12	0	12	2	0
4	G	36	0	36	6	0
5	A	89	0	0	0	0
5	C	79	0	0	0	0
5	E	103	0	0	1	0
5	G	97	0	0	1	0
All	All	8624	0	8168	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:263:LEU:O	4:G:266:SOE:H1	1.11	1.25
1:G:263:LEU:O	4:G:266:SOE:C1	2.07	1.01
1:A:49:ARG:O	1:A:53:GLU:HG3	1.92	0.69
1:C:253:TYR:HD1	1:G:253:TYR:HD1	1.42	0.68
1:E:154:GLU:OE2	3:E:0:SOL:O3	2.11	0.67
1:A:49:ARG:HB3	1:A:49:ARG:HH11	1.60	0.67
1:E:243:GLU:HG2	4:E:264:SOE:O4	1.95	0.66
1:E:33:HIS:ND1	5:E:304:HOH:O	2.17	0.66
1:G:5:ILE:HG22	4:G:265:SOE:O5	1.99	0.62
1:G:45:HIS:HD2	1:G:61:GLU:OE1	1.88	0.56
1:A:253:TYR:HD1	1:E:253:TYR:HD1	1.54	0.55
1:G:240:LEU:HD11	1:G:253:TYR:CD2	2.42	0.55
1:A:46:GLU:OE2	1:A:49:ARG:NH1	2.40	0.54
1:A:206:ASN:HD22	1:A:206:ASN:C	2.10	0.53
1:A:46:GLU:CD	1:A:49:ARG:NH1	2.63	0.51
1:E:60:LEU:HD21	1:E:78:SER:OG	2.11	0.51
1:E:243:GLU:CG	4:E:264:SOE:O4	2.60	0.50
1:A:253:TYR:HD1	1:E:253:TYR:CD1	2.30	0.50
1:C:263:LEU:O	4:G:266:SOE:O5	2.27	0.49
1:E:206:ASN:HD22	1:E:206:ASN:C	2.15	0.49
1:E:240:LEU:HD11	1:E:253:TYR:HD2	1.78	0.48
1:C:206:ASN:HD22	1:C:206:ASN:C	2.17	0.48
1:E:240:LEU:HD11	1:E:253:TYR:CD2	2.48	0.48
1:G:143:ALA:HB3	1:G:186:ASN:OD1	2.13	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:263:LEU:HA	1:C:263:LEU:HD23	1.60	0.47
1:G:5:ILE:CD1	1:G:32:ALA:HB2	2.47	0.45
1:G:54:LYS:HB3	1:G:54:LYS:HE3	1.74	0.45
1:E:52:LYS:NZ	1:E:59:VAL:HB	2.33	0.44
1:C:96:GLY:CA	3:C:0:SOL:O4	2.65	0.44
1:A:263:LEU:HA	1:A:263:LEU:HD23	1.60	0.44
1:C:185:ILE:HD11	1:C:240:LEU:HD12	1.99	0.44
1:C:36:LEU:HB3	1:C:44:LEU:HD11	2.00	0.44
1:E:231:GLU:CD	1:E:231:GLU:H	2.21	0.44
1:G:252:ALA:O	4:G:266:SOE:O4	2.31	0.43
1:A:240:LEU:HD11	1:A:253:TYR:CD2	2.54	0.42
1:A:143:ALA:HB3	1:A:186:ASN:OD1	2.19	0.42
1:G:240:LEU:HD11	1:G:253:TYR:HD2	1.81	0.42
1:A:14:GLY:HA2	2:A:270:NDP:H1B	2.02	0.42
1:A:185:ILE:HD11	1:A:240:LEU:HD12	2.02	0.42
1:C:143:ALA:HB3	1:C:186:ASN:OD1	2.18	0.42
1:E:96:GLY:HA3	3:E:0:SOL:O4	2.19	0.42
1:E:193:PRO:O	1:E:197:LYS:HD3	2.19	0.42
1:G:243:GLU:HG2	4:G:265:SOE:O4	2.19	0.42
1:G:177:LYS:HB2	5:G:486:HOH:O	2.20	0.41
1:E:143:ALA:HB3	1:E:186:ASN:OD1	2.20	0.41
1:C:107:GLU:CD	1:C:107:GLU:H	2.24	0.40
1:C:262:THR:HG22	1:E:149:GLN:HB2	2.03	0.40
1:E:11:VAL:HA	1:E:35:VAL:HB	2.03	0.40
1:G:142:ASN:HD22	1:G:142:ASN:C	2.23	0.40
1:G:38:ALA:O	1:G:63:ALA:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/263 (99%)	254 (97%)	6 (2%)	1 (0%)	38	23
1	C	261/263 (99%)	253 (97%)	8 (3%)	0	100	100
1	E	261/263 (99%)	252 (97%)	9 (3%)	0	100	100
1	G	261/263 (99%)	251 (96%)	10 (4%)	0	100	100
All	All	1044/1052 (99%)	1010 (97%)	33 (3%)	1 (0%)	55	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	98	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/206 (100%)	201 (98%)	5 (2%)	54	40
1	C	206/206 (100%)	202 (98%)	4 (2%)	62	50
1	E	206/206 (100%)	202 (98%)	4 (2%)	62	50
1	G	206/206 (100%)	201 (98%)	5 (2%)	54	40
All	All	824/824 (100%)	806 (98%)	18 (2%)	57	44

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	ARG
1	A	142	ASN
1	A	206	ASN
1	A	232	GLU
1	A	236	PHE
1	C	142	ASN
1	C	206	ASN
1	C	220	GLU
1	C	236	PHE
1	E	44	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	142	ASN
1	E	197	LYS
1	E	206	ASN
1	G	43	ARG
1	G	52	LYS
1	G	142	ASN
1	G	177	LYS
1	G	232	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	92	ASN
1	A	142	ASN
1	A	206	ASN
1	C	92	ASN
1	C	142	ASN
1	C	206	ASN
1	E	40	GLN
1	E	92	ASN
1	E	142	ASN
1	E	206	ASN
1	E	215	GLN
1	G	45	HIS
1	G	92	ASN
1	G	142	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	SOL	A	0	-	11,11,11	0.59	0	10,14,14	1.02	1 (10%)
2	NDP	A	270	-	43,52,52	1.50	5 (11%)	49,80,80	1.95	3 (6%)
3	SOL	C	0	-	11,11,11	0.74	0	10,14,14	1.55	1 (10%)
2	NDP	C	270	-	43,52,52	1.57	5 (11%)	49,80,80	1.85	4 (8%)
3	SOL	E	0	-	11,11,11	0.40	0	10,14,14	1.58	3 (30%)
4	SOE	E	264	-	12,12,12	1.03	1 (8%)	17,18,18	0.98	1 (5%)
3	SOL	E	265	-	11,11,11	0.53	0	10,14,14	1.26	1 (10%)
2	NDP	E	270	-	43,52,52	1.63	6 (13%)	49,80,80	1.87	4 (8%)
3	SOL	G	0	-	11,11,11	0.40	0	10,14,14	0.85	0
4	SOE	G	264	-	12,12,12	0.88	1 (8%)	17,18,18	0.98	1 (5%)
4	SOE	G	265	-	12,12,12	1.06	1 (8%)	17,18,18	1.35	2 (11%)
4	SOE	G	266	-	12,12,12	1.29	2 (16%)	17,18,18	3.15	11 (64%)
2	NDP	G	270	-	43,52,52	1.53	4 (9%)	49,80,80	1.81	4 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SOL	A	0	-	-	0/16/16/16	0/0/0/0
2	NDP	A	270	-	-	0/30/77/77	0/5/5/5
3	SOL	C	0	-	-	0/16/16/16	0/0/0/0
2	NDP	C	270	-	-	0/30/77/77	0/5/5/5
3	SOL	E	0	-	-	0/16/16/16	0/0/0/0
4	SOE	E	264	-	-	0/3/23/23	0/1/1/1
3	SOL	E	265	-	-	1/16/16/16	0/0/0/0
2	NDP	E	270	-	-	0/30/77/77	0/5/5/5
3	SOL	G	0	-	-	1/16/16/16	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SOE	G	264	-	-	0/3/23/23	0/1/1/1
4	SOE	G	265	-	-	0/3/23/23	0/1/1/1
4	SOE	G	266	-	-	0/3/23/23	0/1/1/1
2	NDP	G	270	-	-	0/30/77/77	0/5/5/5

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	270	NDP	C4N-C5N	-4.15	1.40	1.49
2	A	270	NDP	C4N-C5N	-4.09	1.40	1.49
2	G	270	NDP	C4N-C5N	-3.98	1.40	1.49
2	C	270	NDP	C4N-C5N	-3.65	1.41	1.49
4	G	266	SOE	C4-C3	-2.19	1.49	1.53
2	E	270	NDP	P2B-O2B	2.01	1.63	1.59
2	C	270	NDP	C2A-N1A	2.03	1.37	1.33
2	A	270	NDP	C2A-N1A	2.18	1.38	1.33
4	G	266	SOE	O6-C2	2.20	1.44	1.42
4	G	264	SOE	O2-C2	2.24	1.44	1.40
2	E	270	NDP	C2A-N1A	2.42	1.38	1.33
4	E	264	SOE	O6-C2	2.59	1.45	1.42
4	G	265	SOE	O6-C2	2.64	1.45	1.42
2	G	270	NDP	C2A-N3A	3.22	1.37	1.32
2	A	270	NDP	C6N-C5N	3.30	1.39	1.33
2	E	270	NDP	C2A-N3A	3.37	1.37	1.32
2	E	270	NDP	C6N-C5N	3.45	1.39	1.33
2	G	270	NDP	C6N-C5N	3.49	1.39	1.33
2	C	270	NDP	C2A-N3A	3.76	1.38	1.32
2	C	270	NDP	C6N-C5N	4.04	1.40	1.33
2	A	270	NDP	C2A-N3A	4.08	1.39	1.32
2	A	270	NDP	O7N-C7N	5.14	1.37	1.24
2	C	270	NDP	O7N-C7N	5.61	1.38	1.24
2	G	270	NDP	O7N-C7N	5.95	1.39	1.24
2	E	270	NDP	O7N-C7N	6.49	1.40	1.24

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	270	NDP	N3A-C2A-N1A	-11.33	118.99	128.86
2	C	270	NDP	N3A-C2A-N1A	-10.97	119.31	128.86
2	G	270	NDP	N3A-C2A-N1A	-10.87	119.39	128.86
2	E	270	NDP	N3A-C2A-N1A	-10.86	119.40	128.86
4	G	266	SOE	C1-C2-C3	-5.27	100.76	111.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	266	SOE	O5-C5-C4	-4.58	101.17	110.17
4	G	266	SOE	O4-C4-C3	-3.85	101.53	109.65
4	G	266	SOE	O1-C1-C2	-3.59	104.04	111.81
3	C	0	SOL	O3-C3-C4	-3.47	103.09	110.41
2	A	270	NDP	C3N-C2N-N1N	-3.44	118.08	123.08
2	C	270	NDP	C3N-C2N-N1N	-2.94	118.81	123.08
2	G	270	NDP	C3N-C2N-N1N	-2.92	118.84	123.08
4	E	264	SOE	O2-C2-C1	-2.85	106.69	111.20
3	E	0	SOL	O3-C3-C2	-2.83	106.02	110.95
4	G	266	SOE	C6-O6-C2	-2.80	110.90	114.09
4	G	266	SOE	O3-C3-C4	-2.67	104.06	110.07
3	E	0	SOL	O4-C4-C3	-2.61	104.38	109.20
3	A	0	SOL	O3-C3-C4	-2.59	104.94	110.41
2	E	270	NDP	C3N-C2N-N1N	-2.44	119.54	123.08
2	C	270	NDP	O4D-C1D-C2D	-2.22	101.74	106.64
2	G	270	NDP	C1B-N9A-C4A	-2.15	122.92	126.64
3	E	265	SOL	O3-C3-C4	-2.12	105.94	110.41
2	E	270	NDP	O4D-C1D-C2D	-2.09	102.03	106.64
2	C	270	NDP	C4A-C5A-N7A	-2.08	107.40	109.41
3	E	0	SOL	O1-C1-C2	-2.01	107.25	112.66
4	G	265	SOE	O2-C2-C1	-2.00	108.04	111.20
2	G	270	NDP	O3X-P2B-O2X	2.27	116.76	107.61
2	E	270	NDP	O2N-PN-O1N	2.38	124.58	112.28
4	G	266	SOE	O6-C6-C5	2.42	114.81	111.09
2	A	270	NDP	O2N-PN-O1N	2.61	125.77	112.28
4	G	264	SOE	C6-O6-C2	2.76	117.24	114.09
4	G	265	SOE	O3-C3-C2	2.88	114.42	110.02
4	G	266	SOE	C5-C4-C3	3.22	114.01	110.49
4	G	266	SOE	C2-C3-C4	3.26	115.23	110.76
4	G	266	SOE	O2-C2-C3	3.79	114.03	107.74
4	G	266	SOE	C6-C5-C4	5.47	116.58	109.65

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	265	SOL	O2-C2-C1-O1
3	G	0	SOL	O2-C2-C1-O1

There are no ring outliers.

6 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	270	NDP	1	0
3	C	0	SOL	1	0
3	E	0	SOL	2	0
4	E	264	SOE	2	0
4	G	265	SOE	2	0
4	G	266	SOE	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	263/263 (100%)	-0.27	2 (0%) 86 84	5, 10, 20, 25	0
1	C	263/263 (100%)	-0.20	0 100 100	5, 11, 22, 31	0
1	E	263/263 (100%)	-0.19	4 (1%) 74 70	5, 11, 26, 33	0
1	G	263/263 (100%)	-0.22	5 (1%) 67 63	5, 10, 23, 32	0
All	All	1052/1052 (100%)	-0.22	11 (1%) 82 80	5, 11, 22, 33	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	2	ASP	3.0
1	E	204	LYS	3.0
1	E	205	ASP	2.9
1	E	201	GLU	2.7
1	G	197	LYS	2.4
1	E	208	GLY	2.2
1	A	53	GLU	2.1
1	A	40	GLN	2.1
1	G	206	ASN	2.1
1	G	43	ARG	2.0
1	G	201	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	SOL	A	0	12/12	0.64	0.32	11.49	20,36,38,38	0
3	SOL	E	0	12/12	0.74	0.28	7.92	26,36,38,38	0
3	SOL	C	0	12/12	0.71	0.27	7.46	28,36,38,39	0
3	SOL	G	0	12/12	0.65	0.30	5.97	28,40,42,43	0
4	SOE	G	266	12/12	0.69	0.29	5.58	30,32,34,35	0
3	SOL	E	265	12/12	0.64	0.20	3.62	25,31,34,34	0
4	SOE	E	264	12/12	0.91	0.15	1.51	19,20,22,23	0
4	SOE	G	265	12/12	0.91	0.14	1.30	20,22,24,25	0
4	SOE	G	264	12/12	0.95	0.12	0.69	17,19,20,22	0
2	NDP	A	270	48/48	0.96	0.08	-0.45	6,12,17,18	0
2	NDP	E	270	48/48	0.97	0.07	-0.78	8,11,14,15	0
2	NDP	G	270	48/48	0.97	0.07	-0.82	6,12,15,16	0
2	NDP	C	270	48/48	0.97	0.07	-1.04	6,11,14,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.