



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 01:08 am GMT

PDB ID : 2AJL  
Title : X-ray Structure of Novel Biaryl-Based Dipeptidyl peptidase IV inhibitor  
Authors : Qiao, L.; Baumann, C.A.; Crysler, C.S.; Ninan, N.S.; Abad, M.C.; Spurlino, J.C.; DesJarlais, R.L.; Kervinen, J.; Neeper, M.P.; Bayoumy, S.S.  
Deposited on : 2005-08-02  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)
RSRZ outliers	101464	3876 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	I	728	
1	J	728	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	I	768	X	-	-	-
2	NAG	I	771	X	-	-	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	J	767	-	-	X	-
2	NAG	J	769	X	-	-	-
2	NAG	J	770	-	-	-	X
2	NAG	J	771	X	-	-	-
3	JNH	I	1	-	-	-	X
3	JNH	J	1	X	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 12558 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dipeptidyl peptidase 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	I	726	Total	C	N	O	S	0	0	0
			5947	3818	977	1126	26			
1	J	728	Total	C	N	O	S	0	0	0
			5964	3827	982	1129	26			

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	I	1	Total	C	N	O	0	0
			14	8	1	5		
2	I	1	Total	C	N	O	0	0
			14	8	1	5		
2	I	1	Total	C	N	O	0	0
			14	8	1	5		
2	I	1	Total	C	N	O	0	0
			14	8	1	5		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	I	1	Total	C	N	O	0	0
			14	8	1	5		
2	J	1	Total	C	N	O	0	0
			14	8	1	5		
2	J	1	Total	C	N	O	0	0
			14	8	1	5		
2	J	1	Total	C	N	O	0	0
			14	8	1	5		
2	J	1	Total	C	N	O	0	0
			14	8	1	5		
2	J	1	Total	C	N	O	0	0
			14	8	1	5		
2	J	1	Total	C	N	O	0	0
			14	8	1	5		
2	J	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is 1-[2-(S)-AMINO-3-BIPHENYL-4-YL-PROPIONYL]-PYRROLIDINE-2-(S)-CARBONITRILE (three-letter code: JNH) (formula: C<sub>20</sub>H<sub>25</sub>N<sub>3</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	I	1	Total	C	N	O	0	0
			24	20	3	1		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	J	1	Total	C	N	O	0	0
			24	20	3	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	I	190	Total	O	0	0
			190	190		
4	J	213	Total	O	0	0
			213	213		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Dipeptidyl peptidase 4



● Molecule 1: Dipeptidyl peptidase 4





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.28Å 126.88Å 110.83Å 90.00° 99.41° 90.00°	Depositor
Resolution (Å)	64.10 – 2.50 64.40 – 2.51	Depositor EDS
% Data completeness (in resolution range)	90.8 (64.10-2.50) 91.6 (64.40-2.51)	Depositor EDS
$R_{merge}$	0.37	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.76 (at 2.51Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.226 , 0.303 0.245 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	20.7	Xtriage
Anisotropy	0.345	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 48.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.53$ , $\langle L^2 \rangle = 0.37$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	12558	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: JNH, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	I	0.36	0/6119	0.68	4/8322 (0.0%)
1	J	0.37	0/6136	0.67	4/8344 (0.0%)
All	All	0.36	0/12255	0.68	8/16666 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	J	0	1

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	I	766	PRO	CA-C-O	9.57	143.17	120.20
1	J	766	PRO	CA-C-O	9.20	142.28	120.20
1	I	388	GLN	N-CA-C	-6.01	94.77	111.00
1	I	300	LEU	N-CA-C	-5.72	95.55	111.00
1	J	240	PHE	N-CA-C	-5.27	96.78	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	J	700	TYR	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	I	5947	0	5666	532	0
1	J	5964	0	5684	500	0
2	I	70	0	65	8	0
2	J	126	0	117	22	0
3	I	24	0	24	7	0
3	J	24	0	23	7	0
4	I	190	0	0	18	0
4	J	213	0	0	13	0
All	All	12558	0	11579	1029	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 43.

The worst 5 of 1029 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:489:LYS:HE3	1:J:491:LEU:HD21	1.27	1.14
1:I:289:ALA:HB1	1:I:290:PRO:HA	1.28	1.13
1:I:107:ILE:HG13	1:I:114:ILE:HG12	1.23	1.11
1:I:310:ARG:HD3	1:I:329:ASP:OD1	1.51	1.10
1:I:626:ILE:HD13	1:I:639:VAL:HG21	1.34	1.10

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	I	724/728 (100%)	607 (84%)	86 (12%)	31 (4%)	3	3
1	J	726/728 (100%)	632 (87%)	82 (11%)	12 (2%)	11	18
All	All	1450/1456 (100%)	1239 (85%)	168 (12%)	43 (3%)	5	7

5 of 43 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	I	94	THR
1	I	98	PHE
1	I	99	GLY
1	I	140	ARG
1	I	147	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	I	651/653 (100%)	603 (93%)	48 (7%)	16	30
1	J	653/653 (100%)	605 (93%)	48 (7%)	16	30
All	All	1304/1306 (100%)	1208 (93%)	96 (7%)	16	30

5 of 96 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	685	ASN
1	J	137	LEU
1	J	673	LEU
1	I	689	MET
1	J	45	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 58 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	621	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	I	731	GLN
1	J	562	ASN
1	I	679	ASN
1	I	694	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	JNH	I	1	1	26,26,26	1.94	6 (23%)	34,35,35	1.46	7 (20%)
2	NAG	I	767	-	14,14,15	2.76	3 (21%)	15,19,21	2.35	4 (26%)
2	NAG	I	768	-	14,14,15	2.57	3 (21%)	15,19,21	3.54	4 (26%)
2	NAG	I	769	-	14,14,15	2.41	2 (14%)	15,19,21	2.14	4 (26%)
2	NAG	I	770	-	14,14,15	2.56	2 (14%)	15,19,21	2.58	6 (40%)
2	NAG	I	771	-	14,14,15	2.57	2 (14%)	15,19,21	1.77	5 (33%)
3	JNH	J	1	1	26,26,26	1.77	6 (23%)	34,35,35	1.88	8 (23%)
2	NAG	J	767	-	14,14,15	2.58	3 (21%)	15,19,21	3.19	8 (53%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	J	768	-	14,14,15	2.64	2 (14%)	15,19,21	2.99	5 (33%)
2	NAG	J	769	-	14,14,15	2.60	2 (14%)	15,19,21	2.60	6 (40%)
2	NAG	J	770	-	14,14,15	2.84	4 (28%)	15,19,21	2.54	7 (46%)
2	NAG	J	771	-	14,14,15	2.98	4 (28%)	15,19,21	5.99	7 (46%)
2	NAG	J	772	-	14,14,15	2.61	2 (14%)	15,19,21	3.54	6 (40%)
2	NAG	J	773	-	14,14,15	2.81	3 (21%)	15,19,21	2.48	5 (33%)
2	NAG	J	774	-	14,14,15	2.63	3 (21%)	15,19,21	3.23	7 (46%)
2	NAG	J	775	-	14,14,15	2.65	3 (21%)	15,19,21	2.93	8 (53%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	JNH	I	1	1	-	0/18/28/28	0/3/3/3
2	NAG	I	767	-	-	1/6/23/26	0/1/1/1
2	NAG	I	768	-	2/2/5/7	0/6/23/26	0/1/1/1
2	NAG	I	769	-	-	1/6/23/26	0/1/1/1
2	NAG	I	770	-	-	0/6/23/26	0/1/1/1
2	NAG	I	771	-	1/1/5/7	0/6/23/26	0/1/1/1
3	JNH	J	1	1	1/1/4/4	0/18/28/28	0/3/3/3
2	NAG	J	767	-	-	0/6/23/26	0/1/1/1
2	NAG	J	768	-	-	0/6/23/26	0/1/1/1
2	NAG	J	769	-	1/1/5/7	0/6/23/26	0/1/1/1
2	NAG	J	770	-	-	0/6/23/26	0/1/1/1
2	NAG	J	771	-	1/1/5/7	0/6/23/26	0/1/1/1
2	NAG	J	772	-	-	0/6/23/26	0/1/1/1
2	NAG	J	773	-	-	0/6/23/26	0/1/1/1
2	NAG	J	774	-	-	0/6/23/26	0/1/1/1
2	NAG	J	775	-	-	1/6/23/26	0/1/1/1

The worst 5 of 50 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	1	JNH	C20-N3	-7.08	1.29	1.47
3	J	1	JNH	C20-N3	-5.85	1.32	1.47
2	J	771	NAG	C1-C2	-3.57	1.47	1.52
3	I	1	JNH	C4-N1	-2.48	1.42	1.47
3	J	1	JNH	C17-C8	-2.48	1.42	1.49

The worst 5 of 97 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	771	NAG	C2-N2-C7	-17.90	96.83	122.94
2	I	768	NAG	C2-N2-C7	-11.73	105.83	122.94
2	J	771	NAG	O5-C1-C2	-9.84	97.78	111.47
2	J	771	NAG	C1-O5-C5	-9.51	99.06	112.17
2	J	768	NAG	C2-N2-C7	-8.87	110.00	122.94

5 of 6 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	I	768	NAG	C5
2	I	768	NAG	C3
2	I	771	NAG	C5
2	J	769	NAG	C4
2	J	771	NAG	C3

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	I	769	NAG	O7-C7-N2-C2
2	I	767	NAG	O7-C7-N2-C2
2	J	775	NAG	C8-C7-N2-C2

There are no ring outliers.

12 monomers are involved in 44 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	1	JNH	7	0
2	I	767	NAG	2	0
2	I	768	NAG	5	0
2	I	770	NAG	1	0
3	J	1	JNH	7	0
2	J	767	NAG	7	0
2	J	768	NAG	1	0
2	J	769	NAG	3	0
2	J	770	NAG	1	0
2	J	771	NAG	6	0
2	J	773	NAG	1	0
2	J	774	NAG	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	I	726/728 (99%)	0.33	29 (3%) 39 41	3, 19, 50, 80	0
1	J	728/728 (100%)	0.31	30 (4%) 38 40	3, 16, 49, 86	0
All	All	1454/1456 (99%)	0.32	59 (4%) 38 40	3, 18, 49, 86	0

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	99	GLY	7.5
1	J	98	PHE	5.0
1	I	98	PHE	4.7
1	J	277	SER	4.6
1	J	389	ILE	4.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	J	770	14/15	0.65	0.35	6.29	56,62,69,70	0
3	JNH	I	1	24/24	0.86	0.27	3.45	27,32,38,41	0
3	JNH	J	1	24/24	0.86	0.22	2.44	19,24,30,30	0
2	NAG	I	768	14/15	0.83	0.28	1.92	35,40,42,43	0
2	NAG	J	771	14/15	0.81	0.22	1.90	18,21,30,31	0
2	NAG	J	774	14/15	0.83	0.20	1.41	35,38,41,42	0
2	NAG	I	767	14/15	0.74	0.21	0.74	38,40,42,43	0
2	NAG	J	767	14/15	0.78	0.24	-0.22	39,46,49,50	0
2	NAG	J	775	14/15	0.76	0.23	-	48,57,60,64	0
2	NAG	J	768	14/15	0.57	0.28	-	67,70,72,72	0
2	NAG	J	772	14/15	0.77	0.25	-	56,60,65,65	0
2	NAG	I	770	14/15	0.74	0.28	-	77,80,81,82	0
2	NAG	J	769	14/15	0.75	0.24	-	45,50,53,55	0
2	NAG	I	769	14/15	0.70	0.27	-	59,61,65,66	0
2	NAG	I	771	14/15	0.61	0.29	-	78,82,86,86	0
2	NAG	J	773	14/15	0.78	0.18	-	31,38,41,45	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.