



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 06:39 pm GMT

PDB ID : 5AJK  
Title : Crystal structure of variola virus virulence factor F1L in complex with human Bak BH3 domain  
Authors : Kvensakul, M.; Colman, P.M.  
Deposited on : 2015-02-25  
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4993 (2.58-2.50)
Clashscore	112137	5755 (2.58-2.50)
Ramachandran outliers	110173	5652 (2.58-2.50)
Sidechain outliers	110143	5654 (2.58-2.50)
RSRZ outliers	101464	5026 (2.58-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	168	<div> <div>2%</div> <div>85%</div> <div>12%</div> </div>
1	C	168	<div> <div>2%</div> <div>82%</div> <div>17%</div> </div>
1	E	168	<div> <div>6%</div> <div>80%</div> <div>16%</div> </div>
1	G	168	<div> <div>4%</div> <div>79%</div> <div>18%</div> </div>
1	I	168	<div> <div>17%</div> <div>79%</div> <div>18%</div> </div>
1	K	168	<div> <div>7%</div> <div>79%</div> <div>18%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	B	26	<div><div></div><div>4%</div><div>81%</div><div>8%</div><div>12%</div></div>
2	D	26	<div><div></div><div>8%</div><div>88%</div><div></div><div>12%</div></div>
2	F	26	<div><div></div><div>4%</div><div>85%</div><div></div><div>12%</div></div>
2	H	26	<div><div></div><div></div><div>77%</div><div>12%</div><div>12%</div></div>
2	J	26	<div><div></div><div>38%</div><div>92%</div><div></div><div>8%</div></div>
2	L	26	<div><div></div><div>8%</div><div>85%</div><div></div><div>15%</div></div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 15824 atoms, of which 7769 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HOMOLOG OF VACCINIA VIRUS CDS F1L.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	148	Total	C	H	N	O	S	0	0	0
			2386	763	1176	196	237	14			
1	C	139	Total	C	H	N	O	S	0	0	0
			2244	711	1110	184	225	14			
1	E	141	Total	C	H	N	O	S	0	0	0
			2290	732	1131	186	227	14			
1	G	137	Total	C	H	N	O	S	0	0	0
			2222	705	1100	182	221	14			
1	I	137	Total	C	H	N	O	S	0	0	0
			2219	704	1098	182	221	14			
1	K	138	Total	C	H	N	O	S	0	0	0
			2233	708	1105	183	223	14			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	GLY	-	EXPRESSION TAG	UNP Q85365
A	35	PRO	-	EXPRESSION TAG	UNP Q85365
A	36	LEU	-	EXPRESSION TAG	UNP Q85365
A	37	GLY	-	EXPRESSION TAG	UNP Q85365
A	38	SER	-	EXPRESSION TAG	UNP Q85365
C	34	GLY	-	EXPRESSION TAG	UNP Q85365
C	35	PRO	-	EXPRESSION TAG	UNP Q85365
C	36	LEU	-	EXPRESSION TAG	UNP Q85365
C	37	GLY	-	EXPRESSION TAG	UNP Q85365
C	38	SER	-	EXPRESSION TAG	UNP Q85365
E	34	GLY	-	EXPRESSION TAG	UNP Q85365
E	35	PRO	-	EXPRESSION TAG	UNP Q85365
E	36	LEU	-	EXPRESSION TAG	UNP Q85365
E	37	GLY	-	EXPRESSION TAG	UNP Q85365
E	38	SER	-	EXPRESSION TAG	UNP Q85365
G	34	GLY	-	EXPRESSION TAG	UNP Q85365
G	35	PRO	-	EXPRESSION TAG	UNP Q85365

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
G	36	LEU	-	EXPRESSION TAG	UNP Q85365
G	37	GLY	-	EXPRESSION TAG	UNP Q85365
G	38	SER	-	EXPRESSION TAG	UNP Q85365
I	34	GLY	-	EXPRESSION TAG	UNP Q85365
I	35	PRO	-	EXPRESSION TAG	UNP Q85365
I	36	LEU	-	EXPRESSION TAG	UNP Q85365
I	37	GLY	-	EXPRESSION TAG	UNP Q85365
I	38	SER	-	EXPRESSION TAG	UNP Q85365
K	34	GLY	-	EXPRESSION TAG	UNP Q85365
K	35	PRO	-	EXPRESSION TAG	UNP Q85365
K	36	LEU	-	EXPRESSION TAG	UNP Q85365
K	37	GLY	-	EXPRESSION TAG	UNP Q85365
K	38	SER	-	EXPRESSION TAG	UNP Q85365

- Molecule 2 is a protein called BCL-2 HOMOLOGOUS ANTAGONIST/KILLER.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	23	Total	C	H	N	O	S	0	1	0
			360	111	174	38	36	1			
2	D	23	Total	C	H	N	O	S	0	0	0
			341	100	170	35	35	1			
2	F	23	Total	C	H	N	O	S	0	0	0
			352	106	174	35	36	1			
2	H	23	Total	C	H	N	O	S	0	0	0
			352	106	174	35	36	1			
2	J	24	Total	C	H	N	O	S	0	0	0
			363	109	179	36	38	1			
2	L	22	Total	C	H	N	O	S	0	0	0
			341	103	169	34	34	1			

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Cl	0	0
			2	2		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	1	Total	C	H	O	0	0
			7	2	3	2		
4	E	1	Total	C	H	O	0	0
			7	2	3	2		
4	G	1	Total	C	H	O	0	0
			7	2	3	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	30	Total	O	0	0
			30	30		
5	B	5	Total	O	0	0
			5	5		
5	C	17	Total	O	0	0
			17	17		
5	D	2	Total	O	0	0
			2	2		
5	E	10	Total	O	0	0
			10	10		
5	F	5	Total	O	0	0
			5	5		
5	G	12	Total	O	0	0
			12	12		
5	H	2	Total	O	0	0
			2	2		
5	I	7	Total	O	0	0
			7	7		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	J	3	Total 3	O 3	0	0
5	K	3	Total 3	O 3	0	0
5	L	2	Total 2	O 2	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HOMOLOG OF VACCINIA VIRUS CDS F1L




- Molecule 1: HOMOLOG OF VACCINIA VIRUS CDS F1L




- Molecule 1: HOMOLOG OF VACCINIA VIRUS CDS F1L



- Molecule 1: HOMOLOG OF VACCINIA VIRUS CDS F1L




- Molecule 1: HOMOLOG OF VACCINIA VIRUS CDS F1L







- Molecule 1: HOMOLOG OF VACCINIA VIRUS CDS F1L




- Molecule 2: BCL-2 HOMOLOGOUS ANTAGONIST/KILLER




- Molecule 2: BCL-2 HOMOLOGOUS ANTAGONIST/KILLER




- Molecule 2: BCL-2 HOMOLOGOUS ANTAGONIST/KILLER




- Molecule 2: BCL-2 HOMOLOGOUS ANTAGONIST/KILLER



- Molecule 2: BCL-2 HOMOLOGOUS ANTAGONIST/KILLER



- Molecule 2: BCL-2 HOMOLOGOUS ANTAGONIST/KILLER



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	124.77Å 68.76Å 171.60Å 90.00° 109.43° 90.00°	Depositor
Resolution (Å)	39.70 – 2.55 39.70 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.7 (39.70-2.55) 92.9 (39.70-2.55)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.43 (at 2.54Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.197 , 0.240 0.202 , 0.240	Depositor DCC
$R_{free}$ test set	2299 reflections (5.79%)	DCC
Wilson B-factor (Å <sup>2</sup> )	49.8	Xtriage
Anisotropy	0.592	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 49.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	15824	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/1231	0.38	0/1667
1	C	0.23	0/1151	0.38	0/1556
1	E	0.23	0/1178	0.38	0/1593
1	G	0.23	0/1139	0.38	0/1540
1	I	0.23	0/1138	0.38	0/1538
1	K	0.24	0/1145	0.41	0/1548
2	B	0.21	0/189	0.42	0/251
2	D	0.22	0/170	0.43	0/226
2	F	0.24	0/178	0.46	0/237
2	H	0.24	0/178	0.40	0/237
2	J	0.22	0/184	0.41	0/245
2	L	0.22	0/172	0.41	0/229
All	All	0.23	0/8053	0.39	0/10867

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1210	1176	1176	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1134	1110	1110	0	0
1	E	1159	1131	1131	3	0
1	G	1122	1100	1100	3	0
1	I	1121	1098	1098	2	0
1	K	1128	1105	1105	4	0
2	B	186	174	187	2	0
2	D	171	170	167	0	0
2	F	178	174	174	1	0
2	H	178	174	174	2	0
2	J	184	179	179	0	0
2	L	172	169	169	0	0
3	A	2	0	0	0	0
4	C	4	3	3	0	0
4	E	4	3	3	0	0
4	G	4	3	3	0	0
5	A	30	0	0	0	0
5	B	5	0	0	0	0
5	C	17	0	0	0	0
5	D	2	0	0	0	0
5	E	10	0	0	0	0
5	F	5	0	0	0	0
5	G	12	0	0	0	0
5	H	2	0	0	0	0
5	I	7	0	0	0	0
5	J	3	0	0	0	0
5	K	3	0	0	0	0
5	L	2	0	0	0	0
All	All	8055	7769	7779	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:178:ILE:HG21	1:I:178:ILE:HG21	1.95	0.48
1:A:147:ILE:HD11	2:B:75:GLY:CA	2.44	0.47
1:E:140:ILE:HA	2:F:71:MET:HE3	1.97	0.47
1:G:118:ASP:OD2	2:H:88:ARG:NH2	2.48	0.47
1:E:177:THR:HG21	1:K:89:ARG:HD2	1.96	0.46
1:K:124:ASN:OD1	1:K:170:ARG:NH2	2.49	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:MET:O	2:B:70:THR:OG1	2.34	0.44
1:E:168:GLY:O	1:K:89:ARG:NH1	2.50	0.44
1:G:140:ILE:HD12	2:H:71:MET:HE1	2.00	0.43
1:I:66:ILE:HG23	1:I:67:LEU:HD22	2.01	0.41
1:K:144:VAL:O	1:K:148:LEU:HD12	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	144/168 (86%)	141 (98%)	3 (2%)	0	100	100
1	C	137/168 (82%)	133 (97%)	4 (3%)	0	100	100
1	E	137/168 (82%)	135 (98%)	2 (2%)	0	100	100
1	G	135/168 (80%)	133 (98%)	2 (2%)	0	100	100
1	I	135/168 (80%)	133 (98%)	2 (2%)	0	100	100
1	K	136/168 (81%)	130 (96%)	6 (4%)	0	100	100
2	B	22/26 (85%)	22 (100%)	0	0	100	100
2	D	21/26 (81%)	21 (100%)	0	0	100	100
2	F	21/26 (81%)	21 (100%)	0	0	100	100
2	H	21/26 (81%)	21 (100%)	0	0	100	100
2	J	22/26 (85%)	22 (100%)	0	0	100	100
2	L	20/26 (77%)	20 (100%)	0	0	100	100
All	All	951/1164 (82%)	932 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	141/159 (89%)	138 (98%)	3 (2%)	59	81
1	C	133/159 (84%)	132 (99%)	1 (1%)	85	94
1	E	135/159 (85%)	132 (98%)	3 (2%)	57	80
1	G	131/159 (82%)	130 (99%)	1 (1%)	85	94
1	I	131/159 (82%)	129 (98%)	2 (2%)	70	87
1	K	132/159 (83%)	131 (99%)	1 (1%)	85	94
2	B	20/22 (91%)	20 (100%)	0	100	100
2	D	18/22 (82%)	18 (100%)	0	100	100
2	F	19/22 (86%)	19 (100%)	0	100	100
2	H	19/22 (86%)	18 (95%)	1 (5%)	26	45
2	J	20/22 (91%)	20 (100%)	0	100	100
2	L	18/22 (82%)	18 (100%)	0	100	100
All	All	917/1086 (84%)	905 (99%)	12 (1%)	73	89

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	89	ARG
1	A	99	THR
1	A	114	LYS
1	C	147	ILE
1	E	66	ILE
1	E	67	LEU
1	E	68	ASP
1	G	149	MET
2	H	89	TYR
1	I	123	MET
1	I	194	ILE
1	K	200	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no

such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	ACT	C	1202	-	1,3,3	1.38	0	0,3,3	0.00	-
4	ACT	E	1202	-	1,3,3	1.28	0	0,3,3	0.00	-
4	ACT	G	1201	-	1,3,3	1.35	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ACT	C	1202	-	-	0/0/0/0	0/0/0/0
4	ACT	E	1202	-	-	0/0/0/0	0/0/0/0
4	ACT	G	1201	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	148/168 (88%)	0.36	4 (2%)	55	59	37, 56, 98, 128	0
1	C	139/168 (82%)	0.42	4 (2%)	52	56	41, 57, 93, 112	0
1	E	141/168 (83%)	0.46	10 (7%)	17	18	42, 65, 107, 150	0
1	G	137/168 (81%)	0.63	7 (5%)	29	31	42, 63, 100, 132	0
1	I	137/168 (81%)	0.98	28 (20%)	1	1	47, 72, 130, 140	0
1	K	138/168 (82%)	0.73	11 (7%)	13	13	44, 77, 117, 157	0
2	B	23/26 (88%)	0.82	1 (4%)	36	39	46, 60, 106, 133	0
2	D	23/26 (88%)	0.33	2 (8%)	11	11	48, 67, 92, 114	0
2	F	23/26 (88%)	0.37	1 (4%)	36	39	49, 59, 101, 125	0
2	H	23/26 (88%)	0.29	0	100	100	50, 63, 110, 118	0
2	J	24/26 (92%)	1.61	10 (41%)	0	0	77, 89, 144, 159	0
2	L	22/26 (84%)	0.55	2 (9%)	10	10	65, 85, 117, 145	0
All	All	978/1164 (84%)	0.60	80 (8%)	12	13	37, 66, 115, 159	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	90	ASP	6.9
1	I	131	SER	5.9
1	I	138	TYR	5.4
1	G	65	ASN	5.2
2	J	70	THR	4.7
2	J	68	SER	4.4
1	I	132	THR	4.4
1	I	134	SER	4.3
2	J	71	MET	4.2
1	E	67	LEU	4.0
1	I	66	ILE	4.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	L	89	TYR	3.9
1	I	65	ASN	3.9
1	I	135	PHE	3.7
1	K	69	TYR	3.7
1	G	67	LEU	3.6
1	I	71	SER	3.5
2	J	72	GLY	3.5
2	J	69	SER	3.5
1	I	133	GLU	3.3
1	I	69	TYR	3.3
1	K	201	SER	3.3
1	G	64	THR	3.3
1	K	197	ILE	3.3
1	I	121	ARG	3.3
1	G	69	TYR	3.3
1	I	68	ASP	3.2
1	K	200	ILE	3.2
1	E	66	ILE	3.2
1	I	125	ILE	3.2
1	I	129	MET	3.1
2	J	73	GLN	3.1
1	I	200	ILE	3.1
2	J	74	VAL	3.1
1	G	132	THR	2.9
1	K	133	GLU	2.9
1	I	149	MET	2.8
1	I	86	SER	2.8
1	E	72	THR	2.8
1	A	65	ASN	2.8
2	L	90	ASP	2.8
1	I	172	MET	2.8
1	A	54	GLU	2.7
1	I	201	SER	2.7
1	C	63	SER	2.7
1	C	134	SER	2.7
1	K	110	MET	2.7
1	I	142	ASN	2.7
1	A	201	SER	2.7
1	G	68	ASP	2.6
2	D	68	SER	2.6
1	I	127	TYR	2.6
1	E	70	LEU	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	E	48	TYR	2.6
1	I	130	ALA	2.5
1	I	140	ILE	2.5
2	F	90	ASP	2.5
1	I	84	TYR	2.5
2	J	89	TYR	2.5
1	E	49	VAL	2.4
1	I	70	LEU	2.3
1	E	47	ASN	2.3
1	C	133	GLU	2.3
1	A	140	ILE	2.3
2	J	90	ASP	2.3
1	K	68	ASP	2.3
1	K	192	CYS	2.3
1	I	67	LEU	2.2
1	C	69	TYR	2.2
1	E	149	MET	2.2
1	K	191	ASP	2.2
1	K	196	TYR	2.2
1	E	71	SER	2.2
1	I	137	VAL	2.1
2	J	91	SER	2.1
1	G	150	ASP	2.1
1	E	50	TYR	2.1
1	I	72	THR	2.1
1	K	64	THR	2.0
2	D	90	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CL	A	1203	1/1	0.78	0.49	-	83,83,83,83	0
4	ACT	E	1202	4/4	0.82	0.19	-	71,71,86,86	0
4	ACT	G	1201	4/4	0.87	0.14	-	76,77,91,91	1
4	ACT	C	1202	4/4	0.93	0.09	-	72,73,87,87	0
3	CL	A	1202	1/1	0.82	0.62	-	83,83,83,83	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.