



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 07:05 am GMT

PDB ID : 2AKZ
Title : Fluoride Inhibition of Enolase: Crystal Structure of the Inhibitory Complex
Authors : Qin, J.; Chai, G.; Brewer, J.M.; Lovelace, L.L.
Deposited on : 2005-08-04
Resolution : 1.36 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	1063 (1.38-1.34)
Ramachandran outliers	110173	1048 (1.38-1.34)
Sidechain outliers	110143	1048 (1.38-1.34)
RSRZ outliers	101464	1025 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	439	<div> <div>%</div> <div>84% 12% ...</div> </div>
1	B	439	<div> <div>%</div> <div>88% 9% ..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	TRS	A	460	-	X	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7546 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma enolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	435	Total	C	N	O	S	0	6	0
			3295	2086	570	626	13			
1	B	432	Total	C	N	O	S	0	9	0
			3258	2062	564	619	13			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	434	HIS	-	EXPRESSION TAG	UNP P09104
A	435	HIS	-	EXPRESSION TAG	UNP P09104
A	436	HIS	-	EXPRESSION TAG	UNP P09104
A	437	HIS	-	EXPRESSION TAG	UNP P09104
A	438	HIS	-	EXPRESSION TAG	UNP P09104
A	439	HIS	-	EXPRESSION TAG	UNP P09104
B	1434	HIS	-	EXPRESSION TAG	UNP P09104
B	1435	HIS	-	EXPRESSION TAG	UNP P09104
B	1436	HIS	-	EXPRESSION TAG	UNP P09104
B	1437	HIS	-	EXPRESSION TAG	UNP P09104
B	1438	HIS	-	EXPRESSION TAG	UNP P09104
B	1439	HIS	-	EXPRESSION TAG	UNP P09104

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Mg	0	0
			2	2		
2	A	2	Total	Mg	0	0
			2	2		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is FLUORIDE ION (three-letter code: F) (formula: F).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	F	0	0
			2	2		
4	A	2	Total	F	0	0
			2	2		

- Molecule 5 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	496	Total	O	0	0
			496	496		
6	B	471	Total	O	0	0
			471	471		

- Molecule 1: Gamma enolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	109.01Å 118.52Å 67.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.36 9.97 – 1.36	Depositor EDS
% Data completeness (in resolution range)	88.9 (10.00-1.36) 88.9 (9.97-1.36)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.61 (at 1.36Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.110 , 0.144 0.120 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	10.3	Xtriage
Anisotropy	0.557	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 86.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	7546	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, MG, PO4, F

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.38	3/3386 (0.1%)	1.49	41/4584 (0.9%)
1	B	0.69	1/3366 (0.0%)	1.32	18/4559 (0.4%)
All	All	1.09	4/6752 (0.1%)	1.41	59/9143 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	126	GLU	CD-OE2	70.04	2.02	1.25
1	A	3	GLU	CD-OE2	8.29	1.34	1.25
1	B	1003	GLU	CD-OE2	6.22	1.32	1.25
1	A	402	ARG	CZ-NH1	-5.39	1.26	1.33

The worst 5 of 59 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	126	GLU	OE1-CD-OE2	-32.56	84.23	123.30
1	A	131[A]	ARG	NE-CZ-NH1	14.84	127.72	120.30
1	A	131[B]	ARG	NE-CZ-NH1	14.84	127.72	120.30
1	B	1402	ARG	NE-CZ-NH2	13.26	126.93	120.30
1	B	1399	ARG	NE-CZ-NH1	12.79	126.69	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	126	GLU	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3295	0	3239	31	0
1	B	3258	0	3185	21	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
5	A	8	0	12	0	0
6	A	496	0	0	8	0
6	B	471	0	0	8	0
All	All	7546	0	6436	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 51 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:GLU:CD	1:A:126:GLU:OE2	2.02	0.97
1:B:1130:TYR:OH	1:B:1415:GLU:OE1	1.82	0.97
1:B:1265:ASP:O	1:B:1268:ARG:HG2	1.85	0.76
1:A:201:LYS:HE2	6:A:680:HOH:O	1.98	0.63
1:B:1143:ILE:HG13	1:B:1143:ILE:O	2.02	0.59

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	439/439 (100%)	431 (98%)	7 (2%)	1 (0%)	51	19
1	B	439/439 (100%)	429 (98%)	9 (2%)	1 (0%)	51	19
All	All	878/878 (100%)	860 (98%)	16 (2%)	2 (0%)	51	19

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	399	ARG
1	B	1399	ARG

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	335/356 (94%)	325 (97%)	10 (3%)	46	11
1	B	330/356 (93%)	322 (98%)	8 (2%)	54	17
All	All	665/712 (93%)	647 (97%)	18 (3%)	53	15

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	344	ASN
1	A	370	HIS
1	B	1391	GLN
1	A	284	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	337	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	434	HIS
1	A	435	HIS
1	B	1165	GLN
1	A	337	ASN
1	B	1150	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 8 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PO4	A	442	2	4,4,4	2.77	3 (75%)	6,6,6	0.54	0
5	TRS	A	460	-	7,7,7	32.78	3 (42%)	9,9,9	8.11	6 (66%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PO4	B	1442	2	4,4,4	2.43	2 (50%)	6,6,6	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PO4	A	442	2	-	0/0/0/0	0/0/0/0
5	TRS	A	460	-	-	0/9/9/9	0/0/0/0
3	PO4	B	1442	2	-	0/0/0/0	0/0/0/0

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	442	PO4	P-O1	-4.11	1.41	1.50
3	B	1442	PO4	P-O2	-3.55	1.41	1.54
3	B	1442	PO4	P-O4	-2.64	1.45	1.54
3	A	442	PO4	P-O2	-2.58	1.45	1.54
3	A	442	PO4	P-O4	-2.35	1.46	1.54

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	460	TRS	O3-C3-C	-19.80	53.13	110.47
5	A	460	TRS	C3-C-C2	-11.03	79.80	111.06
5	A	460	TRS	O1-C1-C	-6.08	92.85	110.47
5	A	460	TRS	C2-C-N	2.70	113.47	107.73
5	A	460	TRS	C1-C-N	3.03	114.17	107.73

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	435/439 (99%)	-0.50	3 (0%) 87 89	9, 14, 29, 42	0
1	B	432/439 (98%)	-0.51	3 (0%) 87 89	9, 14, 28, 45	0
All	All	867/878 (98%)	-0.51	6 (0%) 87 89	9, 14, 29, 45	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	435	HIS	2.6
1	B	1432	VAL	2.3
1	A	418	ASP	2.3
1	A	420	ALA	2.2
1	B	1141	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	TRS	A	460	8/8	0.95	0.09	3.49	16,19,28,74	0
4	F	A	445	1/1	0.99	0.06	0.70	14,14,14,14	0
4	F	A	444	1/1	1.00	0.05	-0.07	10,10,10,10	0
4	F	B	1445	1/1	0.99	0.04	-1.42	13,13,13,13	0
3	PO4	B	1442	5/5	1.00	0.03	-1.87	8,9,10,10	0
3	PO4	A	442	5/5	1.00	0.03	-2.48	9,9,10,10	0
2	MG	A	440	1/1	1.00	0.02	-2.56	11,11,11,11	0
2	MG	B	1440	1/1	1.00	0.02	-2.62	9,9,9,9	0
4	F	B	1444	1/1	1.00	0.01	-3.15	9,9,9,9	0
2	MG	B	1441	1/1	1.00	0.02	-3.47	9,9,9,9	0
2	MG	A	441	1/1	1.00	0.01	-4.61	10,10,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.