



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 13, 2017 – 03:11 pm GMT

PDB ID : 4AL2
Title : peptide deformylase (Ni-form) with hydrosulfide
Authors : Palm, G.J.; Hinrichs, W.
Deposited on : 2012-02-29
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2542 (2.60-2.60)
Clashscore	112137	2895 (2.60-2.60)
Ramachandran outliers	110173	2848 (2.60-2.60)
Sidechain outliers	110143	2848 (2.60-2.60)
RSRZ outliers	101464	2550 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	186	<div> <div>%</div> <div> <div></div> <div>76%</div> <div>10%</div> <div>•</div> <div>13%</div> </div> </div>
1	B	186	<div> <div>2%</div> <div> <div></div> <div>71%</div> <div>18%</div> <div>•</div> <div>10%</div> </div> </div>
1	C	186	<div> <div>%</div> <div> <div></div> <div>69%</div> <div>17%</div> <div>•</div> <div>13%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3992 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PEPTIDE DEFORMYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	162	Total	C	N	O	S	0	0	0
			1296	814	228	248	6			
1	B	167	Total	C	N	O	S	0	0	0
			1340	841	240	253	6			
1	C	162	Total	C	N	O	S	0	0	0
			1296	814	228	248	6			

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	169	SER	-	EXPRESSION TAG	UNP P0A6K3
A	170	SER	-	EXPRESSION TAG	UNP P0A6K3
A	171	SER	-	EXPRESSION TAG	UNP P0A6K3
A	172	VAL	-	EXPRESSION TAG	UNP P0A6K3
A	173	ASP	-	EXPRESSION TAG	UNP P0A6K3
A	174	LYS	-	EXPRESSION TAG	UNP P0A6K3
A	175	LEU	-	EXPRESSION TAG	UNP P0A6K3
A	176	ALA	-	EXPRESSION TAG	UNP P0A6K3
A	177	ALA	-	EXPRESSION TAG	UNP P0A6K3
A	178	ALA	-	EXPRESSION TAG	UNP P0A6K3
A	179	LEU	-	EXPRESSION TAG	UNP P0A6K3
A	180	GLU	-	EXPRESSION TAG	UNP P0A6K3
A	181	HIS	-	EXPRESSION TAG	UNP P0A6K3
A	182	HIS	-	EXPRESSION TAG	UNP P0A6K3
A	183	HIS	-	EXPRESSION TAG	UNP P0A6K3
A	184	HIS	-	EXPRESSION TAG	UNP P0A6K3
A	185	HIS	-	EXPRESSION TAG	UNP P0A6K3
A	186	HIS	-	EXPRESSION TAG	UNP P0A6K3
B	169	SER	-	EXPRESSION TAG	UNP P0A6K3
B	170	SER	-	EXPRESSION TAG	UNP P0A6K3
B	171	SER	-	EXPRESSION TAG	UNP P0A6K3
B	172	VAL	-	EXPRESSION TAG	UNP P0A6K3
B	173	ASP	-	EXPRESSION TAG	UNP P0A6K3

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	174	LYS	-	EXPRESSION TAG	UNP P0A6K3
B	175	LEU	-	EXPRESSION TAG	UNP P0A6K3
B	176	ALA	-	EXPRESSION TAG	UNP P0A6K3
B	177	ALA	-	EXPRESSION TAG	UNP P0A6K3
B	178	ALA	-	EXPRESSION TAG	UNP P0A6K3
B	179	LEU	-	EXPRESSION TAG	UNP P0A6K3
B	180	GLU	-	EXPRESSION TAG	UNP P0A6K3
B	181	HIS	-	EXPRESSION TAG	UNP P0A6K3
B	182	HIS	-	EXPRESSION TAG	UNP P0A6K3
B	183	HIS	-	EXPRESSION TAG	UNP P0A6K3
B	184	HIS	-	EXPRESSION TAG	UNP P0A6K3
B	185	HIS	-	EXPRESSION TAG	UNP P0A6K3
B	186	HIS	-	EXPRESSION TAG	UNP P0A6K3
C	169	SER	-	EXPRESSION TAG	UNP P0A6K3
C	170	SER	-	EXPRESSION TAG	UNP P0A6K3
C	171	SER	-	EXPRESSION TAG	UNP P0A6K3
C	172	VAL	-	EXPRESSION TAG	UNP P0A6K3
C	173	ASP	-	EXPRESSION TAG	UNP P0A6K3
C	174	LYS	-	EXPRESSION TAG	UNP P0A6K3
C	175	LEU	-	EXPRESSION TAG	UNP P0A6K3
C	176	ALA	-	EXPRESSION TAG	UNP P0A6K3
C	177	ALA	-	EXPRESSION TAG	UNP P0A6K3
C	178	ALA	-	EXPRESSION TAG	UNP P0A6K3
C	179	LEU	-	EXPRESSION TAG	UNP P0A6K3
C	180	GLU	-	EXPRESSION TAG	UNP P0A6K3
C	181	HIS	-	EXPRESSION TAG	UNP P0A6K3
C	182	HIS	-	EXPRESSION TAG	UNP P0A6K3
C	183	HIS	-	EXPRESSION TAG	UNP P0A6K3
C	184	HIS	-	EXPRESSION TAG	UNP P0A6K3
C	185	HIS	-	EXPRESSION TAG	UNP P0A6K3
C	186	HIS	-	EXPRESSION TAG	UNP P0A6K3

- Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Ni 1 1	0	0
2	A	1	Total Ni 1 1	0	0
2	C	1	Total Ni 1 1	0	0

- Molecule 3 is HYDROSULFURIC ACID (three-letter code: H2S) (formula: H₂S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total S 1 1	0	0
3	B	1	Total S 1 1	0	0
3	C	1	Total S 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	21	Total O 21 21	0	0
4	B	22	Total O 22 22	0	0
4	C	11	Total O 11 11	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PEPTIDE DEFORMYLASE



• Molecule 1: PEPTIDE DEFORMYLASE



• Molecule 1: PEPTIDE DEFORMYLASE



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	142.21Å 63.22Å 118.17Å 90.00° 144.28° 90.00°	Depositor
Resolution (Å)	69.00 – 2.60 38.67 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.3 (69.00-2.60) 99.4 (38.67-2.60)	Depositor EDS
R_{merge}	0.22	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.206 , 0.298 0.212 , 0.303	Depositor DCC
R_{free} test set	975 reflections (5.43%)	DCC
Wilson B-factor (Å ²)	46.7	Xtriage
Anisotropy	0.287	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for h+2*l,-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3992	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, H2S

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/1311	0.71	0/1766
1	B	0.52	0/1355	0.73	0/1823
1	C	0.49	0/1311	0.69	0/1766
All	All	0.50	0/3977	0.71	0/5355

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	108	ILE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1296	0	1333	12	0
1	B	1340	0	1388	21	0
1	C	1296	0	1333	13	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	1	0
4	A	21	0	0	2	0
4	B	22	0	0	1	0
4	C	11	0	0	0	0
All	All	3992	0	4054	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 44 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:101:PRO:HD3	1:C:99:LEU:HD21	1.70	0.74
1:A:14:ARG:NH2	1:A:141:LEU:HD12	2.06	0.70
1:B:77:LEU:HA	1:B:108:ILE:HG22	1.75	0.66
1:A:14:ARG:CZ	1:A:141:LEU:HD12	2.28	0.64
1:B:88:GLU:O	1:B:97:ARG:HA	2.00	0.61

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	160/186 (86%)	155 (97%)	5 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	165/186 (89%)	151 (92%)	13 (8%)	1 (1%)	28	53
1	C	160/186 (86%)	148 (92%)	9 (6%)	3 (2%)	9	18
All	All	485/558 (87%)	454 (94%)	27 (6%)	4 (1%)	22	44

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	24	ASN
1	B	166	ALA
1	C	161	LEU
1	C	23	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	144/163 (88%)	136 (94%)	8 (6%)	25	48
1	B	148/163 (91%)	141 (95%)	7 (5%)	30	57
1	C	144/163 (88%)	134 (93%)	10 (7%)	18	36
All	All	436/489 (89%)	411 (94%)	25 (6%)	24	47

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	79	GLU
1	B	114	ASP
1	C	157	LYS
1	B	109	ARG
1	B	143	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	152	GLN
1	C	28	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic and 3 are modelled with single atom - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	162/186 (87%)	-0.21	1 (0%) 89 88	44, 59, 82, 112	0
1	B	167/186 (89%)	-0.17	3 (1%) 69 63	45, 57, 79, 119	0
1	C	162/186 (87%)	-0.01	2 (1%) 79 75	48, 71, 99, 133	0
All	All	491/558 (87%)	-0.13	6 (1%) 79 75	44, 61, 96, 133	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	164	LEU	3.3
1	B	166	ALA	3.1
1	B	167	ARG	2.8
1	C	153	ARG	2.5
1	C	158	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	H2S	A	1002	1/1	0.92	0.15	-0.48	70,70,70,70	0
3	H2S	B	1002	1/1	0.93	0.14	-0.94	78,78,78,78	0
3	H2S	C	1002	1/1	0.92	0.10	-2.53	80,80,80,80	0
2	NI	A	1001	1/1	0.99	0.07	-3.46	57,57,57,57	0
2	NI	C	1001	1/1	0.98	0.07	-3.63	61,61,61,61	0
2	NI	B	1001	1/1	0.99	0.07	-4.60	58,58,58,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.