



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 10:34 pm GMT

PDB ID : 4AM7
Title : ADP-BOUND C-TERMINAL DOMAIN OF ACTIN-RELATED PROTEIN
ARP8 FROM S. CEREVISIAE
Authors : Wuerges, J.; Saravanan, M.; Bose, D.; Cook, N.J.; Zhang, X.; Wigley, D.B.
Deposited on : 2012-03-07
Resolution : 3.25 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1852 (3.32-3.20)
Clashscore	112137	2036 (3.32-3.20)
Ramachandran outliers	110173	2000 (3.32-3.20)
Sidechain outliers	110143	1998 (3.32-3.20)
RSRZ outliers	101464	1861 (3.32-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	655	 5% 60% 27% 7% 5%
1	B	655	 6% 57% 31% 7% 5%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 10126 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACTIN-LIKE PROTEIN ARP8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	623	5036	3219	840	963	14	0	0	0
1	B	623	5036	3219	840	963	14	0	0	0

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	227	MET	-	EXPRESSION TAG	UNP Q12386
A	228	GLY	-	EXPRESSION TAG	UNP Q12386
A	229	SER	-	EXPRESSION TAG	UNP Q12386
A	230	SER	-	EXPRESSION TAG	UNP Q12386
A	231	HIS	-	EXPRESSION TAG	UNP Q12386
A	232	HIS	-	EXPRESSION TAG	UNP Q12386
A	233	HIS	-	EXPRESSION TAG	UNP Q12386
A	234	HIS	-	EXPRESSION TAG	UNP Q12386
A	235	HIS	-	EXPRESSION TAG	UNP Q12386
A	236	HIS	-	EXPRESSION TAG	UNP Q12386
A	237	SER	-	EXPRESSION TAG	UNP Q12386
A	238	SER	-	EXPRESSION TAG	UNP Q12386
A	239	GLY	-	EXPRESSION TAG	UNP Q12386
A	240	LEU	-	EXPRESSION TAG	UNP Q12386
A	241	VAL	-	EXPRESSION TAG	UNP Q12386
A	242	PRO	-	EXPRESSION TAG	UNP Q12386
A	243	ARG	-	EXPRESSION TAG	UNP Q12386
A	244	GLY	-	EXPRESSION TAG	UNP Q12386
A	245	SER	-	EXPRESSION TAG	UNP Q12386
A	246	HIS	-	EXPRESSION TAG	UNP Q12386
A	247	MET	-	EXPRESSION TAG	UNP Q12386
B	227	MET	-	EXPRESSION TAG	UNP Q12386
B	228	GLY	-	EXPRESSION TAG	UNP Q12386
B	229	SER	-	EXPRESSION TAG	UNP Q12386
B	230	SER	-	EXPRESSION TAG	UNP Q12386

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	231	HIS	-	EXPRESSION TAG	UNP Q12386
B	232	HIS	-	EXPRESSION TAG	UNP Q12386
B	233	HIS	-	EXPRESSION TAG	UNP Q12386
B	234	HIS	-	EXPRESSION TAG	UNP Q12386
B	235	HIS	-	EXPRESSION TAG	UNP Q12386
B	236	HIS	-	EXPRESSION TAG	UNP Q12386
B	237	SER	-	EXPRESSION TAG	UNP Q12386
B	238	SER	-	EXPRESSION TAG	UNP Q12386
B	239	GLY	-	EXPRESSION TAG	UNP Q12386
B	240	LEU	-	EXPRESSION TAG	UNP Q12386
B	241	VAL	-	EXPRESSION TAG	UNP Q12386
B	242	PRO	-	EXPRESSION TAG	UNP Q12386
B	243	ARG	-	EXPRESSION TAG	UNP Q12386
B	244	GLY	-	EXPRESSION TAG	UNP Q12386
B	245	SER	-	EXPRESSION TAG	UNP Q12386
B	246	HIS	-	EXPRESSION TAG	UNP Q12386
B	247	MET	-	EXPRESSION TAG	UNP Q12386

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ACTIN-LIKE PROTEIN ARP8



• Molecule 1: ACTIN-LIKE PROTEIN ARP8





4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	137.72Å 89.80Å 149.65Å 90.00° 114.37° 90.00°	Depositor
Resolution (Å)	19.81 – 3.25 19.81 – 3.25	Depositor EDS
% Data completeness (in resolution range)	96.5 (19.81-3.25) 96.5 (19.81-3.25)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.19 (at 3.22Å)	Xtrriage
Refinement program	REFMAC 5.5.0088	Depositor
R, R_{free}	0.219 , 0.284 0.215 , 0.270	Depositor DCC
R_{free} test set	1301 reflections (5.41%)	DCC
Wilson B-factor (Å ²)	98.8	Xtrriage
Anisotropy	0.046	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 64.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.024 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10126	wwPDB-VP
Average B, all atoms (Å ²)	121.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.65	0/5149	0.77	6/6982 (0.1%)
1	B	0.54	0/5149	0.70	4/6982 (0.1%)
All	All	0.60	0/10298	0.74	10/13964 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	342	LYS	N-CA-C	6.07	127.39	111.00
1	A	335	ARG	NE-CZ-NH1	5.77	123.18	120.30
1	A	350	HIS	N-CA-C	5.73	126.47	111.00
1	B	370	ASP	N-CA-C	5.58	126.06	111.00
1	A	858	LEU	CA-CB-CG	5.53	128.01	115.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	349	ALA	Peptide
1	A	688	LYS	Peptide
1	A	690	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5036	0	4985	159	0
1	B	5036	0	4985	138	0
2	A	27	0	12	4	0
2	B	27	0	12	1	0
All	All	10126	0	9994	296	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 296 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:335:ARG:HB3	1:A:339:ARG:HH12	1.06	1.18
1:A:335:ARG:HB3	1:A:339:ARG:NH1	1.73	1.03
1:A:315:LEU:HG	1:A:316:GLN:H	1.27	0.97
1:A:790:ASP:HB2	1:A:793:LYS:HB2	1.44	0.97
1:A:372:SER:HB3	1:A:373:PRO:HD2	1.48	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	621/655 (95%)	542 (87%)	76 (12%)	3 (0%)	32 71
1	B	621/655 (95%)	534 (86%)	83 (13%)	4 (1%)	28 68

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1242/1310 (95%)	1076 (87%)	159 (13%)	7 (1%)	28	68

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	346	PRO
1	B	371	PRO
1	A	372	SER
1	A	855	VAL
1	B	691	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	569/595 (96%)	473 (83%)	96 (17%)	2	12
1	B	569/595 (96%)	462 (81%)	107 (19%)	2	9
All	All	1138/1190 (96%)	935 (82%)	203 (18%)	2	10

5 of 203 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	853	LYS
1	B	367	GLU
1	B	774	ILE
1	A	871	ARG
1	B	314	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	817	HIS
1	B	333	ASN
1	B	691	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	316	GLN
1	B	369	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	ADP	A	1882	-	25,29,29	1.34	3 (12%)	24,45,45	2.16	7 (29%)
2	ADP	B	1882	-	25,29,29	1.65	4 (16%)	24,45,45	2.18	9 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	1882	-	-	0/12/32/32	0/3/3/3
2	ADP	B	1882	-	-	0/12/32/32	0/3/3/3

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1882	ADP	C8-N7	2.05	1.38	1.34
2	A	1882	ADP	PB-O3A	2.73	1.64	1.60
2	A	1882	ADP	C2-N3	2.78	1.36	1.32
2	B	1882	ADP	C2-N3	3.22	1.37	1.32
2	B	1882	ADP	PB-O3A	3.32	1.65	1.60

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1882	ADP	N3-C2-N1	-6.48	123.21	128.86
2	A	1882	ADP	N3-C2-N1	-6.14	123.51	128.86
2	A	1882	ADP	O5'-PA-O1A	-3.36	95.67	109.25
2	B	1882	ADP	O2'-C2'-C1'	-2.76	102.99	111.61
2	B	1882	ADP	C1'-N9-C4	-2.72	121.93	126.64

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1882	ADP	4	0
2	B	1882	ADP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	623/655 (95%)	-0.10	31 (4%) 30 21	61, 97, 174, 284	0
1	B	623/655 (95%)	0.11	40 (6%) 20 15	76, 124, 221, 295	0
All	All	1246/1310 (95%)	0.01	71 (5%) 24 18	61, 110, 212, 295	0

The worst 5 of 71 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	786	ASP	8.5
1	B	782	VAL	7.6
1	B	787	LYS	6.6
1	B	373	PRO	6.5
1	B	689	PRO	5.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ADP	A	1882	27/27	0.84	0.24	0.20	83,89,91,91	0
2	ADP	B	1882	27/27	0.88	0.22	-0.07	92,96,98,99	0

6.5 Other polymers [i](#)

There are no such residues in this entry.