



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 03:30 am GMT

PDB ID : 1APM
Title : 2.0 ANGSTROM REFINED CRYSTAL STRUCTURE OF THE CATALYTIC SUBUNIT OF CAMP-DEPENDENT PROTEIN KINASE COMPLEXED WITH A PEPTIDE INHIBITOR AND DETERGENT
Authors : Knighton, D.R.; Bell, S.M.; Zheng, J.; Teneyck, L.F.; Xuong, N.-H.; Taylor, S.S.; Sowadski, J.M.
Deposited on : 1993-01-18
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	350	 67% 25% . . .
2	I	20	 80% 10% 5% 5%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3075 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP-DEPENDENT PROTEIN KINASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	E	341	Total	C	N	O	P	S	0	1	0
			2696	1749	445	491	3	8			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	139	ALA	SER	ENGINEERED	UNP P05132

- Molecule 2 is a protein called PEPTIDE INHIBITOR PKI(5-24).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	I	20	Total	C	N	O	0	1	0
			164	98	35	31			

- Molecule 3 is N-OCTANE (three-letter code: OCT) (formula: C₈H₁₈).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	E	1	Total C 8 8	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	E	192	Total O 192 192	0	0
4	I	15	Total O 15 15	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: cAMP-DEPENDENT PROTEIN KINASE



• Molecule 2: PEPTIDE INHIBITOR PKI(5-24)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.84Å 75.76Å 81.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 25.67 – 1.95	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-2.00) 90.2 (25.67-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 1.95Å)	Xtriage
Refinement program	TNT, X-PLOR	Depositor
R, R_{free}	0.186 , (Not available) 0.176 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	21.9	Xtriage
Anisotropy	0.239	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 117.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.018 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3075	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, OCT, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	0.94	14/2738 (0.5%)	1.37	36/3714 (1.0%)
2	I	0.88	0/170	1.44	3/226 (1.3%)
All	All	0.94	14/2908 (0.5%)	1.38	39/3940 (1.0%)

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	13	GLU	CD-OE1	6.90	1.33	1.25
1	E	140	GLU	CD-OE2	6.77	1.33	1.25
1	E	230	GLU	CD-OE1	6.20	1.32	1.25
1	E	341	GLU	CD-OE2	6.09	1.32	1.25
1	E	86	GLU	CD-OE1	5.88	1.32	1.25
1	E	127	GLU	CD-OE2	5.75	1.31	1.25
1	E	31	GLU	CD-OE1	5.69	1.31	1.25
1	E	107	GLU	CD-OE1	5.67	1.31	1.25
1	E	208	GLU	CD-OE1	5.53	1.31	1.25
1	E	332	GLU	CD-OE1	5.39	1.31	1.25
1	E	91	GLU	CD-OE1	5.29	1.31	1.25
1	E	346	GLU	CD-OE1	5.24	1.31	1.25
1	E	248	GLU	CD-OE1	5.21	1.31	1.25
1	E	170	GLU	CD-OE1	-5.20	1.20	1.25

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	134	ARG	NE-CZ-NH1	9.29	124.94	120.30
1	E	220	ASP	CB-CG-OD2	-8.15	110.96	118.30
1	E	112	ASP	CB-CG-OD2	-8.07	111.03	118.30
1	E	267	ASP	CB-CG-OD2	-7.92	111.17	118.30
1	E	323	ASP	CB-CG-OD1	-7.88	111.21	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	161	ASP	CB-CG-OD1	-7.71	111.36	118.30
1	E	220	ASP	CB-CG-OD1	7.63	125.17	118.30
1	E	267	ASP	CB-CG-OD1	7.45	125.00	118.30
1	E	184	ASP	CB-CG-OD2	6.94	124.55	118.30
1	E	264	ASP	CB-CG-OD1	-6.85	112.13	118.30
1	E	247	TYR	CB-CG-CD2	-6.78	116.94	121.00
1	E	301	ASP	CB-CG-OD2	6.66	124.30	118.30
1	E	134	ARG	NE-CZ-NH2	-6.59	117.00	120.30
1	E	44	ASP	CB-CG-OD2	-6.57	112.39	118.30
1	E	329	ASP	CB-CG-OD1	-6.56	112.39	118.30
2	I	24	ASP	CB-CG-OD1	6.40	124.06	118.30
1	E	75	ASP	CB-CG-OD1	-6.39	112.55	118.30
1	E	161	ASP	CB-CG-OD2	6.30	123.97	118.30
1	E	60	VAL	CA-CB-CG2	6.25	120.28	110.90
1	E	290	ASP	CB-CG-OD1	-6.23	112.69	118.30
1	E	75	ASP	CB-CG-OD2	6.01	123.71	118.30
1	E	329	ASP	CB-CG-OD2	5.99	123.69	118.30
1	E	264	ASP	CB-CG-OD2	5.97	123.68	118.30
1	E	241	ASP	CB-CG-OD1	5.93	123.63	118.30
1	E	191	VAL	CB-CA-C	-5.92	100.14	111.40
1	E	60	VAL	CA-CB-CG1	5.88	119.73	110.90
1	E	323	ASP	CB-CG-OD2	5.85	123.57	118.30
1	E	276	ASP	CB-CG-OD2	-5.83	113.06	118.30
1	E	301	ASP	CB-CG-OD1	-5.40	113.44	118.30
2	I	24	ASP	CB-CG-OD2	-5.39	113.45	118.30
1	E	93	ARG	NE-CZ-NH2	-5.36	117.62	120.30
2	I	23	HIS	CA-C-N	-5.33	105.47	117.20
1	E	241	ASP	CB-CG-OD2	-5.32	113.51	118.30
1	E	112	ASP	CB-CG-OD1	5.31	123.08	118.30
1	E	328	ASP	CB-CG-OD1	-5.27	113.56	118.30
1	E	276	ASP	CB-CG-OD1	5.25	123.02	118.30
1	E	210	ILE	CA-CB-CG2	5.22	121.34	110.90
1	E	190	ARG	NE-CZ-NH1	5.16	122.88	120.30
1	E	308	ARG	NE-CZ-NH2	-5.04	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	2696	0	2545	49	0
2	I	164	0	155	3	0
3	E	8	0	18	0	0
4	E	192	0	0	7	0
4	I	15	0	0	0	0
All	All	3075	0	2718	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:189:LYS:HG2	1:E:191:VAL:HG22	1.48	0.95
1:E:305:ILE:HD13	1:E:310:VAL:HG11	1.51	0.92
1:E:305:ILE:CD1	1:E:310:VAL:HG11	2.12	0.78
1:E:299:THR:HB	4:E:448:HOH:O	1.84	0.78
1:E:277:LEU:HD12	1:E:277:LEU:H	1.49	0.76
1:E:268:LEU:HD22	1:E:272:LEU:HD22	1.73	0.70
1:E:305:ILE:HD13	1:E:310:VAL:CG1	2.24	0.66
1:E:318:PHE:HB2	4:E:489:HOH:O	1.96	0.65
1:E:172:LEU:HB3	1:E:180:ILE:HD12	1.78	0.65
1:E:303:ILE:N	1:E:303:ILE:HD13	2.12	0.64
1:E:144:ARG:HB2	1:E:296:TRP:CH2	2.35	0.62
1:E:59:LEU:HD22	1:E:60:VAL:N	2.16	0.61
1:E:142:HIS:CD2	1:E:313:PRO:HG2	2.37	0.60
1:E:179:TYR:OH	1:E:310:VAL:HG13	2.03	0.59
2:I:23:HIS:HD2	2:I:24:ASP:HB2	1.67	0.59
1:E:158:HIS:HE1	1:E:220:ASP:OD2	1.87	0.58
1:E:127:GLU:OE1	2:I:18[B]:ARG:NH1	2.37	0.57
1:E:103:LEU:HD22	1:E:185:PHE:HZ	1.71	0.54
1:E:59:LEU:HD22	1:E:60:VAL:H	1.73	0.53
1:E:100:PHE:CG	1:E:101:PRO:HD2	2.44	0.52
1:E:142:HIS:HB2	4:E:537:HOH:O	2.11	0.50
1:E:122:TYR:CE2	1:E:124:ALA:HB2	2.47	0.50
1:E:262:SER:HB2	4:E:522:HOH:O	2.12	0.49
1:E:226:VAL:HG13	1:E:237:PRO:HD2	1.96	0.48
1:E:54:PHE:HD2	1:E:82:LEU:HD12	1.80	0.47
1:E:174:ILE:HD13	1:E:180:ILE:HD13	1.95	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:125:GLY:HA3	1:E:174:ILE:O	2.15	0.46
1:E:210:ILE:HG12	1:E:251:VAL:HG23	1.97	0.46
1:E:157:LEU:O	1:E:162:LEU:HB2	2.16	0.46
1:E:283:ASN:O	1:E:284:LEU:HD23	2.16	0.45
1:E:124:ALA:CB	1:E:318:PHE:CE1	2.99	0.45
1:E:256:ARG:HB3	4:E:607:HOH:O	2.16	0.45
1:E:137:ARG:HD2	1:E:232:ALA:O	2.17	0.44
1:E:265:LEU:HG	1:E:269:LEU:HD22	1.99	0.44
1:E:158:HIS:HD2	4:E:428:HOH:O	2.00	0.44
1:E:55:GLY:HA3	1:E:73:ILE:O	2.18	0.43
1:E:308:ARG:NH1	4:E:530:HOH:O	2.50	0.43
1:E:43:PHE:HA	1:E:61:LYS:O	2.19	0.42
1:E:310:VAL:CG2	1:E:311:GLU:N	2.83	0.42
1:E:201:THR:O	1:E:202:PRO:C	2.58	0.42
1:E:103:LEU:HD22	1:E:185:PHE:CZ	2.53	0.42
1:E:269:LEU:HA	1:E:269:LEU:HD12	1.85	0.42
1:E:40:LEU:HD22	1:E:40:LEU:O	2.20	0.41
1:E:74:LEU:HD12	1:E:74:LEU:HA	1.85	0.41
1:E:100:PHE:CG	1:E:101:PRO:CD	3.04	0.41
1:E:198:LEU:HA	1:E:198:LEU:HD23	1.84	0.41
2:I:23:HIS:CD2	2:I:24:ASP:HB2	2.52	0.40
1:E:163:ILE:HD13	1:E:163:ILE:HG21	1.89	0.40
1:E:40:LEU:HA	1:E:40:LEU:HD23	1.64	0.40
1:E:59:LEU:HD23	1:E:59:LEU:HA	1.73	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	E	338/350 (97%)	328 (97%)	10 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	I	19/20 (95%)	17 (90%)	2 (10%)	0	100	100
All	All	357/370 (96%)	345 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	265/301 (88%)	232 (88%)	33 (12%)	5	3
2	I	16/15 (107%)	14 (88%)	2 (12%)	5	3
All	All	281/316 (89%)	246 (88%)	35 (12%)	6	3

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	14	SER
1	E	19	LEU
1	E	32	THR
1	E	40	LEU
1	E	51	THR
1	E	53	SER
1	E	54	PHE
1	E	56[A]	ARG
1	E	56[B]	ARG
1	E	59	LEU
1	E	60	VAL
1	E	74	LEU
1	E	104	VAL
1	E	107	GLU
1	E	120	MET
1	E	160	LEU
1	E	162	LEU
1	E	180	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	191	VAL
1	E	195	THR
1	E	198	LEU
1	E	210	ILE
1	E	238	PHE
1	E	244	ILE
1	E	263	SER
1	E	268	LEU
1	E	269	LEU
1	E	272	LEU
1	E	273	LEU
1	E	295	LYS
1	E	299	THR
1	E	335	ILE
1	E	348	THR
2	I	22	ILE
2	I	24	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	36	ASN
1	E	67	ASN
1	E	113	ASN
1	E	158	HIS
1	E	271	ASN
2	I	23	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	SEP	E	10	1	9,9,10	1.63	1 (11%)	9,12,14	4.58	3 (33%)
1	TPO	E	197	1	9,10,11	1.08	0	10,14,16	1.50	2 (20%)
1	SEP	E	338	1	9,9,10	0.75	0	9,12,14	3.02	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	E	10	1	-	0/5/8/10	0/0/0/0
1	TPO	E	197	1	-	0/8/11/13	0/0/0/0
1	SEP	E	338	1	-	0/5/8/10	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	10	SEP	CA-C	4.13	1.55	1.50

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	10	SEP	O3P-P-O1P	-9.84	71.99	110.50
1	E	338	SEP	O3P-P-O2P	-6.40	81.78	107.61
1	E	197	TPO	CG2-CB-CA	-2.38	108.81	113.22
1	E	197	TPO	C-CA-N	3.44	116.79	109.86
1	E	10	SEP	OG-CB-CA	3.70	111.81	108.17
1	E	338	SEP	O2P-P-O1P	6.00	133.98	110.50
1	E	10	SEP	O2P-P-O1P	8.52	143.85	110.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	OCT	E	354	-	7,7,7	0.68	0	6,6,6	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OCT	E	354	-	-	0/5/5/5	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	338/350 (96%)	-0.71	1 (0%) 93 93	12, 26, 60, 88	0
2	I	20/20 (100%)	-0.92	0 100 100	14, 20, 72, 92	0
All	All	358/370 (96%)	-0.72	1 (0%) 93 93	12, 26, 61, 92	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	318	PHE	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	SEP	E	10	10/11	0.88	0.10	-	34,51,100,100	0
1	SEP	E	338	10/11	0.97	0.11	-	26,48,58,73	0
1	TPO	E	197	11/12	0.98	0.06	-	12,17,21,29	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	OCT	E	354	8/8	0.92	0.13	0.96	25,31,50,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.