



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 01:29 am GMT

PDB ID : 1APW  
Title : CRYSTALLOGRAPHIC ANALYSIS OF TRANSITION STATE MIMICS  
BOUND TO PENICILLOPEPSIN: DIFLUOROSTATINE-AND DIFLUOR  
OSTATONE-CONTAINING PEPTIDES  
Authors : Sielecki, A.R.; James, M.N.G.  
Deposited on : 1991-12-16  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	E	323	
2	I	5	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	HSY	E	399	X	-	-	-

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 2726 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PENICILLOPEPSIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	323	Total	C	N	O	S	0	0	0
			2366	1479	377	508	2			

- Molecule 2 is a protein called INHIBITOR ISOVALERYL (IVA)-VAL-VAL-DIFLUOROS TATINE-N-METHYLAMINE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	5	Total	C	F	N	O	0	0	1
			35	24	2	4	5			

- Molecule 3 is SUGAR (ALPHA-D-MANNOSE) (three-letter code: MAN) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	E	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is SUGAR (XYLOPYRANOSE) (three-letter code: HSY) (formula: C<sub>5</sub>H<sub>10</sub>O<sub>5</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	E	1	Total	C	O	0	0
			9	5	4		

- Molecule 5 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	E	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is DIMETHYLFORMAMIDE (three-letter code: DMF) (formula: C<sub>3</sub>H<sub>7</sub>NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	I	1	Total	C	N	O	0	0
			5	3	1	1		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	E	291	Total	O	0	0
			291	291		
7	I	4	Total	O	0	0
			4	4		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

#### • Molecule 1: PENICILLOPEPSIN



#### • Molecule 2: INHIBITOR ISOVALERYL (IVA)-VAL-VAL-DIFLUOROSTATINE-N-METHYL AMINE



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.64Å 46.58Å 66.44Å 90.00° 116.12° 90.00°	Depositor
Resolution (Å)	8.00 – 1.80	Depositor
% Data completeness (in resolution range)	(Not available) (8.00-1.80)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, $R_{free}$	0.131 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2726	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IVA, DMF, NME, DFI, SO4, HSY, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	E	1.01	0/2420	1.72	29/3304 (0.9%)
2	I	0.86	0/13	2.05	1/17 (5.9%)
All	All	1.01	0/2433	1.72	30/3321 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	I	0	1

There are no bond length outliers.

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	222	ASP	CB-CG-OD1	7.94	125.44	118.30
1	E	52	SER	O-C-N	7.82	136.49	123.20
1	E	141	PHE	CB-CG-CD1	-7.09	115.84	120.80
1	E	222	ASP	CB-CG-OD2	-6.92	112.07	118.30
1	E	206	ASP	CB-CG-OD1	6.88	124.49	118.30
1	E	119	ASP	CB-CG-OD2	6.19	123.87	118.30
1	E	155	VAL	CG1-CB-CG2	-6.04	101.24	110.90
1	E	131	THR	CA-CB-CG2	5.97	120.75	112.40
1	E	218	LEU	O-C-N	5.97	132.25	122.70
1	E	38	ASP	CB-CG-OD1	5.82	123.54	118.30
1	E	74	SER	CB-CA-C	5.82	121.16	110.10
1	E	1	ALA	CB-CA-C	5.80	118.80	110.10
1	E	69	TYR	CB-CG-CD2	-5.73	117.56	121.00
1	E	196	ASP	CB-CG-OD1	-5.57	113.29	118.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	165	TYR	N-CA-CB	5.53	120.55	110.60
1	E	306	GLN	CB-CA-C	-5.51	99.38	110.40
1	E	18	ILE	CB-CA-C	5.51	122.62	111.60
1	E	256	PHE	CB-CG-CD2	-5.50	116.95	120.80
1	E	43	SER	O-C-N	5.44	131.41	122.70
1	E	307	TYR	N-CA-C	-5.44	96.32	111.00
1	E	295	PHE	CB-CG-CD2	-5.36	117.05	120.80
1	E	308	VAL	O-C-N	5.33	131.23	122.70
1	E	307	TYR	CG-CD1-CE1	-5.28	117.08	121.30
1	E	160	GLN	CG-CD-NE2	5.21	129.20	116.70
1	E	55	SER	N-CA-C	-5.21	96.95	111.00
1	E	274	TYR	CB-CG-CD1	-5.12	117.93	121.00
1	E	250	SER	CB-CA-C	-5.11	100.39	110.10
1	E	142	ASP	CB-CG-OD1	5.09	122.88	118.30
1	E	2	ALA	CA-C-N	-5.05	106.10	117.20
2	I	3	VAL	O-C-N	5.03	130.75	122.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	I	1	DFI	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	2366	0	2181	16	0
2	I	35	0	39	1	0
3	E	11	0	10	0	0
4	E	9	0	7	0	0
5	E	5	0	0	0	0
6	I	5	0	7	2	0
7	E	291	0	0	4	0
7	I	4	0	0	0	0
All	All	2726	0	2244	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:233:VAL:HG13	1:E:251:THR:HG21	1.81	0.62
1:E:121:LEU:HD21	2:I:1:DFI:HD12	1.89	0.55
1:E:160:GLN:HG3	7:E:594:HOH:O	2.09	0.51
1:E:239:ASP:OD1	1:E:279:ASP:OD2	2.29	0.51
1:E:150:GLN:HG2	7:E:616:HOH:O	2.10	0.51
7:E:570:HOH:O	6:I:587:DMF:H21	2.11	0.50
1:E:306:GLN:C	1:E:322:GLN:HG3	2.33	0.48
1:E:172:SER:HA	1:E:175:TYR:CE1	2.52	0.45
7:E:570:HOH:O	6:I:587:DMF:C2	2.66	0.43
1:E:151:PRO:HB2	1:E:312:SER:OG	2.19	0.43
1:E:106:GLN:C	1:E:107:GLN:HG2	2.37	0.42
1:E:40:TRP:HA	1:E:103:GLN:O	2.20	0.42
1:E:125:ALA:HB1	1:E:189:GLY:O	2.20	0.41
1:E:141:PHE:O	1:E:145:LYS:HB3	2.18	0.41
1:E:253:LEU:HA	1:E:254:PRO:HD3	1.94	0.41
1:E:203:GLN:OE1	1:E:228:GLN:HG3	2.21	0.41
1:E:66:LEU:HA	1:E:66:LEU:HD23	1.75	0.40
1:E:71:TRP:CE2	1:E:103:GLN:HB3	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	321/323 (99%)	319 (99%)	2 (1%)	0	100	100
2	I	2/5 (40%)	2 (100%)	0	0	100	100
All	All	323/328 (98%)	321 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	259/259 (100%)	252 (97%)	7 (3%)	50	35
2	I	2/2 (100%)	2 (100%)	0	100	100
All	All	261/261 (100%)	254 (97%)	7 (3%)	50	35

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	18	ILE
1	E	31	ASN
1	E	38	ASP
1	E	39	LEU
1	E	107	GLN
1	E	234	SER
1	E	281	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	13	ASN
1	E	31	ASN
1	E	50	GLN
1	E	107	GLN
1	E	150	GLN
1	E	194	ASN
1	E	228	GLN
1	E	238	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	DFI	I	1	2	9,12,13	1.70	2 (22%)	6,17,19	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DFI	I	1	2	-	0/11/17/20	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	1	DFI	F1-CM	-3.67	1.33	1.37
2	I	1	DFI	F2-CM	-2.09	1.35	1.37

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	I	1	DFI	1	0

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	MAN	E	328	1	11,11,12	0.79	0	13,15,17	1.31	1 (7%)
5	SO4	E	365	-	4,4,4	0.63	0	6,6,6	0.32	0
4	HSY	E	399	1	9,9,10	0.74	0	10,12,14	3.34	3 (30%)
6	DMF	I	587	-	4,4,4	0.22	0	4,4,4	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	E	328	1	-	0/2/19/22	0/1/1/1
5	SO4	E	365	-	-	0/0/0/0	0/0/0/0
4	HSY	E	399	1	3/3/3/4	0/0/14/17	0/1/1/1
6	DMF	I	587	-	-	0/2/2/2	0/0/0/0

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	328	MAN	O2-C2-C1	-3.75	101.56	109.18
4	E	399	HSY	O4-C4-C5	4.26	117.84	109.18
4	E	399	HSY	C5-C4-C3	5.83	117.05	109.65
4	E	399	HSY	C1-O5-C5	6.96	122.25	111.61

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	E	399	HSY	C2
4	E	399	HSY	C3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atom
4	E	399	HSY	C4

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	I	587	DMF	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.