



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 13, 2017 – 07:15 am GMT

PDB ID : 1AQK
Title : THREE-DIMENSIONAL STRUCTURE OF A HUMAN FAB WITH HIGH AFFINITY FOR TETANUS TOXOID
Authors : Faber, C.; Fan, Z.; Edmundson, A.B.
Deposited on : 1997-07-30
Resolution : 1.84 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	3197 (1.86-1.82)
Ramachandran outliers	110173	3164 (1.86-1.82)
Sidechain outliers	110143	3165 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	L	216	
2	H	226	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3597 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FAB B7-15A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	216	Total	C	N	O	S	0	0	0
			1597	995	272	325	5			

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	17	ARG	LYS	CONFLICT	UNP P01842
L	18	VAL	ILE	CONFLICT	UNP P01842
L	23	THR	SER	CONFLICT	UNP P01842
L	25	SER	THR	CONFLICT	UNP P01842
L	26	ASN	SER	CONFLICT	UNP P01842
L	33	PHE	HIS	CONFLICT	UNP P01842
L	34	THR	HIS	CONFLICT	UNP P01842
L	40	HIS	GLN	CONFLICT	UNP P01842
L	41	LEU	VAL	CONFLICT	UNP P01842
L	51	PHE	TYR	CONFLICT	UNP P01842
L	53	ASN	ASP	CONFLICT	UNP P01842
L	54	THR	ASN	CONFLICT	UNP P01842
L	64	PHE	ILE	CONFLICT	UNP P01842
L	81	GLN	ARG	CONFLICT	UNP P01842
L	93	TYR	PHE	CONFLICT	UNP P01842
L	?	-	GLY	DELETION	UNP P01842
L	99	ALA	TRP	CONFLICT	UNP P01842
L	100	ARG	VAL	CONFLICT	UNP P01842
L	104	GLY	ALA	CONFLICT	UNP P01842
L	106	ARG	LYS	CONFLICT	UNP P01842
L	160	ASN	THR	CONFLICT	UNP P01842
L	167	LYS	THR	CONFLICT	UNP P01842
L	213	ALA	THR	CONFLICT	UNP P01842

- Molecule 2 is a protein called FAB B7-15A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	226	Total	C	N	O	S	0	0	0
			1711	1088	289	327	7			

There are 31 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	30	ASN	SER	CONFLICT	UNP P01857
H	33	ALA	GLY	CONFLICT	UNP P01857
H	34	ILE	MET	CONFLICT	UNP P01857
H	50	PHE	ALA	CONFLICT	UNP P01857
H	52	SER	TRP	CONFLICT	UNP P01857
H	57	LYS	ASN	CONFLICT	UNP P01857
H	58	ASN	LYS	CONFLICT	UNP P01857
H	80	PHE	TYR	CONFLICT	UNP P01857
H	81	LEU	MET	CONFLICT	UNP P01857
H	88	PRO	ALA	CONFLICT	UNP P01857
H	93	ILE	VAL	CONFLICT	UNP P01857
H	99	VAL	-	INSERTION	UNP P01857
H	100	LEU	-	INSERTION	UNP P01857
H	101	PHE	GLU	CONFLICT	UNP P01857
H	102	GLN	GLY	CONFLICT	UNP P01857
H	103	GLN	ARG	CONFLICT	UNP P01857
H	104	LEU	TRP	CONFLICT	UNP P01857
H	106	LEU	ARG	CONFLICT	UNP P01857
H	?	-	THR	DELETION	UNP P01857
H	?	-	THR	DELETION	UNP P01857
H	?	-	VAL	DELETION	UNP P01857
H	?	-	THR	DELETION	UNP P01857
H	?	-	THR	DELETION	UNP P01857
H	?	-	ILE	DELETION	UNP P01857
H	?	-	GLY	DELETION	UNP P01857
H	108	ALA	TYR	CONFLICT	UNP P01857
H	109	PRO	TYR	CONFLICT	UNP P01857
H	112	ILE	TYR	CONFLICT	UNP P01857
H	118	MET	LEU	CONFLICT	UNP P01857
H	158	GLN	GLU	CONFLICT	UNP P01857
H	220	LYS	ARG	CONFLICT	UNP P01857

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	161	Total	O	0	0
			161	161		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	L	128	Total 128	O 128	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: FAB B7-15A2



• Molecule 2: FAB B7-15A2



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	40.70Å 78.60Å 74.60Å 90.00° 104.30° 90.00°	Depositor
Resolution (Å)	20.00 – 1.84	Depositor
% Data completeness (in resolution range)	81.0 (20.00-1.84)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	TNT V. 5-E, X-PLOR	Depositor
R, R_{free}	0.185 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3597	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	1.05	7/1629 (0.4%)	1.50	22/2225 (1.0%)
2	H	1.02	3/1747 (0.2%)	1.49	24/2380 (1.0%)
All	All	1.03	10/3376 (0.3%)	1.50	46/4605 (1.0%)

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	46	GLU	CD-OE1	-6.86	1.18	1.25
1	L	202	GLU	CD-OE1	6.29	1.32	1.25
1	L	214	GLU	CD-OE1	5.92	1.32	1.25
2	H	222	GLU	CD-OE1	5.92	1.32	1.25
1	L	187	GLU	CD-OE2	5.81	1.32	1.25

The worst 5 of 46 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	17	ARG	NE-CZ-NH2	-10.69	114.95	120.30
1	L	17	ARG	NE-CZ-NH1	9.83	125.22	120.30
2	H	19	ARG	NE-CZ-NH1	9.68	125.14	120.30
2	H	73	ASP	CB-CG-OD2	-9.05	110.15	118.30
2	H	67	ARG	NE-CZ-NH2	8.53	124.57	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1597	0	1541	92	0
2	H	1711	0	1685	122	0
3	H	161	0	0	24	0
3	L	128	0	0	18	0
All	All	3597	0	3226	202	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 31.

The worst 5 of 202 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:2:ASN:H	1:L:100:ARG:NH2	1.51	1.07
2:H:224:LYS:HZ2	2:H:224:LYS:N	1.52	1.06
2:H:50:PHE:HE1	2:H:57:LYS:HE2	1.29	0.97
2:H:101:PHE:HB2	2:H:105:VAL:HG11	1.47	0.96
2:H:72:ARG:HE	2:H:74:ASN:HD21	1.13	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	214/216 (99%)	194 (91%)	14 (6%)	6 (3%)	6	0
2	H	224/226 (99%)	208 (93%)	13 (6%)	3 (1%)	14	3
All	All	438/442 (99%)	402 (92%)	27 (6%)	9 (2%)	8	1

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	213	ALA
1	L	215	CYS
1	L	97	LEU
2	H	142	SER
1	L	95	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	177/177 (100%)	143 (81%)	34 (19%)	1	0
2	H	191/191 (100%)	165 (86%)	26 (14%)	4	0
All	All	368/368 (100%)	308 (84%)	60 (16%)	3	0

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	185	THR
1	L	215	CYS
2	H	219	LYS
1	L	196	SER
2	H	16	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	58	ASN
2	H	74	ASN
2	H	158	GLN
2	H	31	ASN
2	H	202	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PCA	H	1	2	8,8,9	1.77	1 (12%)	9,10,12	1.39	1 (11%)
1	PCA	L	1	1	8,8,9	2.01	2 (25%)	9,10,12	1.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	H	1	2	-	0/0/11/13	0/1/1/1
1	PCA	L	1	1	-	0/0/11/13	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	1	PCA	CA-C	2.26	1.53	1.50
2	H	1	PCA	CD-N	4.40	1.47	1.34
1	L	1	PCA	CD-N	4.95	1.48	1.34

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	1	PCA	OE-CD-CG	-2.68	121.93	126.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	L	1	PCA	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.