



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 6, 2017 – 06:33 PM EST

PDB ID : 6AQX
Title : Crystal Structure of Z-DNA with 6-fold Twinning_Z4B
Authors : Luo, Z.; Dauter, Z.; Gilski, M.
Deposited on : unknown
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1088 (1.56-1.56)
Clashscore	112137	1132 (1.56-1.56)
RSRZ outliers	101464	1089 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	6	
1	B	6	
1	C	6	
1	D	6	
1	E	6	
1	F	6	
1	G	6	
1	H	6	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 1001 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*CP*GP*CP*GP*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	6	Total	C	N	O	P	0	0	0
			120	57	24	34	5			
1	B	6	Total	C	N	O	P	0	0	0
			120	57	24	34	5			
1	C	6	Total	C	N	O	P	0	0	0
			120	57	24	34	5			
1	D	6	Total	C	N	O	P	0	0	0
			120	57	24	34	5			
1	E	6	Total	C	N	O	P	0	0	0
			120	57	24	34	5			
1	F	6	Total	C	N	O	P	0	0	0
			120	57	24	34	5			
1	G	6	Total	C	N	O	P	0	0	0
			120	57	24	34	5			
1	H	6	Total	C	N	O	P	0	0	0
			120	57	24	34	5			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	O	0	0
			4	4		
2	B	6	Total	O	0	0
			6	6		
2	C	6	Total	O	0	0
			6	6		
2	D	6	Total	O	0	0
			6	6		
2	E	4	Total	O	0	0
			4	4		
2	F	6	Total	O	0	0
			6	6		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	4	Total	O	0	0
			4	4		
2	H	5	Total	O	0	0
			5	5		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(*CP*GP*CP*GP*CP*G)-3')

Chain A: 




- Molecule 1: DNA (5'-D(*CP*GP*CP*GP*CP*G)-3')

Chain B: 



- Molecule 1: DNA (5'-D(*CP*GP*CP*GP*CP*G)-3')

Chain C: 




- Molecule 1: DNA (5'-D(*CP*GP*CP*GP*CP*G)-3')

Chain D: 

There are no outlier residues recorded for this chain.

- Molecule 1: DNA (5'-D(*CP*GP*CP*GP*CP*G)-3')

Chain E: 



- Molecule 1: DNA (5'-D(*CP*GP*CP*GP*CP*G)-3')

Chain F: 

C607
G608
C609
G610
C611
G612

- Molecule 1: DNA (5'-D(*CP*GP*CP*GP*CP*G)-3')

Chain G:  67% 33%

C701
G702
C703
G704
C705
G706

- Molecule 1: DNA (5'-D(*CP*GP*CP*GP*CP*G)-3')

Chain H:  100%

C807
G808
C809
G810
C811
G812

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	35.20 Å 42.22 Å 35.21 Å 90.00° 119.83° 90.00°	Depositor
Resolution (Å)	30.54 – 1.55 30.54 – 1.55	Depositor EDS
% Data completeness (in resolution range)	99.2 (30.54-1.55) 99.7 (30.54-1.55)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.03 (at 1.54 Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.209 , 0.227 0.216 , 0.235	Depositor DCC
R_{free} test set	656 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	15.2	Xtriage
Anisotropy	0.466	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 76.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.101 for -h-l,k,h 0.101 for l,k,-h-l 0.096 for h,-k,-h-l 0.108 for -h-l,-k,l 0.469 for l,-k,h	Xtriage
Reported twinning fraction	0.354 for H, K, L 0.375 for L, -K, H 0.073 for -H, -K, H+L 0.082 for L, K, -H-L 0.061 for -H-L, K, H 0.055 for H+L, -K, -L	Depositor
Outliers	0 of 13133 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	1001	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 90.08 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.6527e-08. The detected translational NCS is most likely*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

also responsible for the elevated intensity ratio.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.67	0/134	1.14	0/205
1	B	0.71	0/134	1.03	0/205
1	C	0.69	0/134	1.16	0/205
1	D	0.73	0/134	1.04	0/205
1	E	0.55	0/134	1.16	0/205
1	F	0.83	0/134	1.03	0/205
1	G	0.61	0/134	1.15	0/205
1	H	0.79	0/134	1.04	0/205
All	All	0.70	0/1072	1.09	0/1640

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	120	0	68	4	0
1	B	120	0	68	7	0
1	C	120	0	68	7	0
1	D	120	0	68	0	0
1	E	120	0	68	1	0
1	F	120	0	68	7	1
1	G	120	0	68	6	0
1	H	120	0	68	6	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	4	0	0	0	0
2	B	6	0	0	0	0
2	C	6	0	0	0	0
2	D	6	0	0	0	0
2	E	4	0	0	0	0
2	F	6	0	0	0	0
2	G	4	0	0	0	0
2	H	5	0	0	0	0
All	All	1001	0	544	29	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:210:DG:H4'	1:G:704:DG:H5'	1.41	1.02
1:H:810:DG:H2'	1:H:811:DC:C5	2.18	0.78
1:F:610:DG:H2'	1:F:611:DC:C5	2.18	0.78
1:F:608:DG:H2'	1:F:608:DG:N3	2.03	0.72
1:H:810:DG:H2'	1:H:811:DC:H5	1.54	0.71
1:F:610:DG:H2'	1:F:611:DC:H5	1.56	0.70
1:B:210:DG:H4'	1:G:704:DG:C5'	2.24	0.66
1:B:210:DG:H1'	1:G:703:DC:OP1	1.99	0.63
1:H:808:DG:H2'	1:H:808:DG:N3	2.14	0.63
1:F:608:DG:H2''	1:F:609:DC:OP1	2.01	0.61
1:H:808:DG:H2''	1:H:809:DC:OP1	2.04	0.56
1:C:303:DC:H5''	1:F:610:DG:H1'	1.88	0.55
1:G:703:DC:H4'	1:G:704:DG:O5'	2.10	0.51
1:B:208:DG:O5'	1:C:306:DG:H1'	2.14	0.48
1:B:208:DG:H1'	1:C:305:DC:OP2	2.14	0.46
1:F:609:DC:H4'	1:F:610:DG:O5'	2.17	0.45
1:C:303:DC:OP1	1:F:610:DG:H1'	2.16	0.45
1:C:303:DC:H4'	1:C:304:DG:O5'	2.17	0.45
1:B:210:DG:C5'	1:G:704:DG:H4'	2.47	0.45
1:A:103:DC:H4'	1:A:104:DG:O5'	2.18	0.44
1:E:503:DC:H4'	1:E:504:DG:O5'	2.18	0.43
1:C:304:DG:H2'	1:C:305:DC:C5	2.54	0.43
1:C:305:DC:O4'	1:C:306:DG:C2	2.73	0.42
1:A:104:DG:H2'	1:A:105:DC:C5	2.55	0.42
1:H:808:DG:N3	1:H:808:DG:C2'	2.83	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:105:DC:O4'	1:A:106:DG:C2	2.74	0.41
1:H:809:DC:H4'	1:H:810:DG:O5'	2.21	0.41
1:A:102:DG:H2'	1:A:103:DC:C6	2.56	0.40
1:B:210:DG:C1'	1:G:703:DC:OP1	2.68	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:807:DC:O5'	1:H:812:DG:O3'[2_556]	2.09	0.11
1:F:607:DC:O5'	1:F:612:DG:O3'[2_655]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	6/6 (100%)	0.83	0 100 100	22, 28, 29, 30	0
1	B	6/6 (100%)	0.92	0 100 100	22, 24, 28, 31	0
1	C	6/6 (100%)	0.41	0 100 100	21, 27, 29, 29	0
1	D	6/6 (100%)	0.29	0 100 100	22, 24, 27, 31	0
1	E	6/6 (100%)	0.72	0 100 100	38, 45, 49, 51	0
1	F	6/6 (100%)	0.76	0 100 100	32, 40, 42, 57	0
1	G	6/6 (100%)	0.88	0 100 100	38, 43, 53, 59	0
1	H	6/6 (100%)	0.88	0 100 100	35, 40, 59, 68	0
All	All	48/48 (100%)	0.71	0 100 100	21, 31, 57, 68	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.