



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 01:23 pm GMT

PDB ID : 1ARO
Title : T7 RNA POLYMERASE COMPLEXED WITH T7 LYSOZYME
Authors : Steitz, T.; Jeruzalmi, D.
Deposited on : 1997-08-08
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	P	883	
2	L	151	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7314 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T7 RNA POLYMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	P	774	6124	3912	1065	1115	32	161	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
P	347	SER	CYS	CONFLICT	UNP P00573
P	723	SER	CYS	ENGINEERED	UNP P00573
P	839	SER	CYS	ENGINEERED	UNP P00573

- Molecule 2 is a protein called T7 LYSOZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	149	1183	742	220	216	5	23	0	0


- Molecule 3 is MERCURY (II) ION (three-letter code: HG) (formula: Hg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	P	6	Total	Hg	0	0
			6	6		
3	L	1	Total	Hg	0	0
			1	1		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: T7 RNA POLYMERASE





● Molecule 2: T7 LYSOZYME



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	273.38Å 95.61Å 63.58Å 90.00° 101.40° 90.00°	Depositor
Resolution (Å)	30.00 – 2.80 33.50 – 2.78	Depositor EDS
% Data completeness (in resolution range)	93.9 (30.00-2.80) 90.1 (33.50-2.78)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.54 (at 2.76Å)	Xtrriage
Refinement program	CNS, X-PLOR	Depositor
R, R_{free}	0.262 , 0.309 0.276 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	68.0	Xtrriage
Anisotropy	0.523	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 83.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.025 for -h-2*1,-k,l	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	7314	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	P	0.69	14/6261 (0.2%)	1.05	41/8460 (0.5%)
2	L	0.57	1/1210 (0.1%)	0.81	1/1630 (0.1%)
All	All	0.68	15/7471 (0.2%)	1.01	42/10090 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	P	0	10
2	L	0	1
All	All	0	11

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	P	74	ILE	N-CA	14.10	1.74	1.46
1	P	559	VAL	C-N	-13.68	1.02	1.34
1	P	499	ASN	C-N	-10.51	1.09	1.34
1	P	497	LEU	C-N	-9.12	1.13	1.34
1	P	74	ILE	CA-CB	9.09	1.75	1.54

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	P	8	LYS	O-C-N	-20.13	90.50	122.70
1	P	559	VAL	O-C-N	-17.78	94.25	122.70
1	P	559	VAL	C-N-CA	14.85	158.83	121.70
1	P	498	GLU	CB-CA-C	-13.31	83.79	110.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	P	499	ASN	CB-CA-C	13.18	136.76	110.40

There are no chirality outliers.

5 of 11 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	P	162	PHE	Sidechain
1	P	497	LEU	Mainchain
1	P	74	ILE	Mainchain
1	P	8	LYS	Mainchain,Peptide
1	P	9	ASN	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	P	6124	0	6086	580	1
2	L	1183	0	1156	88	0
3	L	1	0	0	0	0
3	P	6	0	0	2	0
All	All	7314	0	7242	662	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 47.

The worst 5 of 662 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:P:74:ILE:CB	1:P:74:ILE:CA	1.75	1.55
1:P:74:ILE:N	1:P:74:ILE:CA	1.74	1.45
1:P:73:LEU:C	1:P:74:ILE:HA	1.18	1.45
1:P:74:ILE:N	1:P:75:THR:H	1.10	1.40
1:P:74:ILE:N	1:P:75:THR:N	1.82	1.25

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:P:233:ASN:OD1	1:P:838:SER:O[1_554]	1.63	0.57

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	P	760/883 (86%)	591 (78%)	120 (16%)	49 (6%)	1 4
2	L	147/151 (97%)	110 (75%)	34 (23%)	3 (2%)	9 28
All	All	907/1034 (88%)	701 (77%)	154 (17%)	52 (6%)	2 5

5 of 52 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	P	94	ALA
1	P	197	GLY
1	P	498	GLU
1	P	554	VAL
1	P	584	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	P	642/729 (88%)	608 (95%)	34 (5%)	26 59
2	L	124/125 (99%)	116 (94%)	8 (6%)	20 49
All	All	766/854 (90%)	724 (94%)	42 (6%)	25 57

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	P	452	ILE
1	P	562	LEU
2	L	1036	HIS
1	P	471	ASP
1	P	510	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 39 such sidechains are listed below:

Mol	Chain	Res	Type
1	P	648	GLN
1	P	758	GLN
2	L	1074	HIS
1	P	697	ASN
1	P	737	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	P	774/883 (87%)	0.00	36 (4%) 32 22	19, 77, 112, 123	41 (5%)
2	L	149/151 (98%)	-0.23	3 (2%) 65 56	46, 66, 105, 114	5 (3%)
All	All	923/1034 (89%)	-0.03	39 (4%) 37 26	19, 75, 110, 123	46 (4%)

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	P	233	ASN	6.5
1	P	717	GLU	6.3
1	P	712	ASP	4.6
1	P	751	PHE	4.5
1	P	711	LYS	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	HG	L	903	1/1	0.97	0.11	-1.02	113,113,113,113	0
3	HG	P	907	1/1	0.99	0.04	-4.30	159,159,159,159	0
3	HG	P	908	1/1	0.99	0.03	-4.50	199,199,199,199	0
3	HG	P	905	1/1	0.99	0.12	-	83,83,83,83	0
3	HG	P	906	1/1	0.95	0.16	-	136,136,136,136	0
3	HG	P	904	1/1	0.99	0.15	-	87,87,87,87	0
3	HG	P	909	1/1	0.95	0.15	-	199,199,199,199	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.