



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 07:04 am GMT

PDB ID : 4AUN
Title : Crystal structure, recombinant expression and mutagenesis studies of the bi-functional catalase-phenol oxidase from *Scytalidium thermophilum*
Authors : Yuzugullu, Y.; Trinh, C.H.; Smith, M.A.; Pearson, A.R.; Phillips, S.E.V.; Sutay Kocabas, D.; Bakir, U.; Ogel, Z.B.; McPherson, M.J.
Deposited on : 2012-05-18
Resolution : 1.92 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	6276 (1.94-1.90)
Clashscore	112137	7025 (1.94-1.90)
Ramachandran outliers	110173	6947 (1.94-1.90)
Sidechain outliers	110143	6948 (1.94-1.90)
RSRZ outliers	101464	6332 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	719	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 84%; height: 10px; background-color: red;"></div> <div style="width: 84%; height: 10px; background-color: orange;"></div> <div style="width: 84%; height: 10px; background-color: yellow;"></div> <div style="width: 84%; height: 10px; background-color: green;"></div> <div style="width: 8%; height: 10px; background-color: grey;"></div> </div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%;"></div> </div> <div>84% 8% • 7%</div> </div>
1	B	719	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 85%; height: 10px; background-color: red;"></div> <div style="width: 85%; height: 10px; background-color: orange;"></div> <div style="width: 85%; height: 10px; background-color: yellow;"></div> <div style="width: 85%; height: 10px; background-color: green;"></div> <div style="width: 8%; height: 10px; background-color: grey;"></div> </div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%;"></div> </div> <div>85% 8% • 6%</div> </div>
1	C	719	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 86%; height: 10px; background-color: red;"></div> <div style="width: 86%; height: 10px; background-color: orange;"></div> <div style="width: 86%; height: 10px; background-color: yellow;"></div> <div style="width: 86%; height: 10px; background-color: green;"></div> <div style="width: 7%; height: 10px; background-color: grey;"></div> </div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%;"></div> </div> <div>86% 7% • 6%</div> </div>
1	D	719	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 86%; height: 10px; background-color: red;"></div> <div style="width: 86%; height: 10px; background-color: orange;"></div> <div style="width: 86%; height: 10px; background-color: yellow;"></div> <div style="width: 86%; height: 10px; background-color: green;"></div> <div style="width: 7%; height: 10px; background-color: grey;"></div> </div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%;"></div> </div> <div>86% 7% • 6%</div> </div>
1	E	719	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 83%; height: 10px; background-color: red;"></div> <div style="width: 83%; height: 10px; background-color: orange;"></div> <div style="width: 83%; height: 10px; background-color: yellow;"></div> <div style="width: 83%; height: 10px; background-color: green;"></div> <div style="width: 9%; height: 10px; background-color: grey;"></div> </div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%;"></div> </div> <div>83% 9% • 7%</div> </div>
1	F	719	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 84%; height: 10px; background-color: red;"></div> <div style="width: 84%; height: 10px; background-color: orange;"></div> <div style="width: 84%; height: 10px; background-color: yellow;"></div> <div style="width: 84%; height: 10px; background-color: green;"></div> <div style="width: 7%; height: 10px; background-color: grey;"></div> </div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%;"></div> </div> <div>84% 7% • 7%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	719	 85% 8% • 6%
1	H	719	 84% 8% • 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	HDD	A	900	-	-	-	X
2	HDD	B	900	-	-	-	X
2	HDD	C	900	-	-	-	X
2	HDD	D	900	-	-	-	X
2	HDD	G	900	-	-	-	X
2	HDD	H	900	-	-	-	X

2 Entry composition


There are 4 unique types of molecules in this entry. The entry contains 46567 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CATALASE-PHENOL OXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	671	Total	C	N	O	S	0	19	0
			5357	3388	933	1025	11			
1	B	673	Total	C	N	O	S	0	19	0
			5364	3395	931	1026	12			
1	C	673	Total	C	N	O	S	0	20	0
			5375	3401	937	1025	12			
1	D	674	Total	C	N	O	S	0	18	0
			5368	3396	937	1023	12			
1	E	668	Total	C	N	O	S	0	19	0
			5324	3371	925	1017	11			
1	F	668	Total	C	N	O	S	0	16	0
			5314	3365	928	1010	11			
1	G	673	Total	C	N	O	S	0	21	0
			5374	3402	935	1025	12			
1	H	670	Total	C	N	O	S	0	16	0
			5325	3371	927	1016	11			

- Molecule 2 is CIS-HEME D HYDROXYCHLORIN GAMMA-SPIROLACTONE (three-letter code: HDD) (formula: C₃₄H₃₂FeN₄O₅).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			44	34	1	4	5		
2	B	1	Total	C	Fe	N	O	0	0
			44	34	1	4	5		
2	C	1	Total	C	Fe	N	O	0	0
			44	34	1	4	5		
2	D	1	Total	C	Fe	N	O	0	0
			44	34	1	4	5		
2	E	1	Total	C	Fe	N	O	0	0
			44	34	1	4	5		
2	F	1	Total	C	Fe	N	O	0	0
			44	34	1	4	5		
2	G	1	Total	C	Fe	N	O	0	0
			44	34	1	4	5		
2	H	1	Total	C	Fe	N	O	0	0
			44	34	1	4	5		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	1	Total	Ca	0	0
			1	1		
3	D	1	Total	Ca	0	0
			1	1		
3	E	1	Total	Ca	0	0
			1	1		
3	H	1	Total	Ca	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Ca 1	0	0
3	C	1	Total 1	Ca 1	0	0
3	A	2	Total 2	Ca 2	0	0
3	F	1	Total 1	Ca 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	521	Total 521	O 521	0	0
4	B	420	Total 420	O 420	0	0
4	C	470	Total 470	O 470	0	0
4	D	468	Total 468	O 468	0	0
4	E	341	Total 341	O 341	0	0
4	F	381	Total 381	O 381	0	0
4	G	454	Total 454	O 454	0	0
4	H	350	Total 350	O 350	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CATALASE-PHENOL OXIDASE



• Molecule 1: CATALASE-PHENOL OXIDASE




• Molecule 1: CATALASE-PHENOL OXIDASE






• Molecule 1: CATALASE-PHENOL OXIDASE

Chain D: 86% 7% 6%




• Molecule 1: CATALASE-PHENOL OXIDASE

Chain E: 83% 9% 7%




• Molecule 1: CATALASE-PHENOL OXIDASE

Chain F: 84% 7% 7%





• Molecule 1: CATALASE-PHENOL OXIDASE

Chain G: 85% 8% 6%



● Molecule 1: CATALASE-PHENOL OXIDASE



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	253.37Å 243.29Å 97.06Å 90.00° 104.16° 90.00°	Depositor
Resolution (Å)	69.94 – 1.92 69.94 – 1.92	Depositor EDS
% Data completeness (in resolution range)	99.5 (69.94-1.92) 99.6 (69.94-1.92)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 1.92Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.164 , 0.201 0.163 , 0.199	Depositor DCC
R_{free} test set	21570 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	14.7	Xtriage
Anisotropy	0.442	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	46567	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 48.76 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 8.2225e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, HDD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	2/5536 (0.0%)	0.47	0/7520
1	B	0.44	1/5544 (0.0%)	0.47	0/7533
1	C	0.44	0/5555	0.47	0/7547
1	D	0.45	1/5551 (0.0%)	0.48	0/7539
1	E	0.45	6/5509 (0.1%)	0.46	0/7487
1	F	0.45	2/5490 (0.0%)	0.47	0/7460
1	G	0.44	0/5563	0.48	0/7558
1	H	0.45	4/5498 (0.1%)	0.47	0/7470
All	All	0.45	16/44246 (0.0%)	0.47	0/60114

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	698	SER	C-OXT	5.59	1.33	1.23
1	F	209	TRP	CD2-CE2	5.34	1.47	1.41
1	B	209	TRP	CD2-CE2	5.15	1.47	1.41
1	F	641	TRP	CD2-CE2	5.09	1.47	1.41
1	A	641	TRP	CD2-CE2	5.05	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5357	0	5161	57	0
1	B	5364	0	5173	56	0
1	C	5375	0	5189	51	0
1	D	5368	0	5188	46	0
1	E	5324	0	5134	53	0
1	F	5314	0	5130	47	0
1	G	5374	0	5195	53	0
1	H	5325	0	5133	50	0
2	A	44	0	31	1	0
2	B	44	0	31	2	0
2	C	44	0	31	2	0
2	D	44	0	31	2	0
2	E	44	0	31	3	0
2	F	44	0	31	2	0
2	G	44	0	31	1	0
2	H	44	0	31	2	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
4	A	521	0	0	12	0
4	B	420	0	0	8	0
4	C	470	0	0	15	0
4	D	468	0	0	13	0
4	E	341	0	0	3	0
4	F	381	0	0	8	0
4	G	454	0	0	8	0
4	H	350	0	0	6	0
All	All	46567	0	41551	400	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 400 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:470[A]:THR:HG21	4:B:2340:HOH:O	1.31	1.25
4:A:2194:HOH:O	1:B:127[A]:ARG:O	1.55	1.24

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:470[A]:THR:HG21	4:A:2427:HOH:O	1.35	1.23
1:G:127[B]:ARG:O	4:G:2062:HOH:O	1.53	1.21
4:C:2186:HOH:O	1:E:127[A]:ARG:NH2	1.71	1.17

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	684/719 (95%)	663 (97%)	21 (3%)	0	100	100
1	B	688/719 (96%)	665 (97%)	23 (3%)	0	100	100
1	C	689/719 (96%)	668 (97%)	21 (3%)	0	100	100
1	D	688/719 (96%)	672 (98%)	16 (2%)	0	100	100
1	E	681/719 (95%)	660 (97%)	21 (3%)	0	100	100
1	F	678/719 (94%)	660 (97%)	18 (3%)	0	100	100
1	G	690/719 (96%)	672 (97%)	18 (3%)	0	100	100
1	H	680/719 (95%)	661 (97%)	19 (3%)	0	100	100
All	All	5478/5752 (95%)	5321 (97%)	157 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	575/596 (96%)	555 (96%)	20 (4%)	41	29
1	B	576/596 (97%)	566 (98%)	10 (2%)	66	61
1	C	577/596 (97%)	561 (97%)	16 (3%)	49	39
1	D	576/596 (97%)	564 (98%)	12 (2%)	59	52
1	E	572/596 (96%)	559 (98%)	13 (2%)	56	48
1	F	569/596 (96%)	550 (97%)	19 (3%)	43	32
1	G	578/596 (97%)	565 (98%)	13 (2%)	57	50
1	H	570/596 (96%)	552 (97%)	18 (3%)	44	33
All	All	4593/4768 (96%)	4472 (97%)	121 (3%)	56	42

5 of 121 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	470[B]	THR
1	E	561	SER
1	H	444	SER
1	D	491[B]	GLU
1	E	191	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 67 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	594	GLN
1	E	461	GLN
1	H	167	GLN
1	E	82	HIS
1	E	301	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 9 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HDD	A	900	1,4	39,52,52	2.00	9 (23%)	25,89,89	2.39	8 (32%)
2	HDD	B	900	1,4	39,52,52	1.96	8 (20%)	25,89,89	2.49	12 (48%)
2	HDD	C	900	1,4	39,52,52	1.93	8 (20%)	25,89,89	2.52	12 (48%)
2	HDD	D	900	1,4	39,52,52	2.02	8 (20%)	25,89,89	2.48	11 (44%)
2	HDD	E	900	1,4	39,52,52	1.99	9 (23%)	25,89,89	2.50	11 (44%)
2	HDD	F	900	1,4	39,52,52	1.91	8 (20%)	25,89,89	2.47	11 (44%)
2	HDD	G	900	1,4	39,52,52	1.93	8 (20%)	25,89,89	2.40	11 (44%)
2	HDD	H	900	1,4	39,52,52	1.97	8 (20%)	25,89,89	2.51	13 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HDD	A	900	1,4	-	0/3/89/89	0/1/9/9
2	HDD	B	900	1,4	-	0/3/89/89	0/1/9/9
2	HDD	C	900	1,4	-	0/3/89/89	0/1/9/9
2	HDD	D	900	1,4	-	0/3/89/89	0/1/9/9
2	HDD	E	900	1,4	-	0/3/89/89	0/1/9/9
2	HDD	F	900	1,4	-	0/3/89/89	0/1/9/9
2	HDD	G	900	1,4	-	0/3/89/89	0/1/9/9
2	HDD	H	900	1,4	-	0/3/89/89	0/1/9/9

The worst 5 of 66 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	900	HDD	CHD-C1D	-2.13	1.33	1.36
2	A	900	HDD	CHD-C1D	-2.11	1.33	1.36
2	G	900	HDD	C4C-CHD	2.49	1.46	1.40
2	D	900	HDD	C1C-CHC	2.51	1.46	1.40
2	E	900	HDD	C1C-CHC	2.55	1.46	1.40

The worst 5 of 89 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	900	HDD	C4A-C3A-C2A	-7.76	101.59	107.00
2	B	900	HDD	C4A-C3A-C2A	-7.49	101.79	107.00
2	F	900	HDD	C4A-C3A-C2A	-7.44	101.82	107.00
2	H	900	HDD	C4A-C3A-C2A	-7.23	101.96	107.00
2	D	900	HDD	C4A-C3A-C2A	-7.03	102.10	107.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	900	HDD	1	0
2	B	900	HDD	2	0
2	C	900	HDD	2	0
2	D	900	HDD	2	0
2	E	900	HDD	3	0
2	F	900	HDD	2	0
2	G	900	HDD	1	0
2	H	900	HDD	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	671/719 (93%)	-0.55	5 (0%) 87 89	7, 14, 30, 49	0
1	B	673/719 (93%)	-0.47	2 (0%) 93 94	8, 14, 26, 38	0
1	C	673/719 (93%)	-0.49	1 (0%) 95 95	7, 13, 26, 37	0
1	D	674/719 (93%)	-0.51	2 (0%) 93 94	6, 12, 24, 48	0
1	E	668/719 (92%)	-0.52	3 (0%) 92 93	7, 15, 31, 60	0
1	F	668/719 (92%)	-0.50	5 (0%) 87 89	6, 13, 30, 56	0
1	G	673/719 (93%)	-0.50	2 (0%) 93 94	7, 13, 25, 40	0
1	H	670/719 (93%)	-0.51	4 (0%) 89 91	8, 14, 32, 50	0
All	All	5370/5752 (93%)	-0.51	24 (0%) 92 93	6, 14, 28, 60	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	698	SER	4.8
1	C	618	ALA	4.4
1	A	618	ALA	4.1
1	D	698	SER	3.7
1	F	618	ALA	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	HDD	A	900	44/44	0.96	0.12	2.76	8,13,22,24	0
2	HDD	D	900	44/44	0.96	0.11	2.74	7,11,18,23	0
2	HDD	G	900	44/44	0.96	0.10	2.67	7,13,19,23	0
2	HDD	C	900	44/44	0.95	0.10	2.63	8,14,18,21	0
2	HDD	H	900	44/44	0.95	0.11	2.17	9,13,22,24	0
2	HDD	B	900	44/44	0.96	0.10	2.13	10,14,19,22	0
2	HDD	E	900	44/44	0.96	0.11	1.92	9,14,22,28	0
2	HDD	F	900	44/44	0.96	0.10	1.38	7,12,17,22	0
3	CA	B	1700	1/1	1.00	0.12	-	10,10,10,10	1
3	CA	A	1701	1/1	1.00	0.08	-	8,8,8,8	1
3	CA	G	1700	1/1	0.99	0.09	-	12,12,12,12	1
3	CA	D	1700	1/1	0.99	0.09	-	9,9,9,9	1
3	CA	A	1700	1/1	0.98	0.08	-	19,19,19,19	1
3	CA	F	1700	1/1	1.00	0.09	-	8,8,8,8	1
3	CA	C	1700	1/1	0.99	0.09	-	10,10,10,10	1
3	CA	E	1700	1/1	0.98	0.08	-	8,8,8,8	1
3	CA	H	1700	1/1	1.00	0.09	-	7,7,7,7	1

6.5 Other polymers [i](#)

There are no such residues in this entry.