



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 02:08 am GMT

PDB ID : 2AV7  
Title : Crystal structure of HTLV-1 TAX peptide Bound to Human Class I MHC HLA-A2 with the K66A mutation in the heavy chain.  
Authors : Borbulevych, O.Y.; Baker, B.M.  
Deposited on : 2005-08-29  
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1316 (2.04-2.04)
Clashscore	112137	1394 (2.04-2.04)
Ramachandran outliers	110173	1383 (2.04-2.04)
Sidechain outliers	110143	1383 (2.04-2.04)
RSRZ outliers	101464	1319 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	<div> <div>86%</div> <div>12%</div> <div>•</div> </div>
1	D	275	<div> <div>90%</div> <div>8%</div> <div>•</div> </div>
2	B	100	<div> <div>75%</div> <div>23%</div> <div>•</div> </div>
2	E	100	<div> <div>88%</div> <div>12%</div> </div>
3	C	9	<div> <div>89%</div> <div>11%</div> </div>
3	F	9	<div> <div>78%</div> <div>22%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	701	-	-	-	X
4	GOL	A	702	-	-	-	X
4	GOL	A	703	-	-	-	X
4	GOL	A	704	-	-	-	X
4	GOL	A	719	-	-	-	X
4	GOL	A	720	-	-	-	X
4	GOL	B	705	-	-	-	X
4	GOL	B	712	-	-	-	X
4	GOL	B	713	-	-	-	X
4	GOL	B	716	-	-	-	X
4	GOL	B	718	-	-	-	X
4	GOL	D	708	-	-	-	X
4	GOL	D	714	-	-	-	X
4	GOL	D	715	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7135 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A-2 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	0	0
			2243	1400	408	426	9			
1	D	275	Total	C	N	O	S	0	0	0
			2243	1400	408	426	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	66	ALA	LYS	ENGINEERED	UNP P01892
D	66	ALA	LYS	ENGINEERED	UNP P01892

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			
2	E	100	Total	C	N	O	S	0	1	0
			842	535	144	159	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	INITIATING METHIONINE	UNP P61769
E	0	MET	-	INITIATING METHIONINE	UNP P61769

- Molecule 3 is a protein called Trans-activating transcriptional regulatory peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O		0	0	0
			77	56	9	12				
3	F	9	Total	C	N	O		0	0	0
			77	56	9	12				

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0
4	E	1	Total C O 6 3 3	0	0
4	F	1	Total C O 6 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	223	Total O 224 224	0	1
5	B	99	Total O 99 99	0	0
5	C	8	Total O 8 8	0	0
5	D	251	Total O 251 251	0	0
5	E	106	Total O 106 106	0	0
5	F	8	Total O 8 8	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class I histocompatibility antigen, A-2 alpha chain




- Molecule 1: HLA class I histocompatibility antigen, A-2 alpha chain




- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin




- Molecule 3: Trans-activating transcriptional regulatory peptide



- Molecule 3: Trans-activating transcriptional regulatory peptide

Chain F:  78% 22%





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.32Å 62.53Å 74.80Å 82.13° 76.57° 78.06°	Depositor
Resolution (Å)	10.00 – 2.05 9.99 – 2.05	Depositor EDS
% Data completeness (in resolution range)	96.0 (10.00-2.05) 93.0 (9.99-2.05)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.96 (at 2.06Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.171 , 0.234 0.173 , 0.234	Depositor DCC
$R_{free}$ test set	2612 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	20.4	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 53.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7135	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:  
GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.89	0/2308	1.05	12/3133 (0.4%)
1	D	0.94	3/2308 (0.1%)	1.08	9/3133 (0.3%)
2	B	0.94	0/860	1.10	6/1162 (0.5%)
2	E	0.98	0/871	0.99	0/1176
3	C	0.98	0/80	0.81	0/108
3	F	1.03	0/80	0.87	0/108
All	All	0.93	3/6507 (0.0%)	1.06	27/8820 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	264	GLU	CD-OE1	7.16	1.33	1.25
1	D	189	MET	SD-CE	-6.63	1.40	1.77
1	D	95	VAL	CB-CG2	-5.25	1.41	1.52

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	29	ASP	CB-CG-OD2	9.29	126.66	118.30
1	A	30	ASP	CB-CG-OD2	9.07	126.47	118.30
1	D	30	ASP	CB-CG-OD2	7.65	125.18	118.30
1	A	220	ASP	CB-CG-OD2	7.43	124.99	118.30
1	A	39	ASP	CB-CG-OD2	7.30	124.88	118.30
1	A	183	ASP	CB-CG-OD2	7.10	124.69	118.30
1	A	29	ASP	CB-CG-OD2	6.99	124.59	118.30
1	A	273	ARG	NE-CZ-NH1	-6.77	116.91	120.30
1	D	137	ASP	CB-CG-OD2	6.74	124.37	118.30
1	A	122	ASP	CB-CG-OD2	6.64	124.27	118.30
1	A	119	ASP	CB-CG-OD2	6.55	124.19	118.30
1	D	75	ARG	NE-CZ-NH1	6.42	123.51	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	129	ASP	CB-CG-OD1	6.04	123.73	118.30
2	B	85	VAL	CB-CA-C	-6.00	100.01	111.40
1	D	82	ARG	NE-CZ-NH2	5.75	123.18	120.30
1	A	230	LEU	CA-CB-CG	5.75	128.53	115.30
2	B	99	MET	CA-C-O	5.62	131.91	120.10
1	D	238	ASP	CB-CG-OD2	5.60	123.34	118.30
1	D	225	THR	OG1-CB-CG2	-5.56	97.21	110.00
1	A	102	ASP	CB-CG-OD2	5.56	123.30	118.30
2	B	96	ASP	CB-CG-OD1	5.33	123.10	118.30
2	B	59	ASP	CB-CG-OD1	5.28	123.06	118.30
2	B	76	ASP	CB-CG-OD2	5.28	123.05	118.30
1	A	106	ASP	CB-CG-OD2	5.22	123.00	118.30
1	D	129	ASP	CB-CG-OD1	5.14	122.92	118.30
1	D	102	ASP	CB-CG-OD2	5.10	122.89	118.30
2	B	34	ASP	CB-CG-OD1	5.03	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2243	0	2088	13	0
1	D	2243	0	2088	6	0
2	B	837	0	803	13	0
2	E	842	0	808	6	0
3	C	77	0	79	1	0
3	F	77	0	79	1	0
4	A	42	0	56	5	0
4	B	42	0	56	8	0
4	D	24	0	32	4	0
4	E	6	0	8	0	0
4	F	6	0	8	0	0
5	A	224	0	0	0	0
5	B	99	0	0	1	0
5	C	8	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	251	0	0	0	0
5	E	106	0	0	0	0
5	F	8	0	0	0	0
All	All	7135	0	6105	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:702:GOL:H11	3:C:6:PRO:HA	1.73	0.70
1:A:196:ASP:N	1:A:196:ASP:OD1	2.24	0.69
2:B:19:LYS:HB2	4:B:705:GOL:H31	1.79	0.63
2:B:46:ILE:HG12	4:B:713:GOL:H12	1.79	0.63
4:D:709:GOL:H2	2:E:32:PRO:HB3	1.87	0.55
2:E:3[B]:ARG:NH2	2:E:59:ASP:OD2	2.39	0.55
1:A:244:TRP:HE1	4:B:707:GOL:H11	1.72	0.53
1:A:30:ASP:OD1	4:A:701:GOL:H31	2.08	0.53
1:A:137:ASP:O	1:A:141:GLN:NE2	2.34	0.53
2:B:8:GLN:HA	4:B:707:GOL:H32	1.89	0.53
1:D:94:THR:OG1	4:D:709:GOL:H31	2.07	0.53
2:B:11:SER:O	4:B:706:GOL:H11	2.09	0.52
2:B:10:TYR:HB2	4:B:706:GOL:H12	1.92	0.52
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.44	0.52
1:D:218:GLN:HE21	1:D:221:GLY:HA2	1.75	0.51
1:D:117:ALA:HB2	2:E:60:TRP:CE2	2.46	0.50
1:A:70:HIS:CE1	4:A:702:GOL:H31	2.46	0.50
4:D:715:GOL:H11	2:E:13:HIS:HA	1.93	0.50
2:B:83:ASN:ND2	5:B:744:HOH:O	2.45	0.49
2:E:25:CYS:HB2	2:E:39:LEU:HD21	1.95	0.49
1:A:28:VAL:HG11	1:A:179:LEU:HD13	1.95	0.48
2:B:41:LYS:HD3	4:B:713:GOL:H11	1.96	0.48
1:A:110:LEU:HD13	4:A:703:GOL:H2	1.97	0.46
1:A:181:ARG:NE	1:A:183:ASP:OD2	2.49	0.46
1:D:178:THR:O	1:D:181:ARG:HD3	2.17	0.45
1:D:30:ASP:HA	4:D:708:GOL:H31	1.97	0.45
1:A:144:LYS:HE3	4:A:720:GOL:H31	2.01	0.43
2:E:36:GLU:HB3	2:E:83:ASN:HB2	2.01	0.43
1:A:176:LYS:HE3	1:A:176:LYS:HB3	1.60	0.43
1:A:215:LEU:HD13	1:A:243:LYS:HD3	2.01	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:22:PHE:CE2	2:B:69:GLU:HG3	2.55	0.42
2:B:24:ASN:HB3	2:B:65:LEU:HD11	2.02	0.41
2:B:41:LYS:HD3	4:B:713:GOL:H32	2.02	0.41
2:B:39:LEU:HA	2:B:39:LEU:HD23	1.87	0.41
1:A:11:SER:HB2	1:A:74:HIS:CD2	2.55	0.41
3:F:5:TYR:HA	3:F:6:PRO:HD3	1.96	0.41
1:D:196:ASP:OD2	1:D:196:ASP:N	2.53	0.41
2:B:83:ASN:HD22	2:B:84:HIS:H	1.69	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/275 (99%)	268 (98%)	5 (2%)	0	100	100
1	D	273/275 (99%)	266 (97%)	7 (3%)	0	100	100
2	B	98/100 (98%)	95 (97%)	2 (2%)	1 (1%)	18	8
2	E	99/100 (99%)	98 (99%)	1 (1%)	0	100	100
3	C	7/9 (78%)	7 (100%)	0	0	100	100
3	F	7/9 (78%)	7 (100%)	0	0	100	100
All	All	757/768 (99%)	741 (98%)	15 (2%)	1 (0%)	55	47

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	97	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/230 (100%)	216 (94%)	14 (6%)	22	12
1	D	230/230 (100%)	217 (94%)	13 (6%)	24	14
2	B	95/95 (100%)	89 (94%)	6 (6%)	21	11
2	E	96/95 (101%)	93 (97%)	3 (3%)	45	38
3	C	8/8 (100%)	8 (100%)	0	100	100
3	F	8/8 (100%)	8 (100%)	0	100	100
All	All	667/666 (100%)	631 (95%)	36 (5%)	26	16

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	SER
1	A	11	SER
1	A	14	ARG
1	A	35	ARG
1	A	74	HIS
1	A	75	ARG
1	A	88	SER
1	A	89	GLU
1	A	157	ARG
1	A	176	LYS
1	A	196	ASP
1	A	207	SER
1	A	255	GLN
1	A	268	LYS
2	B	48	LYS
2	B	70	PHE
2	B	83	ASN
2	B	85	VAL
2	B	89	GLN
2	B	98	ASP
1	D	35	ARG
1	D	74	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	88	SER
1	D	110	LEU
1	D	111	ARG
1	D	137	ASP
1	D	176	LYS
1	D	177	GLU
1	D	181	ARG
1	D	196	ASP
1	D	207	SER
1	D	225	THR
1	D	268	LYS
2	E	58	LYS
2	E	70	PHE
2	E	75	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	HIS
1	A	174	ASN
2	B	2	GLN
2	B	83	ASN
1	D	218	GLN
2	E	2	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	A	701	-	5,5,5	0.23	0	5,5,5	0.71	0
4	GOL	A	702	-	5,5,5	0.55	0	5,5,5	0.54	0
4	GOL	A	703	-	5,5,5	0.51	0	5,5,5	1.08	0
4	GOL	A	704	-	5,5,5	0.57	0	5,5,5	0.39	0
4	GOL	A	717	-	5,5,5	0.39	0	5,5,5	0.27	0
4	GOL	A	719	-	5,5,5	0.46	0	5,5,5	0.52	0
4	GOL	A	720	-	5,5,5	0.54	0	5,5,5	0.95	0
4	GOL	B	705	-	5,5,5	0.45	0	5,5,5	1.27	1 (20%)
4	GOL	B	706	-	5,5,5	0.59	0	5,5,5	1.36	1 (20%)
4	GOL	B	707	-	5,5,5	0.60	0	5,5,5	1.03	0
4	GOL	B	712	-	5,5,5	0.44	0	5,5,5	0.80	0
4	GOL	B	713	-	5,5,5	0.53	0	5,5,5	1.24	1 (20%)
4	GOL	B	716	-	5,5,5	0.49	0	5,5,5	0.30	0
4	GOL	B	718	-	5,5,5	0.45	0	5,5,5	0.73	0
4	GOL	D	708	-	5,5,5	0.35	0	5,5,5	0.77	0
4	GOL	D	709	-	5,5,5	0.96	1 (20%)	5,5,5	0.80	0
4	GOL	D	714	-	5,5,5	0.37	0	5,5,5	0.87	0
4	GOL	D	715	-	5,5,5	0.40	0	5,5,5	1.11	0
4	GOL	E	711	-	5,5,5	0.13	0	5,5,5	0.92	0
4	GOL	F	710	-	5,5,5	0.35	0	5,5,5	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	701	-	-	0/4/4/4	0/0/0/0
4	GOL	A	702	-	-	0/4/4/4	0/0/0/0
4	GOL	A	703	-	-	0/4/4/4	0/0/0/0
4	GOL	A	704	-	-	0/4/4/4	0/0/0/0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	717	-	-	0/4/4/4	0/0/0/0
4	GOL	A	719	-	-	0/4/4/4	0/0/0/0
4	GOL	A	720	-	-	0/4/4/4	0/0/0/0
4	GOL	B	705	-	-	0/4/4/4	0/0/0/0
4	GOL	B	706	-	-	0/4/4/4	0/0/0/0
4	GOL	B	707	-	-	0/4/4/4	0/0/0/0
4	GOL	B	712	-	-	0/4/4/4	0/0/0/0
4	GOL	B	713	-	-	0/4/4/4	0/0/0/0
4	GOL	B	716	-	-	0/4/4/4	0/0/0/0
4	GOL	B	718	-	-	0/4/4/4	0/0/0/0
4	GOL	D	708	-	-	0/4/4/4	0/0/0/0
4	GOL	D	709	-	-	0/4/4/4	0/0/0/0
4	GOL	D	714	-	-	0/4/4/4	0/0/0/0
4	GOL	D	715	-	-	0/4/4/4	0/0/0/0
4	GOL	E	711	-	-	0/4/4/4	0/0/0/0
4	GOL	F	710	-	-	0/4/4/4	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	709	GOL	O2-C2	-2.05	1.37	1.43

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	713	GOL	O1-C1-C2	2.14	120.86	110.07
4	B	705	GOL	O3-C3-C2	2.53	122.81	110.07
4	B	706	GOL	O1-C1-C2	2.74	123.85	110.07

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

11 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	701	GOL	1	0
4	A	702	GOL	2	0
4	A	703	GOL	1	0
4	A	720	GOL	1	0
4	B	705	GOL	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	706	GOL	2	0
4	B	707	GOL	2	0
4	B	713	GOL	3	0
4	D	708	GOL	1	0
4	D	709	GOL	2	0
4	D	715	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	275/275 (100%)	-0.39	3 (1%) 80 83	5, 11, 21, 29	0
1	D	275/275 (100%)	-0.41	3 (1%) 80 83	5, 11, 20, 31	0
2	B	100/100 (100%)	-0.39	2 (2%) 65 70	5, 11, 21, 41	0
2	E	100/100 (100%)	-0.59	0 100 100	5, 11, 20, 30	0
3	C	9/9 (100%)	0.02	0 100 100	9, 11, 14, 14	0
3	F	9/9 (100%)	0.89	0 100 100	10, 13, 15, 19	0
All	All	768/768 (100%)	-0.40	8 (1%) 82 85	5, 11, 21, 41	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	99	MET	4.6
2	B	98	ASP	4.0
1	D	196	ASP	3.1
1	D	17	ARG	2.7
1	A	194	VAL	2.6
1	A	196	ASP	2.6
1	D	194	VAL	2.2
1	A	192	HIS	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	A	703	6/6	0.77	0.27	8.68	37,41,44,45	0
4	GOL	B	716	6/6	0.88	0.16	7.32	35,40,41,43	6
4	GOL	D	708	6/6	0.87	0.17	7.03	32,44,45,53	0
4	GOL	D	715	6/6	0.80	0.16	6.40	37,46,48,49	0
4	GOL	A	701	6/6	0.91	0.11	5.67	41,44,45,47	0
4	GOL	A	720	6/6	0.74	0.28	4.93	43,51,52,53	0
4	GOL	B	712	6/6	0.74	0.24	4.30	44,47,47,49	0
4	GOL	B	718	6/6	0.83	0.26	3.98	30,31,31,32	6
4	GOL	A	702	6/6	0.91	0.20	3.62	40,41,45,45	0
4	GOL	B	705	6/6	0.95	0.18	3.39	26,37,42,42	0
4	GOL	D	714	6/6	0.86	0.18	3.12	41,42,44,48	0
4	GOL	A	704	6/6	0.81	0.28	3.04	37,44,44,45	0
4	GOL	A	719	6/6	0.86	0.19	3.02	40,41,43,43	6
4	GOL	B	713	6/6	0.83	0.30	2.80	49,49,50,51	0
4	GOL	D	709	6/6	0.91	0.18	1.93	32,40,41,43	0
4	GOL	F	710	6/6	0.83	0.20	1.79	46,50,51,53	0
4	GOL	B	706	6/6	0.64	0.21	1.31	39,44,45,47	0
4	GOL	A	717	6/6	0.87	0.14	1.19	34,37,38,40	6
4	GOL	E	711	6/6	0.91	0.13	0.87	31,35,36,40	0
4	GOL	B	707	6/6	0.92	0.18	0.74	25,37,41,42	0

## 6.5 Other polymers

There are no such residues in this entry.