



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 04:55 am GMT

PDB ID : 3AVJ  
Title : Crystal structures of novel allosteric peptide inhibitors of HIV integrase in the LEDGF binding site  
Authors : Peat, T.S.; Deadman, J.J.; Newman, J.; Rhodes, D.I.  
Deposited on : 2011-03-05  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3453 (1.70-1.70)
Clashscore	112137	3876 (1.70-1.70)
Ramachandran outliers	110173	3815 (1.70-1.70)
Sidechain outliers	110143	3815 (1.70-1.70)
RSRZ outliers	101464	3491 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	183	
1	B	183	
2	D	8	
2	F	8	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	5	-	-	-	X
3	SO4	B	6	-	-	-	X
5	ACY	A	3	-	-	-	X
5	ACY	B	2	-	-	X	X
5	ACY	B	214	-	-	X	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2902 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	149	Total	C	N	O	S	0	17	0
			1270	817	214	235	4			
1	B	149	Total	C	N	O	S	0	17	0
			1270	813	217	236	4			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	30	MET	-	EXPRESSION TAG	UNP P12497
A	31	GLY	-	EXPRESSION TAG	UNP P12497
A	32	SER	-	EXPRESSION TAG	UNP P12497
A	33	SER	-	EXPRESSION TAG	UNP P12497
A	34	HIS	-	EXPRESSION TAG	UNP P12497
A	35	HIS	-	EXPRESSION TAG	UNP P12497
A	36	HIS	-	EXPRESSION TAG	UNP P12497
A	37	HIS	-	EXPRESSION TAG	UNP P12497
A	38	HIS	-	EXPRESSION TAG	UNP P12497
A	39	HIS	-	EXPRESSION TAG	UNP P12497
A	40	SER	-	EXPRESSION TAG	UNP P12497
A	41	SER	-	EXPRESSION TAG	UNP P12497
A	42	GLY	-	EXPRESSION TAG	UNP P12497
A	43	LEU	-	EXPRESSION TAG	UNP P12497
A	44	VAL	-	EXPRESSION TAG	UNP P12497
A	45	PRO	-	EXPRESSION TAG	UNP P12497
A	46	ARG	-	EXPRESSION TAG	UNP P12497
A	47	GLY	-	EXPRESSION TAG	UNP P12497
A	48	SER	-	EXPRESSION TAG	UNP P12497
A	49	HIS	-	EXPRESSION TAG	UNP P12497
A	56	SER	CYS	ENGINEERED MUTATION	UNP P12497
A	139	ASP	PHE	ENGINEERED MUTATION	UNP P12497
A	185	HIS	PHE	ENGINEERED MUTATION	UNP P12497
B	30	MET	-	EXPRESSION TAG	UNP P12497
B	31	GLY	-	EXPRESSION TAG	UNP P12497

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	32	SER	-	EXPRESSION TAG	UNP P12497
B	33	SER	-	EXPRESSION TAG	UNP P12497
B	34	HIS	-	EXPRESSION TAG	UNP P12497
B	35	HIS	-	EXPRESSION TAG	UNP P12497
B	36	HIS	-	EXPRESSION TAG	UNP P12497
B	37	HIS	-	EXPRESSION TAG	UNP P12497
B	38	HIS	-	EXPRESSION TAG	UNP P12497
B	39	HIS	-	EXPRESSION TAG	UNP P12497
B	40	SER	-	EXPRESSION TAG	UNP P12497
B	41	SER	-	EXPRESSION TAG	UNP P12497
B	42	GLY	-	EXPRESSION TAG	UNP P12497
B	43	LEU	-	EXPRESSION TAG	UNP P12497
B	44	VAL	-	EXPRESSION TAG	UNP P12497
B	45	PRO	-	EXPRESSION TAG	UNP P12497
B	46	ARG	-	EXPRESSION TAG	UNP P12497
B	47	GLY	-	EXPRESSION TAG	UNP P12497
B	48	SER	-	EXPRESSION TAG	UNP P12497
B	49	HIS	-	EXPRESSION TAG	UNP P12497
B	56	SER	CYS	ENGINEERED MUTATION	UNP P12497
B	139	ASP	PHE	ENGINEERED MUTATION	UNP P12497
B	185	HIS	PHE	ENGINEERED MUTATION	UNP P12497

- Molecule 2 is a protein called LEDGF peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	8	Total	C	N	O	S	0	0	0
			62	38	10	13	1			
2	F	8	Total	C	N	O	S	0	0	0
			62	38	10	13	1			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is ACETIC ACID (three-letter code: ACY) (formula: C<sub>2</sub>H<sub>4</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	90	Total	O	0	0
			90	90		
6	B	86	Total	O	0	0
			86	86		
6	D	4	Total	O	0	0
			4	4		
6	F	4	Total	O	0	0
			4	4		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Integrase



- Molecule 1: Integrase



- Molecule 2: LEDGF peptide



There are no outlier residues recorded for this chain.

- Molecule 2: LEDGF peptide



There are no outlier residues recorded for this chain.



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.02Å 71.02Å 67.02Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	61.50 – 1.70 61.50 – 1.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (61.50-1.70) 100.0 (61.50-1.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.94 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.6.0062	Depositor
R, $R_{free}$	0.158 , 0.198 0.155 , 0.196	Depositor DCC
$R_{free}$ test set	2090 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.8	Xtriage
Anisotropy	0.072	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.020 for -h,-k,l 0.477 for h,-h-k,-l 0.021 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2902	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.55% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACY, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.47	8/1318 (0.6%)	1.21	2/1788 (0.1%)
1	B	1.48	9/1319 (0.7%)	1.21	3/1787 (0.2%)
2	D	1.12	0/61	1.18	0/80
2	F	1.33	0/61	1.20	0/80
All	All	1.46	17/2759 (0.6%)	1.21	5/3735 (0.1%)

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	131	TRP	CZ3-CH2	6.93	1.51	1.40
1	B	72	VAL	CB-CG1	-6.83	1.38	1.52
1	A	157	GLU	CG-CD	6.43	1.61	1.51
1	A	81	SER	CB-OG	6.29	1.50	1.42
1	B	79	VAL	CB-CG2	6.22	1.66	1.52
1	A	72	VAL	CB-CG1	-6.17	1.40	1.52
1	A	100	PHE	CD1-CE1	6.10	1.51	1.39
1	B	100	PHE	CD1-CE1	5.95	1.51	1.39
1	B	131	TRP	CZ3-CH2	5.89	1.49	1.40
1	A	138	GLU	CD-OE2	-5.54	1.19	1.25
1	B	110	VAL	CB-CG1	5.36	1.64	1.52
1	B	133	ALA	CA-CB	5.28	1.63	1.52
1	A	132	TRP	CE3-CZ3	5.23	1.47	1.38
1	B	113[A]	VAL	CB-CG1	5.17	1.63	1.52
1	B	113[B]	VAL	CB-CG1	5.17	1.63	1.52
1	A	179	ALA	CA-CB	5.13	1.63	1.52
1	B	131	TRP	CD2-CE2	5.01	1.47	1.41

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	116	ASP	CB-CG-OD1	5.66	123.39	118.30
1	A	83	TYR	CB-CG-CD1	-5.39	117.77	121.00
1	A	139	ASP	CB-CG-OD2	-5.29	113.54	118.30
1	B	71	LYS	CD-CE-NZ	-5.09	100.00	111.70
1	B	173	LYS	CD-CE-NZ	-5.08	100.02	111.70

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1270	0	1302	31	0
1	B	1270	0	1292	34	0
2	D	62	0	63	0	0
2	F	62	0	63	0	0
3	A	20	0	0	1	0
3	B	20	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	4	0	3	0	0
5	B	8	0	6	14	0
6	A	90	0	0	2	0
6	B	86	0	0	2	0
6	D	4	0	0	0	0
6	F	4	0	0	0	0
All	All	2902	0	2729	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87[B]:GLU:OE1	6:A:19:HOH:O	1.65	1.14

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141[B]:ILE:HD12	1:A:142:PRO:HD2	1.29	1.12
1:B:87[B]:GLU:OE1	6:B:269:HOH:O	1.65	1.11
1:A:139:ASP:OD1	1:A:141[B]:ILE:HG22	1.68	0.93
5:B:214:ACY:O	5:B:2:ACY:H3	1.70	0.91
1:B:185:HIS:CD2	5:B:214:ACY:H3	2.07	0.90
1:A:141[B]:ILE:HG23	1:A:143:TYR:CE1	2.09	0.88
1:A:136[A]:LYS:NZ	1:A:138:GLU:OE1	2.11	0.81
1:A:202[B]:ASP:OD1	1:B:209:GLN:NE2	2.14	0.79
5:B:214:ACY:O	5:B:2:ACY:CH3	2.32	0.78
1:A:187:ARG:HH11	1:A:187:ARG:HG2	1.51	0.74
1:B:145:PRO:HA	1:B:148:GLN:HE22	1.53	0.74
1:A:103[B]:LYS:NZ	1:B:87[B]:GLU:OE2	2.25	0.68
1:B:197:GLY:HA3	5:B:2:ACY:H1	1.76	0.68
1:B:57[A]:SER:HB3	1:B:58:PRO:HD3	1.76	0.67
1:A:87[B]:GLU:OE2	1:B:103[B]:LYS:NZ	2.28	0.66
1:B:142:PRO:HB2	1:B:148:GLN:NE2	2.10	0.66
1:B:167[A]:ASP:OD1	6:B:252:HOH:O	2.13	0.66
5:B:214:ACY:C	5:B:2:ACY:CH3	2.74	0.66
1:B:185:HIS:NE2	5:B:214:ACY:H3	2.11	0.65
1:A:141[B]:ILE:HD12	1:A:142:PRO:CD	2.17	0.64
1:A:170[B]:GLU:OE1	6:A:251:HOH:O	2.16	0.61
5:B:214:ACY:C	5:B:2:ACY:H3	2.31	0.60
1:A:187:ARG:HG2	1:A:187:ARG:NH1	2.16	0.60
5:B:214:ACY:C	5:B:2:ACY:H1	2.32	0.59
1:A:202[B]:ASP:CG	1:B:209:GLN:HE22	2.05	0.59
1:A:57[A]:SER:HB3	1:A:58:PRO:CD	2.32	0.58
1:A:160:LYS:NZ	1:A:164:GLN:NE2	2.52	0.58
1:B:205:ALA:O	1:B:209:GLN:HG2	2.04	0.58
3:A:7:SO4:O1	1:B:103[B]:LYS:NZ	2.37	0.57
1:B:145:PRO:O	1:B:148:GLN:NE2	2.39	0.56
1:A:87[B]:GLU:OE2	1:B:103[B]:LYS:HE3	2.06	0.55
1:B:181:PHE:O	1:B:185:HIS:HD2	1.89	0.55
1:A:141[B]:ILE:HG23	1:A:143:TYR:CZ	2.43	0.53
1:A:57[A]:SER:HB3	1:A:58:PRO:HD3	1.90	0.53
1:B:57[A]:SER:CB	1:B:58:PRO:HD3	2.39	0.52
1:A:87[B]:GLU:OE2	1:B:103[B]:LYS:CE	2.57	0.51
1:B:185:HIS:CG	5:B:214:ACY:CH3	2.95	0.49
1:A:57[A]:SER:CB	1:A:58:PRO:CD	2.91	0.48
1:A:103[B]:LYS:HG2	1:B:177:GLN:HG3	1.94	0.48
1:A:127:LYS:NZ	1:A:137:GLN:HE22	2.11	0.48
1:A:141[B]:ILE:CG2	1:A:143:TYR:CZ	2.97	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:185:HIS:CD2	5:B:214:ACY:CH3	2.90	0.48
1:B:185:HIS:CE1	5:B:214:ACY:CH3	2.97	0.48
1:B:63[B]:LEU:HD11	1:B:74:LEU:HD11	1.96	0.47
1:A:187:ARG:CG	1:A:187:ARG:NH1	2.73	0.47
1:B:185:HIS:CG	5:B:214:ACY:H3	2.48	0.47
1:A:141[B]:ILE:CG2	1:A:143:TYR:CE1	2.92	0.45
1:A:187:ARG:HH11	1:A:187:ARG:CG	2.17	0.45
1:A:139:ASP:CG	1:A:141[B]:ILE:HG22	2.36	0.44
1:A:160:LYS:HZ2	1:A:164:GLN:NE2	2.15	0.44
1:B:57[A]:SER:CB	1:B:58:PRO:CD	2.98	0.42
1:B:142:PRO:CB	1:B:148:GLN:NE2	2.82	0.42
1:B:197:GLY:HA3	5:B:2:ACY:CH3	2.47	0.41
1:B:200:ILE:O	1:B:204[B]:ILE:HG12	2.21	0.41
1:A:137:GLN:HE21	1:A:137:GLN:HB3	1.70	0.41
1:A:65:CYS:SG	1:A:72:VAL:HG13	2.61	0.41
1:B:185:HIS:ND1	5:B:214:ACY:H1	2.37	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	161/183 (88%)	160 (99%)	1 (1%)	0	100	100
1	B	161/183 (88%)	159 (99%)	2 (1%)	0	100	100
2	D	6/8 (75%)	5 (83%)	1 (17%)	0	100	100
2	F	6/8 (75%)	5 (83%)	1 (17%)	0	100	100
All	All	334/382 (87%)	329 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	140/150 (93%)	138 (99%)	2 (1%)	71	58
1	B	140/150 (93%)	139 (99%)	1 (1%)	87	81
2	D	7/7 (100%)	7 (100%)	0	100	100
2	F	7/7 (100%)	7 (100%)	0	100	100
All	All	294/314 (94%)	291 (99%)	3 (1%)	75	71

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	167	ASP
1	A	188	LYS
1	B	148	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	137	GLN
1	A	164	GLN
1	B	146	GLN
1	B	148	GLN
1	B	185	HIS
1	B	209	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 2 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	2	-	4,4,4	0.18	0	6,6,6	1.24	0
5	ACY	A	3	-	1,3,3	1.12	0	0,3,3	0.00	-
3	SO4	A	4	-	4,4,4	0.22	0	6,6,6	0.28	0
3	SO4	A	5	-	4,4,4	0.42	0	6,6,6	0.55	0
3	SO4	A	7	-	4,4,4	0.49	0	6,6,6	1.79	2 (33%)
3	SO4	B	1	-	4,4,4	0.30	0	6,6,6	0.90	0
5	ACY	B	2	-	1,3,3	1.89	0	0,3,3	0.00	-
5	ACY	B	214	-	1,3,3	1.95	0	0,3,3	0.00	-
3	SO4	B	3	-	4,4,4	0.19	0	6,6,6	0.36	0
3	SO4	B	6	-	4,4,4	0.39	0	6,6,6	0.57	0
3	SO4	B	8	-	4,4,4	0.57	0	6,6,6	1.03	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	2	-	-	0/0/0/0	0/0/0/0
5	ACY	A	3	-	-	0/0/0/0	0/0/0/0
3	SO4	A	4	-	-	0/0/0/0	0/0/0/0
3	SO4	A	5	-	-	0/0/0/0	0/0/0/0
3	SO4	A	7	-	-	0/0/0/0	0/0/0/0
3	SO4	B	1	-	-	0/0/0/0	0/0/0/0
5	ACY	B	2	-	-	0/0/0/0	0/0/0/0
5	ACY	B	214	-	-	0/0/0/0	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	B	3	-	-	0/0/0/0	0/0/0/0
3	SO4	B	6	-	-	0/0/0/0	0/0/0/0
3	SO4	B	8	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	7	SO4	O3-S-O2	-3.06	92.39	109.26
3	B	8	SO4	O4-S-O1	-2.03	98.07	109.26
3	A	7	SO4	O4-S-O3	2.71	121.16	108.96

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	7	SO4	1	0
5	B	2	ACY	7	0
5	B	214	ACY	12	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	149/183 (81%)	-0.61	0 100 100	12, 18, 41, 51	16 (10%)
1	B	149/183 (81%)	-0.62	0 100 100	12, 18, 40, 58	15 (10%)
2	D	8/8 (100%)	-0.83	0 100 100	17, 22, 28, 29	0
2	F	8/8 (100%)	-0.80	0 100 100	18, 21, 27, 28	0
All	All	314/382 (82%)	-0.63	0 100 100	12, 18, 41, 58	31 (9%)

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	ACY	B	214	4/4	0.94	0.13	6.12	40,46,52,55	0
3	SO4	B	6	5/5	0.99	0.12	5.60	12,16,17,19	5

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	ACY	A	3	4/4	0.93	0.12	5.53	40,44,50,56	0
3	SO4	A	5	5/5	0.99	0.13	5.50	13,16,18,19	5
5	ACY	B	2	4/4	0.93	0.10	2.48	25,31,33,33	4
3	SO4	B	3	5/5	0.98	0.09	1.23	27,35,38,46	5
3	SO4	B	8	5/5	0.97	0.09	0.34	23,31,36,39	5
3	SO4	A	4	5/5	0.98	0.08	0.32	22,31,31,41	5
3	SO4	A	2	5/5	0.99	0.07	-0.10	27,31,41,48	5
3	SO4	B	1	5/5	0.99	0.06	-0.46	27,29,43,47	5
3	SO4	A	7	5/5	0.97	0.06	-1.73	25,34,38,38	5
4	CL	A	213	1/1	0.96	0.06	-	48,48,48,48	0
4	CL	B	213	1/1	0.98	0.05	-	49,49,49,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.