



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 06:09 am GMT

PDB ID : 5AXC  
Title : Crystal structure of mouse SAHH complexed with 3'-keto aristeromycin  
Authors : Kusakabe, Y.; Ishihara, M.; Tanaka, N.  
Deposited on : 2015-07-24  
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1088 (1.56-1.56)
Clashscore	112137	1132 (1.56-1.56)
Ramachandran outliers	110173	1110 (1.56-1.56)
Sidechain outliers	110143	1108 (1.56-1.56)
RSRZ outliers	101464	1089 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	432	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>10%</div> <div>••</div> </div> </div>
1	C	432	<div> <div>3%</div> <div> <div></div> <div>91%</div> <div>8%</div> <div>•</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NA	A	502	-	-	-	X
3	NA	C	502	-	-	-	X

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 7721 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenosylhomocysteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	429	Total	C	N	O	S	0	0	0
			3321	2112	570	614	25			
1	C	430	Total	C	N	O	S	0	0	0
			3329	2116	571	617	25			

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula:  $C_{21}H_{27}N_7O_{14}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		
3	C	1	Total	Na	0	0
			1	1		

- Molecule 4 is (2S,3R,5R)-3-(6-amino-9H-purin-9-yl)-2-hydroxy-5-(hydroxymethyl)cyclopent anone (three-letter code: ARJ) (formula: C<sub>11</sub>H<sub>13</sub>N<sub>5</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			19	11	5	3		
4	C	1	Total	C	N	O	0	0
			19	11	5	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	489	Total	O	0	0
			489	489		
5	C	454	Total	O	0	0
			454	454		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Adenosylhomocysteinase



#### • Molecule 1: Adenosylhomocysteinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.20Å 102.88Å 174.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.55 49.10 – 1.55	Depositor EDS
% Data completeness (in resolution range)	97.2 (50.00-1.55) 97.2 (49.10-1.55)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.71 (at 1.55Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, $R_{free}$	0.179 , 0.210 0.177 , 0.207	Depositor DCC
$R_{free}$ test set	6227 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.3	Xtriage
Anisotropy	0.470	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 48.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.008 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	7721	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, ARJ, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.76	0/3387	0.76	1/4583 (0.0%)
1	C	0.74	0/3395	0.77	6/4594 (0.1%)
All	All	0.75	0/6782	0.76	7/9177 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	205	ARG	NE-CZ-NH2	-6.20	117.20	120.30
1	C	111	LEU	CB-CG-CD1	6.19	121.53	111.00
1	A	263	ASP	CB-CG-OD1	5.32	123.08	118.30
1	C	190	ASP	CB-CG-OD2	-5.31	113.52	118.30
1	C	190	ASP	CB-CG-OD1	5.28	123.05	118.30
1	C	238	ARG	NE-CZ-NH2	-5.15	117.72	120.30
1	C	111	LEU	CA-CB-CG	5.10	127.03	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3321	0	3344	36	0
1	C	3329	0	3348	31	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	44	0	26	0	0
2	C	44	0	26	0	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
4	A	19	0	13	0	0
4	C	19	0	13	1	0
5	A	489	0	0	12	2
5	C	454	0	0	13	2
All	All	7721	0	6770	67	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:MET:HG2	5:A:971:HOH:O	1.40	1.21
1:A:230:GLN:HG2	5:A:860:HOH:O	1.40	1.17
1:C:162:HIS:HB3	5:C:967:HOH:O	1.48	1.12
1:C:162:HIS:HD2	5:C:934:HOH:O	1.44	1.00
1:C:396:GLU:CD	5:C:603:HOH:O	2.01	0.99
1:C:230:GLN:HG2	5:C:931:HOH:O	1.64	0.95
1:C:188:LYS:NZ	5:C:601:HOH:O	2.04	0.88
1:C:188:LYS:HE3	5:C:610:HOH:O	1.80	0.81
1:A:233:ARG:NH1	5:A:601:HOH:O	1.99	0.81
1:A:188:LYS:NZ	5:A:602:HOH:O	2.19	0.75
1:A:4:LYS:HD2	5:A:618:HOH:O	1.85	0.75
1:A:410:THR:H	1:A:413:GLN:HE21	1.35	0.75
1:A:215:VAL:H	1:A:270:ASN:HD22	1.35	0.75
1:A:162:HIS:HD2	5:A:937:HOH:O	1.69	0.74
1:C:215:VAL:H	1:C:270:ASN:HD22	1.33	0.73
1:A:162:HIS:CD2	5:A:937:HOH:O	2.43	0.71
1:C:107:ASP:O	1:C:111:LEU:HD13	1.92	0.70
1:C:186:LYS:HZ2	1:C:191:ASN:HD21	1.41	0.66
1:C:162:HIS:CD2	5:C:934:HOH:O	2.29	0.65
1:A:34:ARG:HA	1:A:37:GLU:HG2	1.84	0.59
1:C:380:PRO:O	1:C:384:HIS:HE1	1.85	0.58
1:A:191:ASN:N	1:A:191:ASN:HD22	2.01	0.58
1:A:380:PRO:O	1:A:384:HIS:HE1	1.87	0.58
1:A:230:GLN:OE1	1:A:233:ARG:NH2	2.29	0.57
1:C:410:THR:H	1:C:413:GLN:HE21	1.53	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:ILE:HD11	1:A:139:ILE:HD12	1.88	0.56
1:C:191:ASN:N	1:C:191:ASN:HD22	2.03	0.56
1:C:110:TYR:HD2	1:C:111:LEU:HD12	1.71	0.56
1:C:186:LYS:NZ	1:C:191:ASN:HD21	2.04	0.55
1:A:200:ILE:HG12	5:A:655:HOH:O	2.08	0.54
1:C:226:LYS:NZ	1:C:251:GLN:HE22	2.06	0.53
1:A:248:ASN:HA	1:A:251:GLN:HE21	1.73	0.53
1:C:351:MET:O	5:C:602:HOH:O	2.19	0.53
1:A:191:ASN:H	1:A:191:ASN:HD22	1.55	0.53
1:C:226:LYS:HZ3	1:C:251:GLN:HE22	1.58	0.52
1:C:215:VAL:H	1:C:270:ASN:ND2	2.05	0.51
1:A:405:LYS:HE3	5:A:924:HOH:O	2.10	0.51
1:A:186:LYS:HZ2	1:A:191:ASN:HD21	1.58	0.51
1:A:215:VAL:H	1:A:270:ASN:ND2	2.07	0.51
1:C:198:SER:OG	1:C:353:HIS:HD2	1.96	0.49
1:A:410:THR:H	1:A:413:GLN:NE2	2.07	0.49
1:C:34:ARG:NH2	5:C:605:HOH:O	2.40	0.47
1:C:389:LYS:HE3	1:C:422:ASN:O	2.15	0.47
1:A:322:LYS:HB2	1:A:323:PRO:HD2	1.96	0.47
1:C:248:ASN:HA	1:C:251:GLN:HE21	1.80	0.47
1:A:428:ASP:CG	5:A:614:HOH:O	2.53	0.47
1:A:125:LEU:HD11	1:A:139:ILE:CD1	2.46	0.45
1:C:4:LYS:HE2	5:C:830:HOH:O	2.17	0.45
1:A:198:SER:OG	1:A:353:HIS:HD2	1.99	0.45
1:C:191:ASN:HD22	1:C:191:ASN:H	1.65	0.45
1:A:186:LYS:NZ	1:A:191:ASN:HD21	2.14	0.44
1:A:184:VAL:HG21	1:A:432:TYR:CE1	2.53	0.43
1:A:181:ASN:HA	1:A:186:LYS:HD2	2.01	0.43
1:C:403:ASN:HB3	5:C:919:HOH:O	2.19	0.42
1:C:55:HIS:CD2	4:C:503:ARJ:H5A'	2.54	0.42
1:A:181:ASN:ND2	1:A:186:LYS:HZ2	2.18	0.42
1:A:199:LEU:HB3	5:A:655:HOH:O	2.18	0.42
1:C:396:GLU:OE1	5:C:603:HOH:O	2.21	0.42
1:A:121:LYS:HB2	1:A:121:LYS:HE2	1.19	0.42
1:A:185:THR:HA	1:A:189:PHE:CD1	2.55	0.42
1:C:226:LYS:NZ	1:C:251:GLN:NE2	2.68	0.41
1:C:238:ARG:HG3	5:C:966:HOH:O	2.20	0.41
1:C:111:LEU:O	1:C:115:GLU:HG3	2.20	0.41
1:A:317:GLU:HB2	1:A:329:TRP:HB2	2.03	0.41
1:A:158:THR:HG22	1:A:162:HIS:CE1	2.56	0.41
1:A:375:HIS:HE1	5:A:980:HOH:O	2.05	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:387:PRO:HD2	1:A:390:LEU:HD12	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:866:HOH:O	5:C:896:HOH:O[2_445]	2.07	0.13
5:A:819:HOH:O	5:C:668:HOH:O[2_445]	2.11	0.09

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	427/432 (99%)	414 (97%)	12 (3%)	1 (0%)	51	23
1	C	428/432 (99%)	417 (97%)	10 (2%)	1 (0%)	51	23
All	All	855/864 (99%)	831 (97%)	22 (3%)	2 (0%)	51	23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	55	HIS
1	C	55	HIS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	353/356 (99%)	343 (97%)	10 (3%)	49	16
1	C	354/356 (99%)	344 (97%)	10 (3%)	49	16
All	All	707/712 (99%)	687 (97%)	20 (3%)	49	16

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LYS
1	A	27	ASN
1	A	87	HIS
1	A	121	LYS
1	A	191	ASN
1	A	335	ARG
1	A	344	LEU
1	A	411	GLU
1	A	415	GLN
1	A	428	ASP
1	C	87	HIS
1	C	94	LYS
1	C	111	LEU
1	C	190	ASP
1	C	191	ASN
1	C	210	MET
1	C	233	ARG
1	C	335	ARG
1	C	344	LEU
1	C	428	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	ASN
1	A	181	ASN
1	A	191	ASN
1	A	248	ASN
1	A	251	GLN
1	A	270	ASN
1	A	353	HIS
1	A	365	GLN
1	A	384	HIS
1	A	413	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	27	ASN
1	C	162	HIS
1	C	181	ASN
1	C	191	ASN
1	C	248	ASN
1	C	251	GLN
1	C	270	ASN
1	C	324	GLN
1	C	353	HIS
1	C	365	GLN
1	C	384	HIS
1	C	413	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAD	A	501	-	41,48,48	1.44	3 (7%)	43,73,73	2.47	9 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	ARJ	A	503	-	17,21,21	1.48	5 (29%)	13,31,31	3.52	6 (46%)
2	NAD	C	501	-	41,48,48	1.24	2 (4%)	43,73,73	2.58	6 (13%)
4	ARJ	C	503	-	17,21,21	1.75	3 (17%)	13,31,31	2.77	6 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	501	-	-	0/22/62/62	0/5/5/5
4	ARJ	A	503	-	-	0/2/22/22	0/3/3/3
2	NAD	C	501	-	-	0/22/62/62	0/5/5/5
4	ARJ	C	503	-	-	0/2/22/22	0/3/3/3

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	503	ARJ	C5'-C4'	-2.35	1.49	1.54
4	A	503	ARJ	C5-C4	2.15	1.45	1.40
2	C	501	NAD	O4D-C1D	2.23	1.44	1.41
4	C	503	ARJ	C2-N3	2.28	1.36	1.32
4	A	503	ARJ	C4-N3	2.38	1.39	1.35
4	A	503	ARJ	C2-N3	2.50	1.36	1.32
4	A	503	ARJ	O3D-C3D	2.78	1.26	1.21
2	A	501	NAD	C2A-N1A	2.85	1.39	1.33
2	A	501	NAD	C2A-N3A	3.78	1.38	1.32
4	C	503	ARJ	O3D-C3D	4.08	1.28	1.21
4	C	503	ARJ	C4-N3	4.88	1.42	1.35
2	C	501	NAD	O7N-C7N	5.91	1.36	1.24
2	A	501	NAD	O7N-C7N	6.39	1.37	1.24

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	501	NAD	N3A-C2A-N1A	-12.34	118.11	128.86
2	A	501	NAD	N3A-C2A-N1A	-12.18	118.25	128.86
4	A	503	ARJ	N3-C2-N1	-9.75	120.37	128.86
2	C	501	NAD	O7N-C7N-C3N	-6.85	111.61	119.62
4	C	503	ARJ	N3-C2-N1	-5.27	124.27	128.86
2	A	501	NAD	C4B-O4B-C1B	-4.10	105.40	109.77

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	503	ARJ	O6'-C6'-C4'	-4.09	96.65	112.01
4	A	503	ARJ	O6'-C6'-C4'	-3.95	97.18	112.01
2	C	501	NAD	C4B-O4B-C1B	-3.92	105.60	109.77
2	C	501	NAD	C5N-C4N-C3N	-3.31	116.45	120.35
4	A	503	ARJ	C4-C5-N7	-3.24	106.28	109.41
4	C	503	ARJ	C4-C5-N7	-3.21	106.31	109.41
2	A	501	NAD	C5N-C4N-C3N	-3.12	116.68	120.35
2	A	501	NAD	O7N-C7N-C3N	-2.52	116.68	119.62
2	A	501	NAD	O7N-C7N-N7N	-2.27	119.35	122.58
4	C	503	ARJ	C2-N1-C6	2.08	122.40	118.77
2	A	501	NAD	C2N-C3N-C4N	2.30	120.89	118.26
4	A	503	ARJ	N6-C6-N1	2.41	123.54	118.77
2	A	501	NAD	C4D-O4D-C1D	2.46	112.39	109.77
4	C	503	ARJ	N6-C6-N1	2.52	123.77	118.77
4	A	503	ARJ	C2-N1-C6	3.08	124.17	118.77
2	A	501	NAD	C2A-N1A-C6A	3.11	124.20	118.77
2	C	501	NAD	C2N-C3N-C4N	3.18	121.89	118.26
4	A	503	ARJ	C6'-C4'-C3D	4.27	119.12	111.47
2	C	501	NAD	C3N-C7N-N7N	4.90	123.37	117.77
4	C	503	ARJ	C6'-C4'-C3D	5.12	120.63	111.47
2	A	501	NAD	C3N-C7N-N7N	5.42	123.96	117.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	503	ARJ	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	429/432 (99%)	-0.03	10 (2%) 61 67	13, 19, 30, 41	0
1	C	430/432 (99%)	0.04	13 (3%) 51 59	13, 20, 32, 52	0
All	All	859/864 (99%)	0.01	23 (2%) 55 62	13, 19, 31, 52	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	3	ASP	5.5
1	C	111	LEU	4.9
1	C	4	LYS	4.1
1	A	4	LYS	3.9
1	A	41	ALA	3.4
1	C	142	LYS	3.4
1	C	377	ASP	3.4
1	C	172	ILE	3.4
1	A	121	LYS	3.2
1	C	376	PRO	3.1
1	C	5	LEU	3.0
1	A	122	ASP	2.8
1	C	122	ASP	2.8
1	C	124	PRO	2.6
1	C	141	THR	2.6
1	C	123	GLY	2.5
1	A	145	GLN	2.4
1	C	121	LYS	2.4
1	A	38	MET	2.2
1	A	216	ALA	2.1
1	A	415	GLN	2.1
1	A	217	VAL	2.0
1	A	224	VAL	2.0



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	NA	C	502	1/1	0.98	0.11	4.92	24,24,24,24	0
3	NA	A	502	1/1	0.98	0.11	4.33	27,27,27,27	0
4	ARJ	C	503	19/19	0.95	0.07	-0.10	12,14,18,22	0
4	ARJ	A	503	19/19	0.94	0.07	-0.31	12,15,20,22	0
2	NAD	C	501	44/44	0.97	0.06	-0.65	12,15,17,18	0
2	NAD	A	501	44/44	0.97	0.06	-1.05	11,15,17,20	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.