



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 10, 2018 – 03:39 PM EST

PDB ID : 6B04  
Title : Crystal structure of CfFPPS2, a lepidopteran type-II farnesyl diphosphate synthase, complexed with [2-(1-methylpyridin-2-yl)-1-phosphono-ethyl]phosphonic acid (inhibitor 1b)  
Authors : Picard, M.-E.; Cusson, M.; Shi, R.  
Deposited on : 2017-09-13  
Resolution : 1.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030736  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030736


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2964 (1.86-1.82)
Clashscore	112137	3197 (1.86-1.82)
Ramachandran outliers	110173	3164 (1.86-1.82)
Sidechain outliers	110143	3165 (1.86-1.82)
RSRZ outliers	101464	2973 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	341	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>12%</div> <div>.</div> </div> </div>
1	B	341	<div> <div>2%</div> <div> <div></div> <div>92%</div> <div>6%</div> <div>..</div> </div> </div>
1	C	341	<div> <div>%</div> <div> <div></div> <div>89%</div> <div>7%</div> <div>..</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	C	405	-	-	-	X
4	EDO	C	406	-	-	-	X

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 9168 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Farnesyl diphosphate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	341	Total	C	N	O	S	0	2	0
			2755	1768	445	521	21			
1	B	338	Total	C	N	O	S	0	1	0
			2722	1749	439	513	21			
1	C	335	Total	C	N	O	S	0	2	0
			2700	1735	434	510	21			

- Molecule 2 is [2-(1-methylpyridin-2-yl)-1-phosphono-ethyl]phosphonic acid (three-letter code: C6J) (formula: C<sub>8</sub>H<sub>14</sub>NO<sub>6</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			17	8	1	6	2		
2	B	1	Total	C	N	O	P	0	0
			17	8	1	6	2		
2	C	1	Total	C	N	O	P	0	0
			17	8	1	6	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	3	Total	Mg	0	0
			3	3		
3	A	3	Total	Mg	0	0
			3	3		
3	C	3	Total	Mg	0	0
			3	3		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	313	Total	O	0	0
			313	313		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	297	Total 297	O 297	0	0
5	C	301	Total 301	O 301	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Farnesyl diphosphate synthase



- Molecule 1: Farnesyl diphosphate synthase



- Molecule 1: Farnesyl diphosphate synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	187.44Å 122.45Å 68.76Å 90.00° 106.60° 90.00°	Depositor
Resolution (Å)	101.18 – 1.83 44.91 – 1.83	Depositor EDS
% Data completeness (in resolution range)	99.7 (101.18-1.83) 99.7 (44.91-1.83)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.94 (at 1.83Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.157 , 0.191 0.167 , 0.199	Depositor DCC
$R_{free}$ test set	6574 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.5	Xtriage
Anisotropy	0.037	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 52.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	9168	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, C6J, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.16	4/2819 (0.1%)	1.06	8/3813 (0.2%)
1	B	1.09	1/2782 (0.0%)	1.01	5/3762 (0.1%)
1	C	1.17	6/2763 (0.2%)	1.04	10/3739 (0.3%)
All	All	1.14	11/8364 (0.1%)	1.04	23/11314 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	176	SER	CA-CB	6.00	1.61	1.52
1	A	127	TYR	CB-CG	-5.99	1.42	1.51
1	C	205[A]	ASN	CG-ND2	5.43	1.46	1.32
1	C	205[B]	ASN	CG-ND2	5.43	1.46	1.32
1	B	379	GLU	CD-OE1	5.41	1.31	1.25
1	C	205[A]	ASN	CB-CG	5.39	1.63	1.51
1	C	205[B]	ASN	CB-CG	5.39	1.63	1.51
1	C	80	THR	CB-CG2	-5.37	1.34	1.52
1	A	157	ARG	CZ-NH2	5.35	1.40	1.33
1	A	181	SER	CB-OG	-5.23	1.35	1.42
1	C	276	CYS	CB-SG	-5.13	1.73	1.81

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	344	ARG	NE-CZ-NH1	8.61	124.60	120.30
1	C	260	LEU	CB-CG-CD2	8.08	124.74	111.00
1	A	364	ARG	NE-CZ-NH2	-7.57	116.52	120.30
1	A	221	MET	CG-SD-CE	7.35	111.96	100.20
1	B	157	ARG	NE-CZ-NH1	7.10	123.85	120.30
1	C	344	ARG	NE-CZ-NH1	6.59	123.60	120.30
1	B	157	ARG	NE-CZ-NH2	-6.56	117.02	120.30
1	B	364	ARG	NE-CZ-NH1	6.56	123.58	120.30
1	A	344	ARG	NE-CZ-NH2	-6.54	117.03	120.30
1	A	151	ASP	CB-CG-OD2	6.38	124.04	118.30
1	A	179	MET	CG-SD-CE	6.28	110.25	100.20
1	C	295	ASP	CB-CG-OD2	-6.05	112.85	118.30
1	A	64	ASP	CB-CG-OD1	-5.61	113.25	118.30
1	C	295	ASP	CB-CG-OD1	5.57	123.31	118.30
1	C	351	GLN	CA-CB-CG	5.52	125.54	113.40
1	C	104	ARG	NE-CZ-NH1	5.47	123.03	120.30
1	A	357	LEU	CB-CG-CD1	5.39	120.16	111.00
1	C	321	ARG	NE-CZ-NH2	5.39	122.99	120.30
1	C	223	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	C	344	ARG	NE-CZ-NH2	-5.27	117.67	120.30
1	C	148	ASP	CB-CG-OD2	5.14	122.93	118.30
1	B	221	MET	CG-SD-CE	5.13	108.41	100.20
1	B	179	MET	CG-SD-CE	5.04	108.26	100.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	222	GLU	Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2755	0	2753	20	0
1	B	2722	0	2720	13	0
1	C	2700	0	2692	10	0
2	A	17	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	17	0	0	0	0
2	C	17	0	0	0	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
3	C	3	0	0	0	0
4	A	4	0	6	0	0
4	B	8	0	12	0	0
4	C	8	0	12	0	0
5	A	313	0	0	7	0
5	B	297	0	0	2	1
5	C	301	0	0	5	2
All	All	9168	0	8195	41	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:255[B]:CYS:SG	5:B:722:HOH:O	2.12	1.06
1:A:353:GLN:OE1	5:A:501:HOH:O	1.93	0.83
1:A:224:ARG:HD2	1:A:229:TYR:CZ	2.19	0.78
1:C:203:LEU:HD11	1:C:255[A]:CYS:SG	2.30	0.72
1:B:157:ARG:NH2	5:B:502:HOH:O	2.24	0.70
1:C:198:THR:HG21	5:C:749:HOH:O	1.92	0.69
1:B:104:ARG:NH1	1:B:140:GLN:OE1	2.27	0.68
1:C:278:GLU:HG3	1:C:372:GLN:HE22	1.58	0.67
1:A:198:THR:HG21	5:A:753:HOH:O	1.96	0.65
1:A:267:VAL:O	1:A:271:LYS:HG3	2.00	0.62
1:B:58:LYS:HD3	1:B:61:SER:H	1.72	0.55
1:A:104:ARG:NH1	1:A:140:GLN:OE1	2.37	0.55
1:B:221:MET:CE	1:B:221:MET:HA	2.36	0.54
1:C:199:ASN:HD22	1:C:199:ASN:H	1.56	0.54
1:A:224:ARG:HD2	1:A:229:TYR:OH	2.07	0.53
1:A:301:LYS:HE2	5:A:766:HOH:O	2.10	0.52
1:A:225:GLN:HG3	1:A:228:ASP:HB3	1.92	0.52
1:A:346:LYS:NZ	5:A:505:HOH:O	2.44	0.50
1:B:252:LEU:HB3	1:B:253:PRO:HD3	1.93	0.49
1:A:116:GLU:OE1	5:A:503:HOH:O	2.20	0.49
1:C:302:MET:HE3	5:C:526:HOH:O	2.12	0.49
1:A:347[A]:GLU:OE1	5:A:502:HOH:O	2.20	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:98:ALA:O	1:B:157:ARG:NH1	2.40	0.47
1:C:222:GLU:HG3	5:C:757:HOH:O	2.15	0.46
1:A:156:ARG:HG2	1:A:157:ARG:HG3	1.97	0.46
1:A:201:VAL:HG11	1:B:201:VAL:HG11	1.97	0.46
1:B:227:SER:HB2	1:B:331:THR:HB	1.99	0.45
1:A:224:ARG:HD2	1:A:229:TYR:CE1	2.52	0.44
1:B:265:ASP:C	1:B:265:ASP:OD1	2.56	0.44
1:A:309:ALA:HA	1:A:334:TYR:CE2	2.53	0.44
1:A:353:GLN:NE2	5:A:513:HOH:O	2.50	0.43
1:B:309:ALA:HA	1:B:334:TYR:CE2	2.53	0.43
1:A:108:THR:OG1	1:A:253:PRO:HB3	2.18	0.43
1:C:309:ALA:HA	1:C:334:TYR:CE2	2.54	0.43
1:A:210:HIS:NE2	1:B:78:GLU:HG3	2.35	0.41
1:C:191:PHE:O	1:C:197:TYR:HB2	2.20	0.41
1:A:215:GLN:O	1:A:219:VAL:HG22	2.21	0.41
1:A:194:LYS:NZ	5:C:511:HOH:O	2.52	0.41
1:B:77:SER:HB3	1:B:185:TYR:CE1	2.57	0.40
1:C:198:THR:HG23	5:C:689:HOH:O	2.21	0.40
1:C:237:TYR:CZ	1:C:241:VAL:HG11	2.56	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:759:HOH:O	5:C:739:HOH:O[1_554]	1.89	0.31
5:C:757:HOH:O	5:C:791:HOH:O[2_656]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	341/341 (100%)	333 (98%)	8 (2%)	0	<b>100</b> <b>100</b>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	335/341 (98%)	328 (98%)	7 (2%)	0	100	100
1	C	333/341 (98%)	325 (98%)	7 (2%)	1 (0%)	44	28
All	All	1009/1023 (99%)	986 (98%)	22 (2%)	1 (0%)	55	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	222	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	306/304 (101%)	298 (97%)	8 (3%)	51	34
1	B	302/304 (99%)	297 (98%)	5 (2%)	66	52
1	C	300/304 (99%)	291 (97%)	9 (3%)	46	27
All	All	908/912 (100%)	886 (98%)	22 (2%)	54	37

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	GLU
1	A	82	VAL
1	A	195	LYS
1	A	222	GLU
1	A	225	GLN
1	A	260	LEU
1	A	264	SER
1	A	387	ARG
1	B	58	LYS
1	B	70	LEU
1	B	157	ARG
1	B	221	MET
1	B	337	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	70	LEU
1	C	104	ARG
1	C	167	GLU
1	C	195	LYS
1	C	199	ASN
1	C	221	MET
1	C	223	ARG
1	C	227	SER
1	C	260	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	199	ASN
1	A	390	HIS
1	B	76	ASN
1	B	199	ASN
1	B	360	GLN
1	C	188	HIS
1	C	199	ASN
1	C	372	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 9 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	C6J	A	401	3	15,17,17	3.99	12 (80%)	22,26,26	3.52	11 (50%)
4	EDO	A	405	-	3,3,3	0.95	0	2,2,2	0.49	0
2	C6J	B	401	3	15,17,17	3.58	9 (60%)	22,26,26	3.51	9 (40%)
4	EDO	B	405	-	3,3,3	0.98	0	2,2,2	0.47	0
4	EDO	B	406	-	3,3,3	0.76	0	2,2,2	1.01	0
2	C6J	C	401	3	15,17,17	2.92	10 (66%)	22,26,26	3.59	14 (63%)
4	EDO	C	405	-	3,3,3	0.77	0	2,2,2	0.84	0
4	EDO	C	406	-	3,3,3	0.39	0	2,2,2	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	C6J	A	401	3	-	0/16/27/27	0/1/1/1
4	EDO	A	405	-	-	0/1/1/1	0/0/0/0
2	C6J	B	401	3	-	0/16/27/27	0/1/1/1
4	EDO	B	405	-	-	0/1/1/1	0/0/0/0
4	EDO	B	406	-	-	0/1/1/1	0/0/0/0
2	C6J	C	401	3	-	0/16/27/27	0/1/1/1
4	EDO	C	405	-	-	0/1/1/1	0/0/0/0
4	EDO	C	406	-	-	0/1/1/1	0/0/0/0

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	C6J	CAA-NAF	-7.09	1.33	1.47
2	A	401	C6J	CAC-CAD	-6.11	1.37	1.53
2	A	401	C6J	CAA-NAF	-5.54	1.36	1.47
2	B	401	C6J	CAC-CAD	-5.37	1.39	1.53
2	C	401	C6J	CAC-CAD	-5.28	1.39	1.53
2	B	401	C6J	PAI-OAO	-4.67	1.47	1.54
2	A	401	C6J	CAD-CAE	-4.43	1.40	1.52

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	C6J	CAB-CAC	-4.37	1.34	1.51
2	C	401	C6J	CAD-CAE	-4.16	1.41	1.52
2	B	401	C6J	CAD-CAE	-4.07	1.41	1.52
2	B	401	C6J	CAB-CAA	-4.04	1.36	1.51
2	B	401	C6J	CAG-CAE	-3.43	1.45	1.53
2	A	401	C6J	CAB-CAA	-3.20	1.39	1.51
2	A	401	C6J	PAK-OAM	-3.14	1.49	1.54
2	B	401	C6J	CAB-CAC	-2.95	1.39	1.51
2	A	401	C6J	CAG-CAE	-2.91	1.46	1.53
2	C	401	C6J	CAB-CAA	-2.90	1.40	1.51
2	C	401	C6J	CAB-CAC	-2.77	1.40	1.51
2	A	401	C6J	PAI-OAP	-2.72	1.50	1.54
2	C	401	C6J	CAA-NAF	-2.53	1.42	1.47
2	C	401	C6J	PAI-OAO	-2.16	1.51	1.54
2	C	401	C6J	CAG-CAE	-2.15	1.48	1.53
2	B	401	C6J	PAI-OAP	2.23	1.58	1.54
2	C	401	C6J	PAK-OAN	3.01	1.55	1.49
2	A	401	C6J	PAK-OAN	3.05	1.55	1.49
2	C	401	C6J	PAK-OAL	4.07	1.61	1.54
2	C	401	C6J	PAI-CAH	4.56	1.88	1.81
2	A	401	C6J	PAK-CAH	4.70	1.88	1.81
2	B	401	C6J	PAI-CAH	5.05	1.88	1.81
2	A	401	C6J	PAI-OAQ	5.52	1.59	1.49
2	A	401	C6J	PAI-OAO	5.53	1.64	1.54

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	C6J	OAL-PAK-OAN	-3.12	105.48	113.41
2	A	401	C6J	OAO-PAI-OAQ	-2.77	106.37	113.41
2	C	401	C6J	OAN-PAK-CAH	-2.55	104.77	112.13
2	C	401	C6J	OAM-PAK-OAN	-2.43	107.22	113.41
2	C	401	C6J	OAL-PAK-OAM	2.24	114.03	107.69
2	B	401	C6J	CAH-CAG-CAE	2.32	116.61	113.41
2	B	401	C6J	CAG-CAE-CAD	2.34	117.52	113.08
2	C	401	C6J	CAB-CAC-CAD	2.40	116.39	111.42
2	C	401	C6J	OAM-PAK-CAH	2.41	114.41	106.88
2	C	401	C6J	CAD-CAE-NAF	2.63	117.11	111.12
2	B	401	C6J	CAD-CAE-NAF	2.72	117.32	111.12
2	A	401	C6J	CAB-CAC-CAD	2.84	117.30	111.42
2	A	401	C6J	CAD-CAE-NAF	3.01	117.97	111.12
2	C	401	C6J	CAH-CAG-CAE	3.08	117.65	113.41

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	401	C6J	CAG-CAE-CAD	3.12	118.99	113.08
2	B	401	C6J	CAA-NAF-CAE	3.24	119.09	112.75
2	C	401	C6J	CAA-NAF-CAE	3.56	119.72	112.75
2	A	401	C6J	CAJ-NAF-CAA	3.89	117.20	110.58
2	C	401	C6J	CAC-CAB-CAA	3.99	119.17	111.21
2	A	401	C6J	CAB-CAA-NAF	4.06	117.97	111.29
2	B	401	C6J	CAJ-NAF-CAA	4.14	117.61	110.58
2	A	401	C6J	CAA-NAF-CAE	4.42	121.42	112.75
2	C	401	C6J	CAJ-NAF-CAA	4.90	118.91	110.58
2	B	401	C6J	CAC-CAB-CAA	5.20	121.56	111.21
2	A	401	C6J	CAC-CAD-CAE	5.41	122.04	110.85
2	A	401	C6J	CAH-CAG-CAE	5.49	120.99	113.41
2	A	401	C6J	CAC-CAB-CAA	5.99	123.14	111.21
2	B	401	C6J	CAC-CAD-CAE	6.01	123.28	110.85
2	C	401	C6J	CAB-CAA-NAF	6.45	121.88	111.29
2	B	401	C6J	CAB-CAA-NAF	7.08	122.92	111.29
2	C	401	C6J	CAC-CAD-CAE	7.11	125.56	110.85
2	C	401	C6J	CAJ-NAF-CAE	8.60	121.27	112.44
2	A	401	C6J	CAJ-NAF-CAE	8.66	121.33	112.44
2	B	401	C6J	CAJ-NAF-CAE	10.06	122.77	112.44

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	341/341 (100%)	-0.38	8 (2%) 61 58	22, 32, 56, 70	0
1	B	338/341 (99%)	-0.50	7 (2%) 64 62	19, 29, 55, 76	0
1	C	335/341 (98%)	-0.50	4 (1%) 79 79	19, 29, 50, 80	0
All	All	1014/1023 (99%)	-0.46	19 (1%) 67 65	19, 30, 54, 80	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	225	GLN	4.4
1	C	226	LYS	4.1
1	A	224	ARG	3.9
1	A	222	GLU	3.8
1	B	221	MET	3.7
1	B	225	GLN	3.5
1	B	226	LYS	3.4
1	B	60	GLU	3.4
1	A	226	LYS	3.4
1	A	221	MET	3.3
1	C	227	SER	3.3
1	B	227	SER	2.7
1	A	223	ARG	2.7
1	B	224	ARG	2.7
1	A	227	SER	2.6
1	A	266	PRO	2.4
1	B	118	PRO	2.4
1	C	228	ASP	2.3
1	C	340	ALA	2.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	EDO	C	405	4/4	0.92	0.13	3.54	28,31,33,36	0
4	EDO	C	406	4/4	0.94	0.12	2.79	39,41,43,46	0
4	EDO	B	406	4/4	0.90	0.13	1.89	39,43,47,49	0
4	EDO	A	405	4/4	0.86	0.10	0.60	42,43,46,47	0
3	MG	B	404	1/1	0.99	0.08	0.14	21,21,21,21	0
2	C6J	A	401	17/17	0.99	0.08	-0.26	21,24,26,28	0
2	C6J	C	401	17/17	0.99	0.07	-0.35	21,23,26,26	0
3	MG	B	402	1/1	0.99	0.06	-0.46	23,23,23,23	0
3	MG	C	404	1/1	0.99	0.05	-0.52	26,26,26,26	0
2	C6J	B	401	17/17	0.99	0.07	-0.60	20,22,24,24	0
3	MG	A	402	1/1	0.99	0.04	-1.20	23,23,23,23	0
3	MG	A	404	1/1	0.99	0.03	-2.23	25,25,25,25	0
3	MG	C	402	1/1	1.00	0.03	-2.77	23,23,23,23	0
4	EDO	B	405	4/4	0.91	0.19	-	44,48,49,50	0
3	MG	A	403	1/1	0.99	0.06	-	25,25,25,25	0
3	MG	B	403	1/1	0.99	0.04	-	21,21,21,21	0
3	MG	C	403	1/1	0.99	0.04	-	21,21,21,21	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.