



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 21, 2018 – 04:35 PM EST

PDB ID : 6B2R
Title : Crystal structure of Xanthomonas campestris OleA H285A
Authors : Jensen, M.R.; Goblirsch, B.R.; Esler, M.A.; Christenson, J.K.; Mohamed, F.A.; Wackett, L.P.; Wilmot, C.M.
Deposited on : 2017-09-20
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	7172 (1.80-1.76)
Clashscore	112137	8247 (1.80-1.76)
Ramachandran outliers	110173	8154 (1.80-1.76)
Sidechain outliers	110143	8153 (1.80-1.76)
RSRZ outliers	101464	7262 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	358	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="width: 100%; height: 10px; background-color: green;"></div> <div style="width: 100%; height: 10px; background-color: yellow;"></div> <div style="width: 100%; height: 10px; background-color: grey;"></div> <div style="position: absolute; left: 0; top: 0;">%</div> <div style="position: absolute; left: 50%; top: 0;">85%</div> <div style="position: absolute; left: 90%; top: 0;">10%</div> <div style="position: absolute; left: 95%; top: 0;">• •</div> </div>
2	B	358	<div> <div style="width: 14%; height: 10px; background-color: red;"></div> <div style="width: 100%; height: 10px; background-color: green;"></div> <div style="width: 100%; height: 10px; background-color: yellow;"></div> <div style="width: 100%; height: 10px; background-color: grey;"></div> <div style="position: absolute; left: 0; top: 0;">14%</div> <div style="position: absolute; left: 50%; top: 0;">83%</div> <div style="position: absolute; left: 80%; top: 0;">12%</div> <div style="position: absolute; left: 95%; top: 0;">5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	401	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5545 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-oxoacyl-[ACP] synthase III.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	344	Total	C	N	O	S	0	8	0
			2667	1674	470	507	16			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q8PDX2
A	2	GLY	-	expression tag	UNP Q8PDX2
A	3	SER	-	expression tag	UNP Q8PDX2
A	4	SER	-	expression tag	UNP Q8PDX2
A	5	HIS	-	expression tag	UNP Q8PDX2
A	6	HIS	-	expression tag	UNP Q8PDX2
A	7	HIS	-	expression tag	UNP Q8PDX2
A	8	HIS	-	expression tag	UNP Q8PDX2
A	9	HIS	-	expression tag	UNP Q8PDX2
A	10	HIS	-	expression tag	UNP Q8PDX2
A	11	SER	-	expression tag	UNP Q8PDX2
A	12	SER	-	expression tag	UNP Q8PDX2
A	13	GLY	-	expression tag	UNP Q8PDX2
A	14	LEU	-	expression tag	UNP Q8PDX2
A	15	VAL	-	expression tag	UNP Q8PDX2
A	16	PRO	-	expression tag	UNP Q8PDX2
A	17	ARG	-	expression tag	UNP Q8PDX2
A	18	GLY	-	expression tag	UNP Q8PDX2
A	19	SER	-	expression tag	UNP Q8PDX2
A	20	HIS	-	expression tag	UNP Q8PDX2
A	285	ALA	HIS	engineered mutation	UNP Q8PDX2

- Molecule 2 is a protein called 3-oxoacyl-[ACP] synthase III.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	339	Total	C	N	O	S	0	4	0
			2605	1636	456	496	17			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MET	-	initiating methionine	UNP Q8PDX2
B	2	GLY	-	expression tag	UNP Q8PDX2
B	3	SER	-	expression tag	UNP Q8PDX2
B	4	SER	-	expression tag	UNP Q8PDX2
B	5	HIS	-	expression tag	UNP Q8PDX2
B	6	HIS	-	expression tag	UNP Q8PDX2
B	7	HIS	-	expression tag	UNP Q8PDX2
B	8	HIS	-	expression tag	UNP Q8PDX2
B	9	HIS	-	expression tag	UNP Q8PDX2
B	10	HIS	-	expression tag	UNP Q8PDX2
B	11	SER	-	expression tag	UNP Q8PDX2
B	12	SER	-	expression tag	UNP Q8PDX2
B	13	GLY	-	expression tag	UNP Q8PDX2
B	14	LEU	-	expression tag	UNP Q8PDX2
B	15	VAL	-	expression tag	UNP Q8PDX2
B	16	PRO	-	expression tag	UNP Q8PDX2
B	17	ARG	-	expression tag	UNP Q8PDX2
B	18	GLY	-	expression tag	UNP Q8PDX2
B	19	SER	-	expression tag	UNP Q8PDX2
B	20	HIS	-	expression tag	UNP Q8PDX2
B	285	ALA	HIS	engineered mutation	UNP Q8PDX2

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	152	Total	O	0	0
			152	152		
4	B	109	Total	O	0	1
			109	109		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3-oxoacyl-[ACP] synthase III



- Molecule 2: 3-oxoacyl-[ACP] synthase III



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	82.18Å 85.61Å 103.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.77 39.53 – 1.77	Depositor EDS
% Data completeness (in resolution range)	99.5 (50.00-1.77) 99.5 (39.53-1.77)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.64 (at 1.77Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.175 , 0.209 0.186 , 0.219	Depositor DCC
R_{free} test set	3562 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	21.0	Xtriage
Anisotropy	0.138	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.017 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5545	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CSO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.32	10/2711 (0.4%)	1.31	29/3671 (0.8%)
2	B	1.30	8/2629 (0.3%)	1.27	26/3557 (0.7%)
All	All	1.31	18/5340 (0.3%)	1.29	55/7228 (0.8%)

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	101	GLU	CD-OE1	11.82	1.38	1.25
1	A	240	ARG	CD-NE	-7.05	1.34	1.46
2	B	216	GLU	CD-OE1	7.05	1.33	1.25
1	A	193	GLU	CD-OE1	5.68	1.31	1.25
1	A	44	GLU	CD-OE2	5.67	1.31	1.25
1	A	191	GLU	CB-CG	-5.57	1.41	1.52
2	B	177	TYR	CE1-CZ	-5.57	1.31	1.38
1	A	117	GLU	CG-CD	5.54	1.60	1.51
1	A	44	GLU	CG-CD	5.39	1.60	1.51
1	A	113	ARG	CZ-NH2	5.35	1.40	1.33
2	B	310	PHE	CB-CG	-5.30	1.42	1.51
2	B	320	SER	CA-CB	5.26	1.60	1.52
2	B	216	GLU	CD-OE2	5.23	1.31	1.25
2	B	101	GLU	CG-CD	5.22	1.59	1.51
1	A	164	TYR	CE2-CZ	-5.10	1.31	1.38
2	B	188	ASP	CB-CG	5.05	1.62	1.51
1	A	286[A]	GLN	CD-OE1	5.03	1.35	1.24
1	A	286[B]	GLN	CD-OE1	5.03	1.35	1.24

All (55) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	223	ARG	NE-CZ-NH2	-11.95	114.33	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	223	ARG	NE-CZ-NH1	11.90	126.25	120.30
2	B	333	ARG	NE-CZ-NH2	-9.85	115.38	120.30
1	A	240	ARG	NE-CZ-NH1	9.40	125.00	120.30
2	B	251	ARG	NE-CZ-NH1	8.81	124.70	120.30
1	A	155	ARG	NE-CZ-NH1	8.78	124.69	120.30
2	B	113	ARG	NE-CZ-NH1	8.15	124.38	120.30
2	B	214	ARG	NE-CZ-NH1	7.75	124.18	120.30
2	B	230	ARG	NE-CZ-NH1	7.55	124.08	120.30
2	B	113	ARG	NE-CZ-NH2	-7.31	116.64	120.30
1	A	302	ASP	CB-CG-OD2	-7.29	111.74	118.30
2	B	243	LEU	CA-CB-CG	7.13	131.70	115.30
2	B	253	LEU	CA-CB-CG	6.93	131.23	115.30
1	A	230	ARG	NE-CZ-NH1	6.89	123.75	120.30
1	A	240	ARG	NE-CZ-NH2	-6.85	116.88	120.30
1	A	251	ARG	NE-CZ-NH2	-6.56	117.02	120.30
2	B	35	ASP	CB-CG-OD1	6.52	124.17	118.30
1	A	302	ASP	CB-CG-OD1	6.49	124.14	118.30
1	A	113	ARG	NE-CZ-NH1	-6.38	117.11	120.30
2	B	286	GLN	CA-CB-CG	-6.36	99.40	113.40
1	A	155	ARG	NE-CZ-NH2	-6.29	117.15	120.30
1	A	137	PHE	CB-CG-CD1	6.29	125.20	120.80
1	A	195	ARG	NE-CZ-NH2	-6.29	117.16	120.30
1	A	21	MET	CA-CB-CG	-6.11	102.92	113.30
1	A	35	ASP	CB-CG-OD1	6.08	123.78	118.30
2	B	280	ASP	CB-CG-OD2	6.07	123.76	118.30
1	A	113	ARG	NE-CZ-NH2	6.00	123.30	120.30
1	A	249	ASP	CB-CG-OD2	-5.95	112.95	118.30
2	B	101	GLU	CG-CD-OE1	5.94	130.19	118.30
1	A	54	ASP	CB-CG-OD1	5.90	123.61	118.30
1	A	234	GLU	OE1-CD-OE2	-5.89	116.23	123.30
2	B	249	ASP	CB-CG-OD2	-5.84	113.04	118.30
2	B	35	ASP	CB-CG-OD2	-5.83	113.05	118.30
2	B	214	ARG	NE-CZ-NH2	-5.79	117.41	120.30
2	B	90	ARG	NE-CZ-NH1	5.78	123.19	120.30
1	A	76	ASP	CB-CG-OD1	5.67	123.40	118.30
2	B	251	ARG	NE-CZ-NH2	-5.63	117.49	120.30
1	A	72	ARG	NE-CZ-NH1	5.59	123.09	120.30
1	A	48	ARG	NE-CZ-NH2	5.49	123.04	120.30
2	B	95	ASP	CB-CG-OD1	-5.46	113.38	118.30
1	A	333	ARG	NE-CZ-NH1	-5.44	117.58	120.30
1	A	289	ARG	NE-CZ-NH1	5.41	123.00	120.30
2	B	159	ARG	NE-CZ-NH1	5.41	123.00	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	101	GLU	CG-CD-OE2	-5.38	107.54	118.30
1	A	76	ASP	CB-CG-OD2	-5.33	113.51	118.30
1	A	163	ASP	CB-CG-OD1	5.33	123.09	118.30
2	B	333	ARG	CG-CD-NE	-5.30	100.66	111.80
2	B	339	ARG	NE-CZ-NH1	-5.28	117.66	120.30
1	A	240	ARG	CB-CG-CD	5.20	125.12	111.60
2	B	113	ARG	CD-NE-CZ	5.19	130.87	123.60
2	B	339	ARG	NE-CZ-NH2	5.15	122.88	120.30
2	B	234	GLU	OE1-CD-OE2	-5.14	117.13	123.30
2	B	320	SER	CA-CB-OG	-5.14	97.33	111.20
1	A	280	ASP	CB-CG-OD2	5.09	122.88	118.30
1	A	243	LEU	CB-CG-CD2	5.01	119.52	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2667	0	2695	12	0
2	B	2605	0	2637	6	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	A	152	0	0	1	0
4	B	109	0	0	1	0
All	All	5545	0	5348	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:265[A]:PHE:CD2	1:A:299:PHE:CE1	2.68	0.81
1:A:24:GLN:NE2	4:A:501:HOH:O	2.33	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:265[A]:PHE:HD2	1:A:299:PHE:CE1	2.16	0.60
1:A:265[A]:PHE:HD2	1:A:299:PHE:CZ	2.20	0.58
1:A:265[A]:PHE:CD2	1:A:299:PHE:CZ	2.98	0.50
1:A:254:LEU:HB2	1:A:291[B]:HIS:CE1	2.48	0.49
2:B:313:HIS:HB3	2:B:316:ILE:HD11	1.94	0.49
2:B:237:LYS:HE2	4:B:503:HOH:O	2.12	0.49
2:B:269:LYS:HG2	2:B:274:TRP:O	2.13	0.48
1:A:254:LEU:HD23	1:A:254:LEU:C	2.34	0.48
1:A:288:SER:HA	1:A:310:PHE:CZ	2.49	0.47
1:A:143[B]:CYS:SG	1:A:317:GLY:HA2	2.54	0.47
1:A:333:ARG:O	1:A:333:ARG:HG3	2.14	0.47
2:B:77:GLN:O	2:B:78:ASP:HB2	2.15	0.47
1:A:336:LYS:HE3	1:A:358:TRP:O	2.17	0.43
1:A:265[A]:PHE:CD2	1:A:299:PHE:CD1	3.08	0.42
2:B:50:GLN:HB3	2:B:51:PRO:HD3	2.02	0.41
2:B:100:ILE:HD11	2:B:130:VAL:HG12	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/358 (98%)	342 (98%)	8 (2%)	0	100	100
2	B	339/358 (95%)	327 (96%)	11 (3%)	1 (0%)	44	27
All	All	689/716 (96%)	669 (97%)	19 (3%)	1 (0%)	55	37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	55	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/289 (99%)	279 (98%)	6 (2%)	59	43
2	B	275/288 (96%)	266 (97%)	9 (3%)	43	24
All	All	560/577 (97%)	545 (97%)	15 (3%)	51	32

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	72	ARG
1	A	223	ARG
1	A	243	LEU
1	A	265[A]	PHE
1	A	265[B]	PHE
1	A	298	SER
2	B	59	LYS
2	B	60	THR
2	B	72	ARG
2	B	191	GLU
2	B	192	GLU
2	B	196	ASN
2	B	248	THR
2	B	303	PRO
2	B	335	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CSO	B	143[A]	2	4,6,7	2.04	1 (25%)	1,6,8	1.62	0
2	CSO	B	143[B]	2	5,5,7	1.97	1 (20%)	2,5,8	1.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CSO	B	143[A]	2	-	0/1/5/7	0/0/0/0
2	CSO	B	143[B]	2	-	0/1/4/7	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	143[B]	CSO	CB-SG	-3.70	1.73	1.81
2	B	143[A]	CSO	CA-C	3.47	1.54	1.50

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	401	-	5,5,5	0.39	0	5,5,5	0.58	0
3	GOL	B	401	-	5,5,5	0.58	0	5,5,5	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	401	-	-	0/4/4/4	0/0/0/0
3	GOL	B	401	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	344/358 (96%)	-0.01	4 (1%) 79 79	13, 21, 36, 55	0
2	B	338/358 (94%)	0.61	49 (14%) 3 3	13, 23, 58, 86	0
All	All	682/716 (95%)	0.30	53 (7%) 14 14	13, 22, 52, 86	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	243	LEU	8.2
2	B	58	ILE	8.1
2	B	244	ASP	5.7
2	B	194	PHE	5.7
2	B	53	TYR	5.5
2	B	59	LYS	5.2
2	B	193	GLU	5.0
2	B	187	PRO	5.0
2	B	191	GLU	4.8
2	B	56	LEU	4.7
2	B	60	THR	4.7
2	B	189	VAL	4.6
2	B	20	HIS	4.1
2	B	77	GLN	3.6
2	B	54	ASP	3.6
2	B	238	LEU	3.5
2	B	195	ARG	3.4
1	A	299	PHE	3.3
2	B	78	ASP	3.3
2	B	185	THR	3.2
2	B	48	ARG	3.1
2	B	249	ASP	3.1
2	B	245	ARG	3.1
2	B	192	GLU	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	152	ILE	3.0
1	A	243	LEU	3.0
2	B	248	THR	3.0
2	B	186	SER	3.0
2	B	247	VAL	3.0
2	B	239	CYS	2.9
2	B	184	MET	2.7
2	B	57	GLY	2.7
1	A	300	GLY	2.7
2	B	242	ASN	2.6
2	B	55	ARG	2.6
2	B	197	GLU	2.6
2	B	278	GLU	2.6
2	B	175	LEU	2.6
2	B	182	GLU	2.6
2	B	196	ASN	2.6
1	A	15	VAL	2.5
2	B	270	GLN	2.4
2	B	51	PRO	2.3
2	B	188	ASP	2.2
2	B	276	VAL	2.2
2	B	274	TRP	2.2
2	B	240	ARG	2.1
2	B	241	GLY	2.1
2	B	190	THR	2.1
2	B	246	MET	2.1
2	B	251	ARG	2.0
2	B	176	VAL	2.0
2	B	237	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	CSO	B	143[A]	7/8	0.94	0.15	-	16,17,20,20	7
2	CSO	B	143[B]	6/8	0.94	0.15	-	16,16,17,18	6

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	A	401	6/6	0.88	0.19	5.54	33,40,42,53	0
3	GOL	B	401	6/6	0.97	0.14	1.82	25,26,28,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.