



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 4, 2017 – 06:56 PM EDT

PDB ID : 2B3F  
Title : Thermus thermophilus Glucose/Galactose Binding Protein Bound With Galactose  
Authors : Cuneo, M.J.; Changela, A.; Warren, J.J.; Beese, L.S.; Hellinga, H.W.  
Deposited on : unknown  
Resolution : 1.56 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1088 (1.56-1.56)
Clashscore	112137	1132 (1.56-1.56)
Ramachandran outliers	110173	1110 (1.56-1.56)
Sidechain outliers	110143	1108 (1.56-1.56)
RSRZ outliers	101464	1089 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	400	<div> <div>2%</div> <div> <div></div> <div>95%</div> <div></div> </div> <div>..</div> </div>
1	B	400	<div> <div>3%</div> <div> <div></div> <div>96%</div> <div></div> </div> <div>..</div> </div>
1	C	400	<div> <div>4%</div> <div> <div></div> <div>95%</div> <div></div> </div> <div>..</div> </div>
1	D	400	<div> <div>0%</div> <div> <div></div> <div>95%</div> <div></div> </div> <div>..</div> </div>
1	E	400	<div> <div>2%</div> <div> <div></div> <div>97%</div> <div></div> </div> <div>..</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	400	<div><div></div><div>2%</div><div>96%</div><div></div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 20905 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called glucose-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	392	Total	C	N	O	S	0	0	0
			3036	1942	535	545	14			
1	B	392	Total	C	N	O	S	0	0	0
			3036	1942	535	545	14			
1	C	391	Total	C	N	O	S	0	0	0
			3028	1936	534	544	14			
1	D	392	Total	C	N	O	S	0	0	0
			3036	1942	535	545	14			
1	E	392	Total	C	N	O	S	0	0	0
			3036	1942	535	545	14			
1	F	392	Total	C	N	O	S	0	0	0
			3036	1942	535	545	14			

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	INITIATING METHIONINE	UNP Q72KX2
A	133	GLY	GLU	SEE REMARK 999	UNP Q72KX2
A	144	LYS	GLU	SEE REMARK 999	UNP Q72KX2
A	240	ILE	VAL	SEE REMARK 999	UNP Q72KX2
A	308	SER	PHE	SEE REMARK 999	UNP Q72KX2
A	395	HIS	-	EXPRESSION TAG	UNP Q72KX2
A	396	HIS	-	EXPRESSION TAG	UNP Q72KX2
A	397	HIS	-	EXPRESSION TAG	UNP Q72KX2
A	398	HIS	-	EXPRESSION TAG	UNP Q72KX2
A	399	HIS	-	EXPRESSION TAG	UNP Q72KX2
A	400	HIS	-	EXPRESSION TAG	UNP Q72KX2
B	1	MET	-	INITIATING METHIONINE	UNP Q72KX2
B	133	GLY	GLU	SEE REMARK 999	UNP Q72KX2
B	144	LYS	GLU	SEE REMARK 999	UNP Q72KX2
B	240	ILE	VAL	SEE REMARK 999	UNP Q72KX2
B	308	SER	PHE	SEE REMARK 999	UNP Q72KX2
B	395	HIS	-	EXPRESSION TAG	UNP Q72KX2

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	396	HIS	-	EXPRESSION TAG	UNP Q72KX2
B	397	HIS	-	EXPRESSION TAG	UNP Q72KX2
B	398	HIS	-	EXPRESSION TAG	UNP Q72KX2
B	399	HIS	-	EXPRESSION TAG	UNP Q72KX2
B	400	HIS	-	EXPRESSION TAG	UNP Q72KX2
C	1	MET	-	INITIATING METHIONINE	UNP Q72KX2
C	133	GLY	GLU	SEE REMARK 999	UNP Q72KX2
C	144	LYS	GLU	SEE REMARK 999	UNP Q72KX2
C	240	ILE	VAL	SEE REMARK 999	UNP Q72KX2
C	308	SER	PHE	SEE REMARK 999	UNP Q72KX2
C	395	HIS	-	EXPRESSION TAG	UNP Q72KX2
C	396	HIS	-	EXPRESSION TAG	UNP Q72KX2
C	397	HIS	-	EXPRESSION TAG	UNP Q72KX2
C	398	HIS	-	EXPRESSION TAG	UNP Q72KX2
C	399	HIS	-	EXPRESSION TAG	UNP Q72KX2
C	400	HIS	-	EXPRESSION TAG	UNP Q72KX2
D	1	MET	-	INITIATING METHIONINE	UNP Q72KX2
D	133	GLY	GLU	SEE REMARK 999	UNP Q72KX2
D	144	LYS	GLU	SEE REMARK 999	UNP Q72KX2
D	240	ILE	VAL	SEE REMARK 999	UNP Q72KX2
D	308	SER	PHE	SEE REMARK 999	UNP Q72KX2
D	395	HIS	-	EXPRESSION TAG	UNP Q72KX2
D	396	HIS	-	EXPRESSION TAG	UNP Q72KX2
D	397	HIS	-	EXPRESSION TAG	UNP Q72KX2
D	398	HIS	-	EXPRESSION TAG	UNP Q72KX2
D	399	HIS	-	EXPRESSION TAG	UNP Q72KX2
D	400	HIS	-	EXPRESSION TAG	UNP Q72KX2
E	1	MET	-	INITIATING METHIONINE	UNP Q72KX2
E	133	GLY	GLU	SEE REMARK 999	UNP Q72KX2
E	144	LYS	GLU	SEE REMARK 999	UNP Q72KX2
E	240	ILE	VAL	SEE REMARK 999	UNP Q72KX2
E	308	SER	PHE	SEE REMARK 999	UNP Q72KX2
E	395	HIS	-	EXPRESSION TAG	UNP Q72KX2
E	396	HIS	-	EXPRESSION TAG	UNP Q72KX2
E	397	HIS	-	EXPRESSION TAG	UNP Q72KX2
E	398	HIS	-	EXPRESSION TAG	UNP Q72KX2
E	399	HIS	-	EXPRESSION TAG	UNP Q72KX2
E	400	HIS	-	EXPRESSION TAG	UNP Q72KX2
F	1	MET	-	INITIATING METHIONINE	UNP Q72KX2
F	133	GLY	GLU	SEE REMARK 999	UNP Q72KX2
F	144	LYS	GLU	SEE REMARK 999	UNP Q72KX2
F	240	ILE	VAL	SEE REMARK 999	UNP Q72KX2

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	308	SER	PHE	SEE REMARK 999	UNP Q72KX2
F	395	HIS	-	EXPRESSION TAG	UNP Q72KX2
F	396	HIS	-	EXPRESSION TAG	UNP Q72KX2
F	397	HIS	-	EXPRESSION TAG	UNP Q72KX2
F	398	HIS	-	EXPRESSION TAG	UNP Q72KX2
F	399	HIS	-	EXPRESSION TAG	UNP Q72KX2
F	400	HIS	-	EXPRESSION TAG	UNP Q72KX2

- Molecule 2 is BETA-D-GALACTOSE (three-letter code: GAL) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		
2	C	1	Total	C	O	0	0
			12	6	6		
2	D	1	Total	C	O	0	0
			12	6	6		
2	E	1	Total	C	O	0	0
			12	6	6		
2	F	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	436	Total 436	O 436	0	0
3	B	451	Total 451	O 451	0	0
3	C	408	Total 408	O 408	0	0
3	D	472	Total 472	O 472	0	0
3	E	451	Total 451	O 451	0	0
3	F	407	Total 407	O 407	0	0



- Molecule 1: glucose-binding protein



- 


- 

- [illegible]

-



● Molecule 1: glucose-binding protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	121.38 Å   134.96 Å   159.72 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	103.14 – 1.56 42.18 – 1.56	Depositor EDS
% Data completeness (in resolution range)	94.7 (103.14-1.56) 94.7 (42.18-1.56)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.02 (at 1.56 Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.174   ,   0.205 0.188   ,   0.215	Depositor DCC
$R_{free}$ test set	17741 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.6	Xtriage
Anisotropy	0.276	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 55.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	20905	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 70.26 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.2184e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.54	0/3116	0.65	0/4235
1	B	0.52	0/3116	0.64	1/4235 (0.0%)
1	C	0.52	0/3108	0.65	1/4224 (0.0%)
1	D	0.54	0/3116	0.65	1/4235 (0.0%)
1	E	0.52	0/3116	0.66	2/4235 (0.0%)
1	F	0.53	0/3116	0.66	1/4235 (0.0%)
All	All	0.53	0/18688	0.65	6/25399 (0.0%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	80	ARG	NE-CZ-NH1	7.53	124.06	120.30
1	B	297	ARG	NE-CZ-NH1	6.27	123.44	120.30
1	C	83	ASP	CB-CG-OD1	5.48	123.23	118.30
1	F	47	ARG	NE-CZ-NH2	-5.43	117.59	120.30
1	D	83	ASP	CB-CG-OD1	5.30	123.07	118.30
1	E	80	ARG	NE-CZ-NH2	-5.01	117.80	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3036	0	3012	7	0
1	B	3036	0	3012	4	0
1	C	3028	0	3001	6	0
1	D	3036	0	3012	6	0
1	E	3036	0	3012	1	0
1	F	3036	0	3012	5	0
2	A	12	0	12	0	0
2	B	12	0	12	0	0
2	C	12	0	12	0	0
2	D	12	0	12	0	0
2	E	12	0	12	0	0
2	F	12	0	12	0	0
3	A	436	0	0	0	0
3	B	451	0	0	0	0
3	C	408	0	0	0	0
3	D	472	0	0	1	0
3	E	451	0	0	0	0
3	F	407	0	0	0	0
All	All	20905	0	18133	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:169:GLN:OE1	1:F:368:LEU:HD11	1.48	1.11
1:C:169:GLN:OE1	1:C:368:LEU:HD11	1.86	0.75
1:D:47:ARG:HD3	1:D:354:GLU:OE2	1.95	0.67
1:A:364:MET:O	1:A:368:LEU:HD22	1.97	0.64
1:F:270:GLN:H	1:F:270:GLN:CD	2.02	0.63
1:A:23:LEU:HD23	1:A:298:LEU:HD11	1.84	0.59
1:D:117:ASN:HB3	1:D:309:ASN:HD21	1.69	0.57
1:A:20:LEU:HG	1:A:308:SER:HB3	1.87	0.57
1:B:169:GLN:OE1	1:B:368:LEU:HD11	2.04	0.57
1:A:169:GLN:OE1	1:A:368:LEU:HD21	2.06	0.56
1:F:20:LEU:HG	1:F:308:SER:HB3	1.88	0.55
1:B:126:TYR:CD1	1:B:131:LEU:HD22	2.45	0.52
1:A:34:ILE:HG12	1:C:154:GLN:HA	1.91	0.52
1:D:20:LEU:HG	1:D:308:SER:HB3	1.90	0.52
1:A:212:LEU:HD21	1:A:368:LEU:CD1	2.43	0.49
1:A:363:VAL:O	1:A:366:ILE:HG13	2.17	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:169:GLN:OE1	1:F:368:LEU:CD1	2.40	0.45
1:E:54:MET:O	1:E:80:ARG:NH2	2.51	0.44
1:C:364:MET:O	1:C:368:LEU:HG	2.18	0.43
1:D:94:LEU:HA	1:D:94:LEU:HD12	1.92	0.42
1:B:107:TYR:CE2	1:B:108:LYS:HG3	2.54	0.42
1:C:126:TYR:CE1	1:C:262:ALA:HB3	2.54	0.41
1:F:119:HIS:CE1	1:F:314:SER:HB3	2.56	0.41
1:C:34:ILE:N	1:C:34:ILE:HD12	2.36	0.41
1:D:14:GLY:N	1:D:15:PRO:CD	2.84	0.41
1:D:270:GLN:NE2	3:D:3628:HOH:O	2.54	0.40
1:B:119:HIS:CE1	1:B:314:SER:HB3	2.57	0.40
1:C:119:HIS:CE1	1:C:314:SER:HB3	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	390/400 (98%)	385 (99%)	5 (1%)	0	100	100
1	B	390/400 (98%)	385 (99%)	5 (1%)	0	100	100
1	C	389/400 (97%)	382 (98%)	7 (2%)	0	100	100
1	D	390/400 (98%)	383 (98%)	7 (2%)	0	100	100
1	E	390/400 (98%)	386 (99%)	4 (1%)	0	100	100
1	F	390/400 (98%)	385 (99%)	5 (1%)	0	100	100
All	All	2339/2400 (98%)	2306 (99%)	33 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	309/316 (98%)	306 (99%)	3 (1%)	80	61
1	B	309/316 (98%)	308 (100%)	1 (0%)	94	87
1	C	308/316 (98%)	305 (99%)	3 (1%)	80	61
1	D	309/316 (98%)	308 (100%)	1 (0%)	94	87
1	E	309/316 (98%)	306 (99%)	3 (1%)	80	61
1	F	309/316 (98%)	307 (99%)	2 (1%)	89	77
All	All	1853/1896 (98%)	1840 (99%)	13 (1%)	87	74

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	273	PHE
1	A	276	LEU
1	A	368	LEU
1	B	276	LEU
1	C	270	GLN
1	C	276	LEU
1	C	341	ARG
1	D	276	LEU
1	E	213	ASP
1	E	276	LEU
1	E	341	ARG
1	F	213	ASP
1	F	276	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	309	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GAL	A	3310	-	12,12,12	0.67	0	17,17,17	0.84	0
2	GAL	B	3311	-	12,12,12	0.67	0	17,17,17	0.72	0
2	GAL	C	3312	-	12,12,12	0.82	0	17,17,17	0.82	1 (5%)
2	GAL	D	3313	-	12,12,12	0.74	0	17,17,17	0.89	0
2	GAL	E	3314	-	12,12,12	0.60	0	17,17,17	0.91	0
2	GAL	F	3315	-	12,12,12	0.89	0	17,17,17	0.88	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	A	3310	-	-	0/2/22/22	0/1/1/1
2	GAL	B	3311	-	-	0/2/22/22	0/1/1/1
2	GAL	C	3312	-	-	0/2/22/22	0/1/1/1
2	GAL	D	3313	-	-	0/2/22/22	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	E	3314	-	-	0/2/22/22	0/1/1/1
2	GAL	F	3315	-	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3312	GAL	O4-C4-C3	-2.32	105.31	110.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	392/400 (98%)	0.02	8 (2%) 65 71	12, 20, 33, 39	0
1	B	392/400 (98%)	0.08	12 (3%) 49 58	13, 21, 33, 53	0
1	C	391/400 (97%)	0.20	17 (4%) 36 41	14, 22, 35, 54	0
1	D	392/400 (98%)	-0.09	4 (1%) 82 85	12, 20, 28, 36	0
1	E	392/400 (98%)	0.02	6 (1%) 74 79	14, 22, 30, 38	0
1	F	392/400 (98%)	0.07	8 (2%) 65 71	13, 21, 33, 38	0
All	All	2351/2400 (97%)	0.05	55 (2%) 61 67	12, 21, 32, 54	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	392	LEU	11.4
1	B	391	ARG	6.5
1	C	391	ARG	6.3
1	C	258	GLY	4.1
1	C	390	GLY	3.8
1	C	256	LYS	3.5
1	C	259	THR	3.5
1	C	129	ALA	3.2
1	D	392	LEU	3.2
1	A	154	GLN	3.2
1	B	259	THR	3.1
1	C	132	LYS	3.1
1	A	133	GLY	3.0
1	B	132	LYS	3.0
1	B	38	VAL	3.0
1	E	392	LEU	2.8
1	B	390	GLY	2.8
1	F	26	GLN	2.7
1	C	108	LYS	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	256	LYS	2.7
1	D	30	GLY	2.7
1	B	99	LYS	2.7
1	E	94	LEU	2.7
1	C	167	TRP	2.6
1	F	134	TRP	2.6
1	A	140	ARG	2.6
1	C	99	LYS	2.6
1	F	33	VAL	2.6
1	F	154	GLN	2.5
1	C	29	PRO	2.5
1	F	32	GLU	2.5
1	B	167	TRP	2.4
1	E	44	VAL	2.4
1	A	129	ALA	2.4
1	B	39	THR	2.4
1	E	132	LYS	2.4
1	C	38	VAL	2.3
1	C	77	VAL	2.3
1	A	369	GLN	2.3
1	A	26	GLN	2.2
1	F	1	MET	2.2
1	F	133	GLY	2.2
1	A	134	TRP	2.2
1	E	134	TRP	2.2
1	C	107	TYR	2.1
1	C	79	ASN	2.1
1	E	140	ARG	2.1
1	B	131	LEU	2.1
1	B	108	LYS	2.1
1	A	392	LEU	2.1
1	C	140	ARG	2.0
1	C	133	GLY	2.0
1	D	140	ARG	2.0
1	D	338	ARG	2.0
1	F	131	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GAL	A	3310	12/12	0.95	0.10	0.36	11,12,13,16	0
2	GAL	F	3315	12/12	0.93	0.10	-0.03	12,13,15,17	0
2	GAL	E	3314	12/12	0.96	0.08	-0.15	13,14,16,17	0
2	GAL	C	3312	12/12	0.96	0.09	-0.16	11,14,15,18	0
2	GAL	D	3313	12/12	0.97	0.08	-0.30	10,12,13,15	0
2	GAL	B	3311	12/12	0.97	0.09	-0.61	11,13,14,19	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.