



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 10, 2018 – 07:58 AM EST

PDB ID : 6B3P
Title : Crystal structure of CBMbc (family CBM26) from Eubacterium rectale Amy13K in Complex with Maltoheptaose
Authors : Cockburn, D.W.; Wawrzak, Z.; Perez Medina, K.; Koropatkin, N.M.
Deposited on : 2017-09-22
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	<div> <div>9%</div> <div>96%</div> <div>.</div> </div>
1	B	209	<div> <div>6%</div> <div>98%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	409	-	-	-	X
3	EDO	A	412	-	-	-	X
3	EDO	A	413	-	-	-	X
3	EDO	A	414	-	-	-	X
3	EDO	A	415	-	-	-	X
3	EDO	B	409	-	-	-	X
3	EDO	B	410	-	-	-	X
3	EDO	B	411	-	-	-	X
3	EDO	B	413	-	-	-	X
3	EDO	B	414	-	-	-	X
4	FMT	A	417	-	-	-	X
4	FMT	A	420	-	-	-	X
4	FMT	B	416	-	-	-	X
4	FMT	B	423	-	-	-	X
4	FMT	B	424	-	-	-	X
4	FMT	B	425	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7344 atoms, of which 3265 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Amy13K.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	209	Total	C	H	N	O	S	0	2	0
			3129	1024	1498	260	341	6			
1	B	209	Total	C	H	N	O	S	0	0	0
			3106	1016	1489	259	336	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	179	MET	-	initiating methionine	UNP D4JJZ5
B	179	MET	-	initiating methionine	UNP D4JJZ5

- Molecule 2 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			23	6	11	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			22	6	10	6		
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			22	6	11	5		
2	B	1	Total	C	H	O	0	0
			22	6	10	6		
2	B	1	Total	C	H	O	0	0
			21	6	10	5		
2	B	1	Total	C	H	O	0	0
			22	6	11	5		
2	B	1	Total	C	H	O	0	0
			23	6	11	6		
2	B	1	Total	C	H	O	0	0
			21	6	10	5		
2	B	1	Total	C	H	O	0	0
			21	6	10	5		
2	B	1	Total	C	H	O	0	0
			21	6	10	5		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	B	1	Total	C	H	O	0	0
			10	2	6	2		
3	B	1	Total	C	H	O	0	0
			10	2	6	2		
3	B	1	Total	C	H	O	0	0
			10	2	6	2		
3	B	1	Total	C	H	O	0	0
			10	2	6	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	H	O	0	0
			10	2	6	2		
3	B	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 4 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			5	1	2	2		
4	A	1	Total	C	H	O	0	0
			5	1	2	2		
4	A	1	Total	C	H	O	0	0
			5	1	2	2		
4	A	1	Total	C	H	O	0	0
			5	1	2	2		
4	A	1	Total	C	H	O	0	0
			5	1	2	2		
4	A	1	Total	C	H	O	0	0
			5	1	2	2		
4	A	1	Total	C	H	O	0	0
			5	1	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	H	O	0	0
			5	1	2	2		
4	B	1	Total	C	H	O	0	0
			5	1	2	2		
4	B	1	Total	C	H	O	0	0
			5	1	2	2		
4	B	1	Total	C	H	O	0	0
			5	1	2	2		
4	B	1	Total	C	H	O	0	0
			5	1	2	2		
4	B	1	Total	C	H	O	0	0
			5	1	2	2		
4	B	1	Total	C	H	O	0	0
			5	1	2	2		
4	B	1	Total	C	H	O	0	0
			5	1	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	268	Total	O	0	0
			268	268		
5	B	284	Total	O	0	0
			284	284		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Amy13K



- Molecule 1: Amy13K



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	134.20Å 134.20Å 231.13Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.15 – 2.01 28.18 – 2.01	Depositor EDS
% Data completeness (in resolution range)	99.7 (28.15-2.01) 99.7 (28.18-2.01)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 2.01Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.184 , 0.196 0.184 , 0.196	Depositor DCC
R_{free} test set	1995 reflections (2.44%)	DCC
Wilson B-factor (Å ²)	31.8	Xtriage
Anisotropy	0.204	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 55.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7344	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FMT, GLC, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/1677	0.44	0/2291
1	B	0.28	0/1660	0.43	0/2268
All	All	0.26	0/3337	0.43	0/4559

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1631	1498	1501	5	1
1	B	1617	1489	1492	1	0
2	A	79	72	69	0	1
2	B	79	72	69	0	2
3	A	32	48	48	0	0
3	B	32	48	48	0	0
4	A	27	18	9	0	0
4	B	30	20	10	1	0
5	A	268	0	0	0	0
5	B	284	0	0	1	0
All	All	4079	3265	3246	7	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:423:FMT:O2	5:B:501:HOH:O	2.14	0.64
1:A:298:MET:CE	1:A:310:VAL:HG12	2.42	0.49
1:A:181:LEU:HD22	1:A:229:LEU:HD21	1.98	0.45
1:A:200:GLU:O	1:A:201:THR:OG1	2.27	0.45
1:B:387:GLU:H	1:B:387:GLU:CD	2.19	0.44
1:A:250:ASP:OD1	1:A:250:ASP:N	2.52	0.43
1:A:179:MET:N	1:A:229:LEU:O	2.55	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:405:GLC:O1	2:B:407:GLC:C4[10_555]	1.56	0.64
1:A:202:ASN:HD21	2:B:403:GLC:HO3[12_564]	1.29	0.31

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	209/209 (100%)	203 (97%)	6 (3%)	0	100	100
1	B	207/209 (99%)	200 (97%)	7 (3%)	0	100	100
All	All	416/418 (100%)	403 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/172 (101%)	173 (99%)	1 (1%)	89	92
1	B	172/172 (100%)	169 (98%)	3 (2%)	66	70
All	All	346/344 (101%)	342 (99%)	4 (1%)	75	80

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	250	ASP
1	B	229	LEU
1	B	250	ASP
1	B	385	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

49 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	A	401	2	12,12,12	0.64	0	17,17,17	1.02	0
2	GLC	A	402	2	11,11,12	0.88	0	13,15,17	1.02	1 (7%)
2	GLC	A	403	2	11,11,12	1.07	0	13,15,17	1.05	0
2	GLC	A	404	2	11,11,12	0.95	0	13,15,17	1.11	1 (7%)
2	GLC	A	405	2	12,12,12	1.02	1 (8%)	17,17,17	0.81	0
2	GLC	A	406	2	11,11,12	0.94	0	13,15,17	0.84	1 (7%)
2	GLC	A	407	2	11,11,12	0.55	0	13,15,17	1.04	1 (7%)
3	EDO	A	408	-	3,3,3	0.47	0	2,2,2	0.30	0
3	EDO	A	409	-	3,3,3	0.47	0	2,2,2	0.35	0
3	EDO	A	410	-	3,3,3	0.45	0	2,2,2	0.21	0
3	EDO	A	411	-	3,3,3	0.47	0	2,2,2	0.30	0
3	EDO	A	412	-	3,3,3	0.42	0	2,2,2	0.50	0
3	EDO	A	413	-	3,3,3	0.50	0	2,2,2	0.30	0
3	EDO	A	414	-	3,3,3	0.48	0	2,2,2	0.34	0
3	EDO	A	415	-	3,3,3	0.45	0	2,2,2	0.35	0
4	FMT	A	416	-	2,2,2	0.70	0	1,1,1	0.43	0
4	FMT	A	417	-	2,2,2	0.74	0	1,1,1	0.52	0
4	FMT	A	418	-	2,2,2	0.75	0	1,1,1	0.44	0
4	FMT	A	419	-	2,2,2	0.69	0	1,1,1	0.44	0
4	FMT	A	420	-	2,2,2	0.72	0	1,1,1	0.50	0
4	FMT	A	421	-	2,2,2	0.70	0	1,1,1	0.44	0
4	FMT	A	422	-	2,2,2	0.71	0	1,1,1	0.46	0
4	FMT	A	423	-	2,2,2	0.76	0	1,1,1	0.50	0
4	FMT	A	424	-	2,2,2	0.67	0	1,1,1	0.44	0
2	GLC	B	401	2	12,12,12	0.83	0	17,17,17	0.76	0
2	GLC	B	402	2	11,11,12	1.25	1 (9%)	13,15,17	1.35	1 (7%)
2	GLC	B	403	2	11,11,12	1.27	1 (9%)	13,15,17	0.77	1 (7%)
2	GLC	B	404	2	12,12,12	0.55	0	17,17,17	1.14	2 (11%)
2	GLC	B	405	2	11,11,12	0.94	0	13,15,17	1.05	1 (7%)
2	GLC	B	406	2	11,11,12	1.05	0	13,15,17	1.23	2 (15%)
2	GLC	B	407	2	11,11,12	1.02	1 (9%)	13,15,17	0.89	0
3	EDO	B	408	-	3,3,3	0.45	0	2,2,2	0.30	0
3	EDO	B	409	-	3,3,3	0.46	0	2,2,2	0.33	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	B	410	-	3,3,3	0.45	0	2,2,2	0.32	0
3	EDO	B	411	-	3,3,3	0.45	0	2,2,2	0.36	0
3	EDO	B	412	-	3,3,3	0.46	0	2,2,2	0.29	0
3	EDO	B	413	-	3,3,3	0.50	0	2,2,2	0.25	0
3	EDO	B	414	-	3,3,3	0.40	0	2,2,2	0.67	0
3	EDO	B	415	-	3,3,3	0.45	0	2,2,2	0.33	0
4	FMT	B	416	-	2,2,2	0.71	0	1,1,1	0.50	0
4	FMT	B	417	-	2,2,2	0.78	0	1,1,1	0.47	0
4	FMT	B	418	-	2,2,2	0.70	0	1,1,1	0.46	0
4	FMT	B	419	-	2,2,2	0.70	0	1,1,1	0.45	0
4	FMT	B	420	-	2,2,2	0.71	0	1,1,1	0.50	0
4	FMT	B	421	-	2,2,2	0.67	0	1,1,1	0.43	0
4	FMT	B	422	-	2,2,2	0.69	0	1,1,1	0.45	0
4	FMT	B	423	-	2,2,2	0.70	0	1,1,1	0.46	0
4	FMT	B	424	-	2,2,2	0.73	0	1,1,1	0.48	0
4	FMT	B	425	-	2,2,2	0.62	0	1,1,1	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	401	2	-	0/2/22/22	0/1/1/1
2	GLC	A	402	2	-	0/2/19/22	0/1/1/1
2	GLC	A	403	2	-	0/2/19/22	0/1/1/1
2	GLC	A	404	2	-	0/2/19/22	0/1/1/1
2	GLC	A	405	2	-	0/2/22/22	0/1/1/1
2	GLC	A	406	2	-	0/2/19/22	0/1/1/1
2	GLC	A	407	2	-	0/2/19/22	0/1/1/1
3	EDO	A	408	-	-	0/1/1/1	0/0/0/0
3	EDO	A	409	-	-	0/1/1/1	0/0/0/0
3	EDO	A	410	-	-	0/1/1/1	0/0/0/0
3	EDO	A	411	-	-	0/1/1/1	0/0/0/0
3	EDO	A	412	-	-	0/1/1/1	0/0/0/0
3	EDO	A	413	-	-	0/1/1/1	0/0/0/0
3	EDO	A	414	-	-	0/1/1/1	0/0/0/0
3	EDO	A	415	-	-	0/1/1/1	0/0/0/0
4	FMT	A	416	-	-	0/0/0/0	0/0/0/0
4	FMT	A	417	-	-	0/0/0/0	0/0/0/0
4	FMT	A	418	-	-	0/0/0/0	0/0/0/0
4	FMT	A	419	-	-	0/0/0/0	0/0/0/0
4	FMT	A	420	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FMT	A	421	-	-	0/0/0/0	0/0/0/0
4	FMT	A	422	-	-	0/0/0/0	0/0/0/0
4	FMT	A	423	-	-	0/0/0/0	0/0/0/0
4	FMT	A	424	-	-	0/0/0/0	0/0/0/0
2	GLC	B	401	2	-	0/2/22/22	0/1/1/1
2	GLC	B	402	2	-	0/2/19/22	0/1/1/1
2	GLC	B	403	2	-	0/2/19/22	0/1/1/1
2	GLC	B	404	2	-	0/2/22/22	0/1/1/1
2	GLC	B	405	2	-	0/2/19/22	0/1/1/1
2	GLC	B	406	2	-	0/2/19/22	0/1/1/1
2	GLC	B	407	2	-	0/2/19/22	0/1/1/1
3	EDO	B	408	-	-	0/1/1/1	0/0/0/0
3	EDO	B	409	-	-	0/1/1/1	0/0/0/0
3	EDO	B	410	-	-	0/1/1/1	0/0/0/0
3	EDO	B	411	-	-	0/1/1/1	0/0/0/0
3	EDO	B	412	-	-	0/1/1/1	0/0/0/0
3	EDO	B	413	-	-	0/1/1/1	0/0/0/0
3	EDO	B	414	-	-	0/1/1/1	0/0/0/0
3	EDO	B	415	-	-	0/1/1/1	0/0/0/0
4	FMT	B	416	-	-	0/0/0/0	0/0/0/0
4	FMT	B	417	-	-	0/0/0/0	0/0/0/0
4	FMT	B	418	-	-	0/0/0/0	0/0/0/0
4	FMT	B	419	-	-	0/0/0/0	0/0/0/0
4	FMT	B	420	-	-	0/0/0/0	0/0/0/0
4	FMT	B	421	-	-	0/0/0/0	0/0/0/0
4	FMT	B	422	-	-	0/0/0/0	0/0/0/0
4	FMT	B	423	-	-	0/0/0/0	0/0/0/0
4	FMT	B	424	-	-	0/0/0/0	0/0/0/0
4	FMT	B	425	-	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	403	GLC	O5-C1	-2.84	1.39	1.43
2	B	402	GLC	O5-C1	-2.74	1.39	1.43
2	B	407	GLC	O5-C1	-2.48	1.39	1.43
2	A	405	GLC	O5-C5	-2.09	1.39	1.44

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	406	GLC	O5-C1-C2	-2.86	106.31	110.79

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	407	GLC	O5-C1-C2	-2.51	106.86	110.79
2	A	402	GLC	C1-O5-C5	2.01	114.94	112.17
2	B	403	GLC	C1-O5-C5	2.03	114.96	112.17
2	A	406	GLC	C1-O5-C5	2.04	114.98	112.17
2	B	404	GLC	C1-O5-C5	2.11	117.20	113.39
2	B	405	GLC	C1-O5-C5	2.20	115.19	112.17
2	B	404	GLC	O5-C5-C4	2.43	114.13	109.66
2	A	404	GLC	C1-O5-C5	2.67	115.85	112.17
2	B	406	GLC	C1-O5-C5	3.22	116.61	112.17
2	B	402	GLC	C1-O5-C5	3.80	117.41	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	405	GLC	0	1
2	B	403	GLC	0	1
2	B	407	GLC	0	1
4	B	423	FMT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	209/209 (100%)	0.53	19 (9%)	10 10	23, 33, 67, 97	0
1	B	209/209 (100%)	0.32	12 (5%)	24 25	24, 34, 59, 93	0
All	All	418/418 (100%)	0.43	31 (7%)	15 15	23, 34, 64, 97	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	323	ALA	7.0
1	B	323	ALA	6.0
1	B	326	LEU	5.7
1	A	326	LEU	5.6
1	A	384	ASP	5.3
1	A	311	GLY	4.2
1	A	324	ASP	3.8
1	A	387	GLU	3.7
1	A	289	GLY	3.5
1	B	384	ASP	3.4
1	B	387	GLU	3.3
1	A	375	ASP	3.3
1	A	293	THR	3.2
1	A	292	SER	3.1
1	B	324	ASP	3.1
1	A	327	GLY	3.0
1	B	289	GLY	3.0
1	A	283	ILE	3.0
1	A	349	ASN	2.9
1	B	321	ALA	2.9
1	B	327	GLY	2.7
1	A	379	TYR	2.7
1	A	320	SER	2.6
1	A	385	SER	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	349	ASN	2.4
1	A	348	ASN	2.4
1	B	292	SER	2.3
1	B	322	ASN	2.3
1	A	321	ALA	2.1
1	B	220	GLU	2.1
1	A	288	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	EDO	A	409	4/4	0.81	0.25	8.45	32,54,76,76	0
4	FMT	A	420	3/3	0.97	0.26	7.35	18,21,59,71	0
3	EDO	B	413	4/4	0.63	0.32	6.57	55,67,78,78	0
4	FMT	B	424	3/3	0.96	0.23	6.46	22,27,54,65	0
4	FMT	B	416	3/3	0.97	0.27	5.53	20,24,57,60	0
3	EDO	A	414	4/4	0.79	0.22	5.47	32,48,67,67	0
4	FMT	B	425	3/3	0.92	0.20	5.40	33,37,68,82	0
3	EDO	A	415	4/4	0.79	0.22	4.84	43,55,84,84	0
3	EDO	A	413	4/4	0.64	0.28	4.06	49,65,102,102	0
3	EDO	A	412	4/4	0.86	0.27	4.02	36,48,67,67	0
3	EDO	B	411	4/4	0.81	0.44	3.67	59,71,78,78	0
4	FMT	A	417	3/3	0.99	0.21	3.32	11,14,56,68	0
3	EDO	B	414	4/4	0.91	0.28	3.16	21,36,53,53	0
4	FMT	B	423	3/3	0.97	0.18	2.62	18,22,81,97	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	EDO	B	409	4/4	0.82	0.22	2.33	38,53,64,64	0
3	EDO	B	410	4/4	0.91	0.30	2.19	34,52,56,63	0
3	EDO	A	411	4/4	0.79	0.28	1.95	51,67,70,81	0
3	EDO	B	408	4/4	0.92	0.14	1.52	42,51,55,67	0
4	FMT	A	422	3/3	0.96	0.17	1.50	20,24,51,61	0
4	FMT	B	421	3/3	0.95	0.16	1.50	29,35,56,67	0
3	EDO	B	415	4/4	0.82	0.30	1.35	62,118,157,157	0
4	FMT	B	418	3/3	0.93	0.14	0.87	36,51,64,77	0
3	EDO	A	408	4/4	0.85	0.20	0.36	53,63,73,74	0
2	GLC	A	404	11/12	0.90	0.13	0.24	29,37,46,47	1
2	GLC	B	403	11/12	0.93	0.12	0.06	31,40,51,53	0
2	GLC	B	401	12/12	0.94	0.12	-0.39	25,39,48,49	1
4	FMT	A	421	3/3	0.95	0.12	-0.59	44,46,55,60	0
3	EDO	A	410	4/4	0.96	0.09	-0.62	36,45,55,62	0
2	GLC	A	406	11/12	0.94	0.10	-0.88	34,38,43,46	0
2	GLC	A	405	12/12	0.93	0.10	-0.96	34,43,53,60	1
2	GLC	B	406	11/12	0.92	0.09	-1.08	29,34,39,45	0
2	GLC	B	402	11/12	0.96	0.09	-1.44	25,30,34,40	0
2	GLC	A	403	11/12	0.96	0.08	-1.95	25,30,36,41	0
4	FMT	B	419	3/3	0.91	0.27	-	61,64,76,83	0
2	GLC	B	405	11/12	0.94	0.10	-	26,34,43,49	0
4	FMT	B	422	3/3	0.79	0.14	-	61,61,76,91	0
2	GLC	A	407	11/12	0.83	0.34	-	56,76,90,91	0
4	FMT	B	420	3/3	0.93	0.23	-	43,55,66,71	0
2	GLC	B	404	12/12	0.74	0.36	-	38,79,98,108	0
4	FMT	B	417	3/3	0.96	0.18	-	24,29,38,59	0
4	FMT	A	419	3/3	0.97	0.21	-	19,23,52,62	0
4	FMT	A	418	3/3	0.96	0.21	-	15,18,44,45	0
2	GLC	A	401	12/12	0.82	0.34	-	33,69,88,100	0
2	GLC	A	402	11/12	0.94	0.10	-	24,31,38,41	0
4	FMT	A	424	3/3	0.97	0.18	-	25,30,59,71	0
2	GLC	B	407	11/12	0.88	0.14	-	33,42,51,58	1
3	EDO	B	412	4/4	0.88	0.18	-	48,57,67,67	0
4	FMT	A	423	3/3	0.96	0.23	-	21,26,58,70	0
4	FMT	A	416	3/3	0.89	0.22	-	55,61,73,74	0

6.5 Other polymers [i](#)

There are no such residues in this entry.