



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:40 pm GMT

PDB ID : 1B5F
Title : NATIVE CARDOSIN A FROM CYNARA CARDUNCULUS L.
Authors : Frazao, C.; Bento, I.; Carrondo, M.A.
Deposited on : 1999-01-06
Resolution : 1.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	4841 (1.74-1.70)
Ramachandran outliers	110173	4771 (1.74-1.70)
Sidechain outliers	110143	4771 (1.74-1.70)
RSRZ outliers	101464	4426 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	239	<div> <div>8%</div> <div>82%</div> <div>16%</div> <div>.</div> </div>
1	C	239	<div> <div>10%</div> <div>75%</div> <div>21%</div> <div>.</div> </div>
2	B	87	<div> <div>7%</div> <div>83%</div> <div>15%</div> <div>.</div> </div>
2	D	87	<div> <div>7%</div> <div>89%</div> <div>10%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	403	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 5842 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (CARDOSIN A).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	239	Total	C	N	O	S	0	7	0
			1870	1188	305	369	8			
1	C	238	Total	C	N	O	S	0	8	0
			1870	1189	303	372	6			

- Molecule 2 is a protein called PROTEIN (CARDOSIN A).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	87	Total	C	N	O	S	0	1	0
			668	432	103	128	5			
2	D	87	Total	C	N	O	S	0	0	0
			663	429	103	126	5			

- Molecule 3 is a polymer of unknown type called SUGAR (NAG-FUC-NAG-MAN-MAN).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	5	Total	C	N	O	0	0
			60	34	2	24		

- Molecule 4 is a polymer of unknown type called SUGAR (NAG-FUC-NAG-MAN).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	4	Total	C	N	O	0	1
			53	30	3	20		

- Molecule 5 is a polymer of unknown type called SUGAR (NAG-FUC-NAG-MAN-MAN-MAN).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	6	Total	C	N	O	0	0
			71	40	2	29		

- Molecule 6 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	D	4	Total	C	N	O	0	1
			59	34	2	23		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	199	Total	O	0	0
			199	199		
7	B	71	Total	O	0	0
			71	71		
7	C	192	Total	O	0	0
			192	192		
7	D	66	Total	O	0	0
			66	66		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PROTEIN (CARDOSIN A)




• Molecule 1: PROTEIN (CARDOSIN A)



• Molecule 2: PROTEIN (CARDOSIN A)



• Molecule 2: PROTEIN (CARDOSIN A)



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	116.96Å 87.19Å 81.30Å 90.00° 104.43° 90.00°	Depositor
Resolution (Å)	34.80 – 1.72 34.78 – 1.72	Depositor EDS
% Data completeness (in resolution range)	99.8 (34.80-1.72) 95.6 (34.78-1.72)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.25 (at 1.72Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.206 , 0.256 0.195 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	19.1	Xtriage
Anisotropy	0.285	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 76.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5842	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/1932	1.08	3/2623 (0.1%)
1	C	0.42	0/1944	1.12	2/2639 (0.1%)
2	B	0.42	0/687	1.12	1/930 (0.1%)
2	D	0.44	0/678	1.02	2/918 (0.2%)
All	All	0.42	0/5241	1.09	8/7110 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	204	THR	C-N-CA	6.38	135.69	122.30
1	A	149	ARG	CD-NE-CZ	6.17	132.23	123.60
2	D	274	TYR	CB-CG-CD1	5.97	124.58	121.00
1	A	138	TYR	CA-CB-CG	5.77	124.36	113.40
2	D	274	TYR	CA-CB-CG	5.37	123.61	113.40
2	B	274	TYR	CA-CB-CG	5.12	123.12	113.40
1	C	176	ARG	CD-NE-CZ	5.08	130.72	123.60
1	A	176	ARG	NE-CZ-NH1	5.03	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1870	0	1763	27	0
1	C	1870	0	1762	41	0
2	B	668	0	651	9	0
2	D	663	0	649	5	0
3	A	60	0	52	0	0
4	B	53	0	42	1	0
5	C	71	0	61	4	0
6	D	59	0	53	6	0
7	A	199	0	0	5	0
7	B	71	0	0	2	0
7	C	192	0	0	7	0
7	D	66	0	0	1	0
All	All	5842	0	5033	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:225:THR:O	1:C:229[A]:THR:HG22	1.89	0.72
4:B:501[A]:NAG:H82	7:B:1106:HOH:O	1.93	0.68
2:B:245:LEU:HG	7:B:1389:HOH:O	1.94	0.68
1:C:229[B]:THR:HG21	7:C:1198:HOH:O	1.94	0.68
1:C:238:ASN:OD1	2:D:245:LEU:HA	1.96	0.66
1:A:45[A]:CYS:SG	1:A:47[A]:SER:HB2	2.37	0.65
1:C:147:LYS:HE3	7:C:1386:HOH:O	1.97	0.62
1:C:137:TRP:CZ2	1:C:141:LEU:HD21	2.35	0.61
1:A:137:TRP:CZ2	1:A:141:LEU:HD21	2.38	0.59
1:A:66:GLU:HG2	7:A:1078:HOH:O	2.03	0.59
1:A:184:VAL:HA	1:A:191:GLN:O	2.04	0.58
6:D:501:NAG:O4	6:D:503:NAG:H83	2.05	0.57
1:C:137:TRP:NE1	1:C:141:LEU:HD11	2.20	0.57
1:C:238:ASN:HB2	2:D:245:LEU:HD22	1.86	0.57
1:A:37:VAL:HG22	7:A:1400:HOH:O	2.06	0.54
6:D:504:MAN:O2	6:D:504:MAN:H62	2.07	0.54
1:A:137:TRP:NE1	1:A:141:LEU:HD11	2.23	0.54
1:A:109:ASP:O	1:A:113:LEU:HB3	2.08	0.53
1:C:160(A)[B]:GLU:O	1:C:160(B):GLU:HG3	2.08	0.53
1:A:218:THR:HG21	7:A:1379:HOH:O	2.09	0.53
1:C:48:LYS:HD2	1:C:113:LEU:CD2	2.39	0.52
1:A:48:LYS:HG3	1:A:48:LYS:O	2.11	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:114:HIS:HB2	7:C:1387:HOH:O	2.09	0.50
1:C:238:ASN:CG	2:D:246:GLN:H	2.15	0.50
1:A:237:ALA:O	1:A:238:ASN:HB3	2.12	0.50
2:B:244:GLU:O	2:B:244:GLU:HG3	2.12	0.50
5:C:401:NAG:O4	5:C:403:NAG:O7	2.30	0.49
1:C:73:ILE:O	1:C:79:SER:HB2	2.12	0.49
1:C:72:ALA:CB	1:C:81:THR:HG22	2.43	0.48
6:D:501:NAG:H4	6:D:502[B]:FUC:O5	2.13	0.48
1:C:159:VAL:CG1	1:C:160(A)[B]:GLU:HG3	2.44	0.48
1:C:157:ARG:HB3	7:C:1190:HOH:O	2.14	0.48
1:A:26:LYS:HG3	7:A:1420:HOH:O	2.13	0.48
1:A:48:LYS:O	1:A:51:ARG:HG2	2.13	0.48
1:C:78:GLY:HA3	1:C:111:VAL:HG21	1.96	0.47
1:C:39:TRP:CD2	1:C:80:ILE:HG12	2.49	0.47
1:A:137:TRP:CE2	1:A:141:LEU:HD11	2.50	0.47
1:A:224:PRO:HD2	2:B:299:TRP:CE2	2.49	0.47
1:C:231:ILE:O	1:C:235:ILE:HG12	2.14	0.47
6:D:501:NAG:O7	6:D:502[B]:FUC:H5	2.14	0.47
2:B:244:GLU:OE1	2:B:244:GLU:HA	2.15	0.47
1:C:132:ILE:HD12	1:C:133:SER:N	2.29	0.46
2:D:243:GLU:O	2:D:244:GLU:HB2	2.16	0.46
1:C:39:TRP:HE1	1:C:75:TYR:HH	1.62	0.45
1:A:75:TYR:N	1:A:78:GLY:O	2.49	0.45
5:C:401:NAG:H83	5:C:402:FUC:O2	2.16	0.45
1:C:159:VAL:HG13	1:C:160(A)[B]:GLU:HG3	1.98	0.45
1:C:176:ARG:HD2	7:D:1156:HOH:O	2.17	0.45
1:A:114:HIS:O	1:A:115:ARG:HD2	2.17	0.45
2:B:278(B):LYS:O	2:B:280:GLU:N	2.50	0.45
1:A:141:LEU:HD23	1:A:146:VAL:CG2	2.46	0.44
1:C:60:ASP:OD1	1:C:60:ASP:N	2.50	0.44
1:A:172:PRO:HA	1:A:175:PHE:CE1	2.52	0.44
2:B:245:LEU:N	2:B:245:LEU:HD23	2.32	0.44
1:C:10:ARG:NE	7:C:1452:HOH:O	2.50	0.44
1:A:157:ARG:NH1	1:A:157:ARG:HG3	2.32	0.44
1:A:30:ILE:HD11	1:A:117:PHE:CE2	2.52	0.44
1:C:99:GLN:OE1	1:C:136:VAL:HA	2.18	0.44
1:A:9:ASP:OD2	1:A:10:ARG:NH2	2.50	0.44
1:C:75:TYR:N	1:C:78:GLY:O	2.49	0.44
1:A:157:ARG:NH2	7:A:1477:HOH:O	2.50	0.44
1:A:160:ASP:OD1	1:A:160(A):GLU:N	2.51	0.43
1:C:75:TYR:HB2	1:C:78:GLY:O	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:161:GLU:HB3	7:C:1211:HOH:O	2.17	0.43
1:C:17[B]:GLU:OE2	1:C:28:THR:OG1	2.29	0.43
6:D:501:NAG:H2	6:D:502[B]:FUC:O5	2.17	0.43
1:C:169:GLY:C	1:C:170:LEU:HD23	2.39	0.43
1:A:0:GLY:HA2	1:A:147:LYS:HE3	2.01	0.43
5:C:401:NAG:H83	5:C:402:FUC:C1	2.49	0.42
1:C:199:ILE:O	1:C:202[A]:LYS:HB3	2.19	0.42
6:D:501:NAG:H2	6:D:502[B]:FUC:C5	2.49	0.42
2:B:282:CYS:C	2:B:283:ILE:HD12	2.40	0.42
1:A:113:LEU:HG	1:A:114:HIS:ND1	2.34	0.42
1:C:66:GLU:HG3	1:C:67:ASN:N	2.24	0.42
1:C:48:LYS:HD2	1:C:113:LEU:HD21	2.01	0.42
1:A:1:SER:OG	1:A:147:LYS:HG2	2.19	0.42
1:C:87:ASP:OD1	1:C:88:SER:N	2.51	0.42
1:C:160:ASP:HA	7:C:1291:HOH:O	2.20	0.41
2:B:293:LEU:O	2:B:295:LEU:HB2	2.21	0.41
1:C:100:ASP:OD2	5:C:401:NAG:O6	2.30	0.41
1:C:108:ALA:HB1	1:C:110:ASN:ND2	2.35	0.41
1:C:17[B]:GLU:OE1	1:C:27:PHE:O	2.39	0.41
1:C:45:CYS:SG	1:C:47:SER:HB3	2.60	0.41
2:D:248:ASP:OD1	2:D:250:ASN:HB2	2.21	0.41
1:C:75:TYR:CZ	1:C:120:ILE:HD12	2.55	0.40
1:A:180:THR:O	2:B:321:GLY:HA2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	244/239 (102%)	231 (95%)	12 (5%)	1 (0%)	38	20
1	C	244/239 (102%)	235 (96%)	9 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	86/87 (99%)	83 (96%)	2 (2%)	1 (1%)	15	3
2	D	85/87 (98%)	81 (95%)	3 (4%)	1 (1%)	15	3
All	All	659/652 (101%)	630 (96%)	26 (4%)	3 (0%)	32	14

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	279	GLY
2	D	279	GLY
1	A	1	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/199 (104%)	196 (95%)	10 (5%)	29	9
1	C	207/199 (104%)	193 (93%)	14 (7%)	18	4
2	B	72/71 (101%)	67 (93%)	5 (7%)	18	4
2	D	71/71 (100%)	68 (96%)	3 (4%)	34	13
All	All	556/540 (103%)	524 (94%)	32 (6%)	24	7

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	LYS
1	A	44	LYS
1	A	46(A)	ASN
1	A	66	GLU
1	A	77	THR
1	A	106	ASP
1	A	147	LYS
1	A	160	ASP
1	A	198	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	202	LYS
2	B	278(B)	LYS
2	B	281	GLN
2	B	289	MET
2	B	293	LEU
2	B	318	LEU
1	C	45	CYS
1	C	48	LYS
1	C	51	ARG
1	C	63	THR
1	C	66	GLU
1	C	77	THR
1	C	80	ILE
1	C	103	GLU
1	C	106	ASP
1	C	109	ASP
1	C	110	ASN
1	C	115	ARG
1	C	160(A)[A]	GLU
1	C	160(A)[B]	GLU
2	D	244	GLU
2	D	278(B)	LYS
2	D	293	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	187	GLN
1	A	191	GLN
1	A	211	GLN
2	B	317	ASN
1	C	191	GLN
1	C	211	GLN
2	D	317	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

21 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	401	1,3	14,14,15	0.77	0	15,19,21	1.57	4 (26%)
3	FUC	A	402	3	9,10,11	0.55	0	13,14,16	0.95	1 (7%)
3	NAG	A	403	3	14,14,15	1.02	1 (7%)	15,19,21	1.66	3 (20%)
3	BMA	A	404	3	11,11,12	1.12	1 (9%)	13,15,17	1.98	5 (38%)
3	MAN	A	405	3	11,11,12	0.79	0	13,15,17	1.89	4 (30%)
4	NAG	B	501[A]	-	14,14,15	0.88	0	15,19,21	0.97	1 (6%)
4	NAG	B	501[B]	-	14,14,15	0.89	0	15,19,21	0.63	0
4	FUC	B	502	4	9,10,11	0.60	0	13,14,16	1.02	0
4	NAG	B	503	4	14,14,15	0.87	1 (7%)	15,19,21	1.20	2 (13%)
4	BMA	B	504	4	11,11,12	0.52	0	13,15,17	1.88	3 (23%)
5	NAG	C	401	1,5	14,14,15	0.76	0	15,19,21	1.80	2 (13%)
5	FUC	C	402	5	9,10,11	0.73	0	13,14,16	1.02	0
5	NAG	C	403	5	14,14,15	0.70	0	15,19,21	1.03	1 (6%)
5	BMA	C	404	5	11,11,12	0.77	0	13,15,17	1.36	1 (7%)
5	MAN	C	405	5	11,11,12	0.88	0	13,15,17	1.58	2 (15%)
5	MAN	C	406	5	11,11,12	0.77	0	13,15,17	2.51	4 (30%)
6	NAG	D	501	2,6	14,14,15	0.88	0	15,19,21	1.27	2 (13%)
6	FUC	D	502[A]	6	9,10,11	0.72	0	13,14,16	1.14	1 (7%)
6	FUC	D	502[B]	6	9,10,11	0.89	0	13,14,16	2.02	3 (23%)
6	NAG	D	503	6	14,14,15	0.88	1 (7%)	15,19,21	1.19	1 (6%)
6	MAN	D	504	6	11,11,12	0.97	0	13,15,17	2.19	4 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	401	1,3	-	0/6/23/26	0/1/1/1
3	FUC	A	402	3	-	0/0/17/20	0/1/1/1
3	NAG	A	403	3	-	0/6/23/26	0/1/1/1
3	BMA	A	404	3	-	0/2/19/22	0/1/1/1
3	MAN	A	405	3	-	0/2/19/22	0/1/1/1
4	NAG	B	501[A]	-	-	0/6/23/26	0/1/1/1
4	NAG	B	501[B]	-	-	0/6/23/26	0/1/1/1
4	FUC	B	502	4	-	0/0/17/20	0/1/1/1
4	NAG	B	503	4	-	0/6/23/26	0/1/1/1
4	BMA	B	504	4	-	0/2/19/22	0/1/1/1
5	NAG	C	401	1,5	-	0/6/23/26	0/1/1/1
5	FUC	C	402	5	-	0/0/17/20	0/1/1/1
5	NAG	C	403	5	-	0/6/23/26	0/1/1/1
5	BMA	C	404	5	-	0/2/19/22	0/1/1/1
5	MAN	C	405	5	-	0/2/19/22	0/1/1/1
5	MAN	C	406	5	-	0/2/19/22	0/1/1/1
6	NAG	D	501	2,6	-	0/6/23/26	0/1/1/1
6	FUC	D	502[A]	6	-	0/0/17/20	0/1/1/1
6	FUC	D	502[B]	6	-	0/0/17/20	0/1/1/1
6	NAG	D	503	6	-	0/6/23/26	0/1/1/1
6	MAN	D	504	6	-	0/2/19/22	1/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	403	NAG	O5-C1	-2.99	1.38	1.43
6	D	503	NAG	O5-C1	-2.43	1.39	1.43
4	B	503	NAG	O5-C1	-2.01	1.40	1.43
3	A	404	BMA	C2-C3	2.81	1.56	1.52

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	405	MAN	C1-C2-C3	-4.89	103.45	109.65
5	C	406	MAN	C1-C2-C3	-4.32	104.18	109.65
6	D	504	MAN	C1-C2-C3	-4.06	104.50	109.65
5	C	405	MAN	C1-C2-C3	-3.60	105.08	109.65
3	A	401	NAG	C2-N2-C7	-3.52	117.81	122.94
3	A	404	BMA	C2-C3-C4	-3.50	104.77	110.88

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	404	BMA	O3-C3-C4	-3.44	102.88	110.36
4	B	504	BMA	C1-C2-C3	-3.11	105.71	109.65
5	C	404	BMA	O3-C3-C4	-3.00	103.84	110.36
6	D	502[A]	FUC	C2-C3-C4	-2.98	105.67	110.88
3	A	404	BMA	C1-C2-C3	-2.95	105.91	109.65
3	A	403	NAG	C8-C7-N2	-2.91	110.86	116.11
5	C	406	MAN	O5-C1-C2	-2.78	106.44	110.79
6	D	501	NAG	C2-N2-C7	-2.69	119.02	122.94
5	C	405	MAN	O5-C1-C2	-2.68	106.60	110.79
3	A	405	MAN	O5-C1-C2	-2.39	107.05	110.79
3	A	401	NAG	O3-C3-C4	-2.38	105.17	110.36
5	C	401	NAG	C4-C3-C2	-2.38	107.53	111.02
3	A	404	BMA	O2-C2-C3	-2.24	105.77	110.17
4	B	503	NAG	C2-N2-C7	-2.18	119.77	122.94
4	B	501[A]	NAG	O7-C7-N2	2.02	125.81	121.92
3	A	401	NAG	C1-C2-N2	2.11	114.10	110.49
5	C	406	MAN	O3-C3-C2	2.15	113.93	110.02
4	B	504	BMA	O3-C3-C2	2.21	114.05	110.02
3	A	401	NAG	C1-O5-C5	2.41	115.49	112.17
5	C	403	NAG	C1-O5-C5	2.44	115.53	112.17
6	D	504	MAN	O3-C3-C2	2.46	114.50	110.02
3	A	402	FUC	C1-C2-C3	2.51	112.83	109.65
6	D	502[B]	FUC	C1-O5-C5	2.55	118.04	112.39
3	A	405	MAN	C6-C5-C4	2.59	119.07	113.00
3	A	404	BMA	C1-O5-C5	2.67	115.84	112.17
3	A	403	NAG	O3-C3-C4	2.67	116.17	110.36
4	B	503	NAG	C4-C3-C2	2.75	115.04	111.02
3	A	405	MAN	O2-C2-C3	2.78	115.63	110.17
6	D	501	NAG	C1-O5-C5	2.93	116.20	112.17
6	D	502[B]	FUC	O5-C1-C2	3.13	115.70	110.79
6	D	503	NAG	C1-O5-C5	3.64	117.19	112.17
3	A	403	NAG	C1-O5-C5	3.76	117.34	112.17
6	D	504	MAN	C1-O5-C5	4.01	117.69	112.17
6	D	504	MAN	O2-C2-C3	4.33	118.68	110.17
4	B	504	BMA	O2-C2-C3	4.97	119.94	110.17
5	C	401	NAG	C1-O5-C5	5.16	119.27	112.17
5	C	406	MAN	O2-C2-C3	5.67	121.31	110.17
6	D	502[B]	FUC	C1-C2-C3	5.72	116.91	109.65

There are no chirality outliers.

There are no torsion outliers.

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	D	504	MAN	C1-C2-C3-C4-C5-O5

8 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	501[A]	NAG	1	0
5	C	401	NAG	4	0
5	C	402	FUC	2	0
5	C	403	NAG	1	0
6	D	501	NAG	5	0
6	D	502[B]	FUC	4	0
6	D	503	NAG	1	0
6	D	504	MAN	1	0

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	239/239 (100%)	0.30	20 (8%)	12	14	15, 24, 69, 132	0
1	C	238/239 (99%)	0.52	23 (9%)	8	10	15, 27, 64, 132	0
2	B	87/87 (100%)	0.02	6 (6%)	18	21	14, 21, 63, 89	0
2	D	87/87 (100%)	0.01	6 (6%)	18	21	13, 22, 63, 97	0
All	All	651/652 (99%)	0.30	55 (8%)	12	14	13, 25, 68, 132	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	46	ILE	9.9
1	A	77	THR	7.0
2	B	279	GLY	6.3
1	A	46	ILE	6.0
1	A	160	ASP	6.0
2	D	279	GLY	5.5
1	C	74	ILE	5.4
1	A	76	GLY	5.4
2	B	280	GLU	5.0
1	C	46(A)	ASN	4.9
1	A	74	ILE	4.8
2	D	243	GLU	4.3
1	A	238	ASN	4.2
1	A	111	VAL	4.1
1	C	110	ASN	4.1
1	C	70	PHE	4.0
1	C	111	VAL	3.9
1	C	77	THR	3.8
1	A	78	GLY	3.7
1	A	114	HIS	3.7
1	C	76	GLY	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	114	HIS	3.6
1	C	160(A)[A]	GLU	3.6
1	C	80	ILE	3.6
1	A	51	ARG	3.5
1	A	110	ASN	3.4
2	D	280(A)	ALA	3.4
1	C	109	ASP	3.4
1	A	75	TYR	3.3
2	B	278(B)	LYS	3.3
1	C	78	GLY	3.3
2	B	243	GLU	3.2
2	D	280	GLU	3.1
1	C	108	ALA	3.1
1	C	112	PHE	3.0
2	D	244	GLU	3.0
1	A	46(A)	ASN	2.9
2	B	244	GLU	2.8
1	C	238	ASN	2.8
2	D	278(B)	LYS	2.7
1	A	113	LEU	2.7
1	C	48	LYS	2.6
1	A	45[A]	CYS	2.6
1	A	0	GLY	2.5
1	A	159	VAL	2.5
1	A	160(A)	GLU	2.5
1	C	66	GLU	2.4
1	C	64	TYR	2.4
1	A	48	LYS	2.4
1	A	18	ILE	2.2
1	C	50	CYS	2.2
1	C	47	SER	2.2
1	C	45	CYS	2.1
2	B	278(A)	GLY	2.1
1	C	59[A]	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	A	403	14/15	0.91	0.13	2.38	28,35,51,79	0
5	MAN	C	405	11/12	0.87	0.15	1.49	27,34,40,44	0
5	NAG	C	401	14/15	0.81	0.17	1.13	32,47,65,87	0
3	NAG	A	401	14/15	0.93	0.10	0.28	21,30,53,53	0
4	NAG	B	503	14/15	0.86	0.21	-	40,59,87,94	0
4	NAG	B	501[A]	14/15	0.90	0.16	-	29,41,54,57	4
3	MAN	A	405	11/12	0.79	0.20	-	46,52,63,79	0
5	NAG	C	403	14/15	0.93	0.11	-	24,33,70,80	0
6	MAN	D	504	11/12	0.36	0.46	-	134,142,148,153	0
6	FUC	D	502[A]	10/11	0.57	0.40	-	54,79,103,105	10
4	BMA	B	504	11/12	0.44	0.31	-	99,103,112,136	0
5	MAN	C	406	11/12	0.44	0.44	-	105,124,135,137	0
5	FUC	C	402	10/11	0.82	0.27	-	50,57,80,81	0
6	FUC	D	502[B]	10/11	0.57	0.40	-	45,69,79,82	10
3	BMA	A	404	11/12	0.81	0.17	-	38,47,54,60	0
4	NAG	B	501[B]	14/15	0.90	0.16	-	29,41,54,57	4
4	FUC	B	502	10/11	0.76	0.23	-	65,75,82,89	0
5	BMA	C	404	11/12	0.90	0.12	-	27,36,56,89	0
3	FUC	A	402	10/11	0.80	0.17	-	42,54,60,86	0
6	NAG	D	503	14/15	0.58	0.36	-	62,88,125,127	0
6	NAG	D	501	14/15	0.85	0.16	-	32,54,65,66	0

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.