



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 11:55 am GMT

PDB ID : 3B6P  
Title : Structure of TREX1 in complex with a nucleotide and inhibitor ions (sodium and zinc)  
Authors : Brucet, M.; Querol-Audi, J.; Fita, I.; Celada, A.  
Deposited on : 2007-10-29  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	<div> <div>18%</div> <div> <div></div> <div>82%</div> <div>6%</div> <div>12%</div> </div> </div>
1	B	247	<div> <div>19%</div> <div> <div></div> <div>84%</div> <div>•</div> <div>12%</div> </div> </div>
1	C	247	<div> <div>19%</div> <div> <div></div> <div>85%</div> <div>•</div> <div>12%</div> </div> </div>
1	D	247	<div> <div>15%</div> <div> <div></div> <div>82%</div> <div>6%</div> <div>12%</div> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7299 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Three prime repair exonuclease 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	217	Total	C	N	O	S	0	1	0
			1684	1066	295	313	10			
1	B	217	Total	C	N	O	S	0	1	0
			1684	1066	295	313	10			
1	C	217	Total	C	N	O	S	0	1	0
			1684	1066	295	313	10			
1	D	217	Total	C	N	O	S	0	1	0
			1684	1066	295	313	10			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	EXPRESSION TAG	UNP Q91XB0
A	0	LYS	-	EXPRESSION TAG	UNP Q91XB0
A	1	HIS	-	EXPRESSION TAG	UNP Q91XB0
A	2	HIS	-	EXPRESSION TAG	UNP Q91XB0
A	3	HIS	-	EXPRESSION TAG	UNP Q91XB0
A	4	HIS	-	EXPRESSION TAG	UNP Q91XB0
A	5	HIS	-	EXPRESSION TAG	UNP Q91XB0
A	6	HIS	-	EXPRESSION TAG	UNP Q91XB0
A	7	PRO	-	EXPRESSION TAG	UNP Q91XB0
A	8	MET	-	EXPRESSION TAG	UNP Q91XB0
B	-1	MET	-	EXPRESSION TAG	UNP Q91XB0
B	0	LYS	-	EXPRESSION TAG	UNP Q91XB0
B	1	HIS	-	EXPRESSION TAG	UNP Q91XB0
B	2	HIS	-	EXPRESSION TAG	UNP Q91XB0
B	3	HIS	-	EXPRESSION TAG	UNP Q91XB0
B	4	HIS	-	EXPRESSION TAG	UNP Q91XB0
B	5	HIS	-	EXPRESSION TAG	UNP Q91XB0
B	6	HIS	-	EXPRESSION TAG	UNP Q91XB0
B	7	PRO	-	EXPRESSION TAG	UNP Q91XB0
B	8	MET	-	EXPRESSION TAG	UNP Q91XB0
C	-1	MET	-	EXPRESSION TAG	UNP Q91XB0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	LYS	-	EXPRESSION TAG	UNP Q91XB0
C	1	HIS	-	EXPRESSION TAG	UNP Q91XB0
C	2	HIS	-	EXPRESSION TAG	UNP Q91XB0
C	3	HIS	-	EXPRESSION TAG	UNP Q91XB0
C	4	HIS	-	EXPRESSION TAG	UNP Q91XB0
C	5	HIS	-	EXPRESSION TAG	UNP Q91XB0
C	6	HIS	-	EXPRESSION TAG	UNP Q91XB0
C	7	PRO	-	EXPRESSION TAG	UNP Q91XB0
C	8	MET	-	EXPRESSION TAG	UNP Q91XB0
D	-1	MET	-	EXPRESSION TAG	UNP Q91XB0
D	0	LYS	-	EXPRESSION TAG	UNP Q91XB0
D	1	HIS	-	EXPRESSION TAG	UNP Q91XB0
D	2	HIS	-	EXPRESSION TAG	UNP Q91XB0
D	3	HIS	-	EXPRESSION TAG	UNP Q91XB0
D	4	HIS	-	EXPRESSION TAG	UNP Q91XB0
D	5	HIS	-	EXPRESSION TAG	UNP Q91XB0
D	6	HIS	-	EXPRESSION TAG	UNP Q91XB0
D	7	PRO	-	EXPRESSION TAG	UNP Q91XB0
D	8	MET	-	EXPRESSION TAG	UNP Q91XB0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Zn 1 1	0	0
2	A	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Na 1 1	0	0
3	A	1	Total Na 1 1	0	0
3	D	1	Total Na 1 1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	Na	0	0
			1	1		

- Molecule 4 is THYMIDINE-5'-PHOSPHATE (three-letter code: TMP) (formula:  $C_{10}H_{15}N_2O_8P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			21	10	2	8	1		
4	B	1	Total	C	N	O	P	0	0
			21	10	2	8	1		
4	C	1	Total	C	N	O	P	0	0
			21	10	2	8	1		
4	D	1	Total	C	N	O	P	0	0
			21	10	2	8	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	125	Total	O	0	0
			125	125		
5	B	103	Total	O	0	0
			103	103		
5	C	105	Total	O	0	0
			105	105		
5	D	138	Total	O	0	0
			138	138		



- Molecule 1: Three prime repair exonuclease 1





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.77Å 81.50Å 92.54Å 90.00° 103.11° 90.00°	Depositor
Resolution (Å)	24.18 – 2.30 24.18 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (24.18-2.30) 98.2 (24.18-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.62 (at 2.19Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.246 , 0.285 0.264 , 0.303	Depositor DCC
$R_{free}$ test set	2155 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.2	Xtriage
Anisotropy	0.082	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 44.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	7299	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.06 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.1256e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TMP, NA, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.30	0/1724	0.49	0/2350
1	B	0.30	0/1724	0.48	0/2350
1	C	0.30	0/1724	0.49	0/2350
1	D	0.30	0/1724	0.50	0/2350
All	All	0.30	0/6896	0.49	0/9400

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1684	0	1686	6	0
1	B	1684	0	1686	4	0
1	C	1684	0	1686	1	0
1	D	1684	0	1686	7	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	21	0	13	0	0
4	B	21	0	13	0	0
4	C	21	0	13	0	0
4	D	21	0	13	0	0
5	A	125	0	0	0	0
5	B	103	0	0	1	0
5	C	105	0	0	0	0
5	D	138	0	0	0	0
All	All	7299	0	6796	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:ALA:HB3	1:A:65:ASP:HB2	1.91	0.51
1:B:37:LEU:HD21	1:B:205:LEU:HD22	1.94	0.49
1:D:38:ALA:HB3	1:D:65:ASP:HB2	1.96	0.48
1:B:121:LEU:HD11	1:B:147:LEU:HD22	1.95	0.47
1:D:121:LEU:HD11	1:D:147:LEU:HD22	1.97	0.46
1:A:121:LEU:HD11	1:A:147:LEU:HD22	1.99	0.45
1:C:121:LEU:HD11	1:C:147:LEU:HD22	1.98	0.45
1:A:43:ALA:HB1	1:A:61:PRO:HB3	2.00	0.43
1:B:73:PRO:HD2	5:B:1081:HOH:O	2.18	0.43
1:B:14:LEU:HD21	1:B:219:VAL:HG11	2.00	0.43
1:A:93:GLU:HB3	1:D:49:ILE:HD12	2.01	0.42
1:A:53:HIS:HA	1:A:54:PRO:HA	1.90	0.42
1:A:24:LEU:HB3	1:A:25:PRO:HD2	2.01	0.42
1:D:21:ALA:HB3	1:D:129:TYR:OH	2.20	0.41
1:D:103:ASN:HA	1:D:106:ILE:HD12	2.01	0.41
1:D:37:LEU:HD21	1:D:205:LEU:HD22	2.03	0.41
1:D:43:ALA:HB1	1:D:61:PRO:HB3	2.02	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/247 (87%)	210 (98%)	4 (2%)	0	100	100
1	B	214/247 (87%)	207 (97%)	7 (3%)	0	100	100
1	C	214/247 (87%)	207 (97%)	6 (3%)	1 (0%)	32	39
1	D	214/247 (87%)	210 (98%)	4 (2%)	0	100	100
All	All	856/988 (87%)	834 (97%)	21 (2%)	1 (0%)	55	67

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	50	SER

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/212 (88%)	184 (98%)	3 (2%)	68	82
1	B	187/212 (88%)	185 (99%)	2 (1%)	78	89
1	C	187/212 (88%)	183 (98%)	4 (2%)	59	76
1	D	187/212 (88%)	185 (99%)	2 (1%)	78	89
All	All	748/848 (88%)	737 (98%)	11 (2%)	70	83

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	133	LEU
1	A	148	ASP
1	A	205	LEU
1	B	133	LEU
1	B	153	VAL
1	C	64	VAL
1	C	133	LEU
1	C	153	VAL
1	C	205	LEU
1	D	133	LEU
1	D	153	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	GLN
1	A	125	ASN
1	A	217	GLN
1	B	125	ASN
1	B	189	GLN
1	B	217	GLN
1	C	46	ASN
1	C	125	ASN
1	C	189	GLN
1	C	217	GLN
1	D	125	ASN
1	D	217	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	TMP	A	1000	3,2	17,22,22	0.70	0	24,33,33	2.07	4 (16%)
4	TMP	B	1000	3,2	17,22,22	0.71	0	24,33,33	2.09	4 (16%)
4	TMP	C	1000	3,2	17,22,22	0.70	0	24,33,33	2.09	4 (16%)
4	TMP	D	1000	3,2	17,22,22	0.71	0	24,33,33	2.08	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TMP	A	1000	3,2	-	0/6/22/22	0/2/2/2
4	TMP	B	1000	3,2	-	0/6/22/22	0/2/2/2
4	TMP	C	1000	3,2	-	0/6/22/22	0/2/2/2
4	TMP	D	1000	3,2	-	0/6/22/22	0/2/2/2

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	1000	TMP	C5-C4-N3	-5.69	118.97	125.24
4	B	1000	TMP	C5-C4-N3	-5.63	119.03	125.24
4	A	1000	TMP	C5-C4-N3	-5.63	119.04	125.24
4	D	1000	TMP	C5-C4-N3	-5.57	119.10	125.24
4	B	1000	TMP	C2'-C1'-N1	-2.40	108.56	114.23
4	C	1000	TMP	C2'-C1'-N1	-2.33	108.74	114.23
4	D	1000	TMP	C2'-C1'-N1	-2.29	108.81	114.23
4	A	1000	TMP	C2'-C1'-N1	-2.21	109.00	114.23
4	C	1000	TMP	O4'-C1'-N1	2.53	112.04	107.78
4	B	1000	TMP	O4'-C1'-N1	2.54	112.05	107.78

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1000	TMP	O4'-C1'-N1	2.61	112.17	107.78
4	A	1000	TMP	O4'-C1'-N1	2.67	112.28	107.78
4	D	1000	TMP	C4-N3-C2	6.80	121.11	115.16
4	B	1000	TMP	C4-N3-C2	6.81	121.11	115.16
4	A	1000	TMP	C4-N3-C2	6.81	121.12	115.16
4	C	1000	TMP	C4-N3-C2	6.92	121.21	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	217/247 (87%)	1.26	44 (20%) <b>1</b> <b>1</b>	25, 28, 33, 39	0
1	B	217/247 (87%)	1.30	48 (22%) <b>1</b> <b>1</b>	25, 28, 34, 37	0
1	C	217/247 (87%)	1.26	47 (21%) <b>1</b> <b>1</b>	26, 28, 36, 39	0
1	D	217/247 (87%)	1.12	38 (17%) <b>2</b> <b>2</b>	25, 28, 34, 39	0
All	All	868/988 (87%)	1.24	177 (20%) <b>1</b> <b>1</b>	25, 28, 34, 39	0

All (177) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	49	ILE	10.5
1	B	165	ALA	9.0
1	C	48	SER	8.8
1	A	165	ALA	8.7
1	B	49	ILE	8.2
1	C	47	THR	7.1
1	D	165	ALA	7.0
1	B	48	SER	6.5
1	B	50	SER	6.4
1	B	46	ASN	6.4
1	B	148	ASP	6.1
1	C	50	SER	6.1
1	A	50	SER	5.9
1	D	164	GLN	5.9
1	C	53	HIS	5.8
1	A	164	GLN	5.8
1	D	10	HIS	5.4
1	A	10	HIS	5.3
1	B	9	GLY	5.3
1	A	49	ILE	5.2
1	D	51	GLN	5.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	163	GLU	5.1
1	A	59	ARG	5.1
1	B	164	GLN	4.9
1	B	234	GLY	4.9
1	B	47	THR	4.9
1	C	46	ASN	4.8
1	B	10	HIS	4.8
1	C	10	HIS	4.8
1	A	51	GLN	4.7
1	B	163	GLU	4.7
1	D	59	ARG	4.6
1	B	161	ALA	4.6
1	D	49	ILE	4.5
1	B	162	LEU	4.4
1	C	9	GLY	4.3
1	B	53	HIS	4.2
1	D	52	GLY	4.1
1	C	177	TYR	4.0
1	B	175	LYS	4.0
1	C	26	SER	3.9
1	A	122	VAL	3.9
1	C	51	GLN	3.9
1	A	26	SER	3.8
1	C	80	GLY	3.8
1	B	121	LEU	3.8
1	D	234	GLY	3.7
1	C	163	GLU	3.7
1	B	217	GLN	3.7
1	B	51	GLN	3.6
1	A	19	LEU	3.5
1	D	122	VAL	3.5
1	C	164	GLN	3.5
1	B	80	GLY	3.5
1	D	163	GLU	3.5
1	A	148	ASP	3.5
1	A	52	GLY	3.5
1	C	17	LEU	3.5
1	B	59	ARG	3.4
1	A	53	HIS	3.4
1	D	142	SER	3.4
1	D	148	ASP	3.4
1	C	128	ARG	3.3

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	44	LEU	3.3
1	A	42	ARG	3.3
1	A	175	LYS	3.3
1	C	42	ARG	3.2
1	D	53	HIS	3.2
1	C	121	LEU	3.2
1	A	16	PHE	3.2
1	C	99	ARG	3.1
1	A	234	GLY	3.1
1	A	17	LEU	3.1
1	B	142	SER	3.1
1	C	188	TRP	3.1
1	C	16	PHE	3.1
1	C	122	VAL	3.0
1	B	122	VAL	3.0
1	B	128	ARG	3.0
1	B	177	TYR	3.0
1	B	160	LYS	3.0
1	C	15	ILE	3.0
1	B	44	LEU	2.9
1	A	9	GLY	2.9
1	B	188	TRP	2.9
1	D	19	LEU	2.9
1	B	15	ILE	2.9
1	D	15	ILE	2.8
1	A	143	THR	2.8
1	B	17	LEU	2.8
1	C	148	ASP	2.8
1	D	34	LEU	2.8
1	A	142	SER	2.8
1	D	42	ARG	2.8
1	C	79	PRO	2.8
1	A	157	ALA	2.8
1	A	131	PHE	2.8
1	A	184	THR	2.8
1	C	230	LYS	2.8
1	D	50	SER	2.7
1	C	175	LYS	2.7
1	B	16	PHE	2.7
1	D	16	PHE	2.7
1	B	79	PRO	2.6
1	C	142	SER	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	175	LYS	2.6
1	B	67	LEU	2.6
1	C	36	LEU	2.6
1	A	15	ILE	2.6
1	D	9	GLY	2.6
1	B	230	LYS	2.6
1	A	123	ALA	2.6
1	C	157	ALA	2.6
1	B	19	LEU	2.5
1	B	69	LEU	2.5
1	A	80	GLY	2.5
1	C	143	THR	2.5
1	A	34	LEU	2.5
1	B	210	TRP	2.5
1	A	208	CYS	2.5
1	B	222	HIS	2.4
1	D	36	LEU	2.4
1	C	165	ALA	2.4
1	D	80	GLY	2.4
1	B	184	THR	2.4
1	D	26	SER	2.4
1	C	14	LEU	2.4
1	D	208	CYS	2.4
1	D	67	LEU	2.3
1	B	220	ASP	2.3
1	D	35	CYS	2.3
1	D	17	LEU	2.3
1	D	37	LEU	2.3
1	C	19	LEU	2.3
1	C	52	GLY	2.3
1	A	162	LEU	2.3
1	B	52	GLY	2.3
1	D	157	ALA	2.3
1	D	184	THR	2.3
1	D	77	CYS	2.3
1	A	140	ARG	2.3
1	B	83	GLU	2.2
1	A	217	GLN	2.2
1	C	28	ARG	2.2
1	C	220	ASP	2.2
1	A	218	TRP	2.2
1	A	204	LEU	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	108	LEU	2.2
1	B	118	PRO	2.2
1	D	39	VAL	2.2
1	C	59	ARG	2.2
1	A	35	CYS	2.2
1	C	67	LEU	2.1
1	C	109	ARG	2.1
1	C	45	GLU	2.1
1	A	117	GLN	2.1
1	C	69	LEU	2.1
1	D	144	PRO	2.1
1	A	77	CYS	2.1
1	B	119[A]	CYS	2.1
1	A	57	VAL	2.1
1	B	107	LEU	2.1
1	B	191	PRO	2.1
1	A	161	ALA	2.1
1	D	69	LEU	2.1
1	A	18	ASP	2.1
1	B	213	GLN	2.1
1	A	47	THR	2.1
1	A	153	VAL	2.1
1	C	176	SER	2.1
1	D	123	ALA	2.1
1	D	139	ALA	2.1
1	C	31	VAL	2.1
1	D	153	VAL	2.1
1	A	67	LEU	2.0
1	C	83	GLU	2.0
1	B	31	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	TMP	B	1000	21/21	0.89	0.17	-0.38	23,24,24,24	0
4	TMP	D	1000	21/21	0.90	0.15	-0.69	25,25,26,27	0
4	TMP	C	1000	21/21	0.90	0.16	-0.83	26,27,27,27	0
4	TMP	A	1000	21/21	0.89	0.15	-0.87	22,23,24,24	0
2	ZN	A	800	1/1	1.00	0.05	-2.21	28,28,28,28	0
2	ZN	B	800	1/1	0.99	0.04	-3.04	25,25,25,25	0
2	ZN	D	800	1/1	0.99	0.03	-3.14	28,28,28,28	0
2	ZN	C	800	1/1	0.98	0.03	-4.09	27,27,27,27	0
3	NA	C	801	1/1	0.74	0.20	-	31,31,31,31	0
3	NA	D	801	1/1	0.92	0.25	-	25,25,25,25	0
3	NA	A	801	1/1	0.82	0.24	-	22,22,22,22	0
3	NA	B	801	1/1	0.88	0.17	-	25,25,25,25	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.