



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 28, 2017 – 08:46 pm GMT

PDB ID : 1B7Y
Title : PHENYLALANYL TRNA SYNTHETASE COMPLEXED WITH PHENYL
ALANINYL-ADENYLATE
Authors : Reshetnikova, L.; Moor, N.; Lavrik, O.; Vassilyev, D.G.
Deposited on : 1999-01-26
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : recal29047
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recal29047


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3846 (2.50-2.50)
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)
RSRZ outliers	101464	3876 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	350	<div> <div>2%</div> <div>33% 34% 9% 24%</div> </div>
2	B	785	<div> <div>5%</div> <div>43% 43% 12%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MG	A	1001	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8439 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (PHENYLALANYL-TRNA SYNTHETASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	265	Total	C	N	O	S	0	0	0
			2112	1382	359	364	7			


- Molecule 2 is a protein called PROTEIN (PHENYLALANYL-TRNA SYNTHETASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	775	Total	C	N	O	S	0	0	0
			6054	3879	1078	1087	10			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is ADENOSINE-5'-[PHENYLALANINOL-PHOSPHATE] (three-letter code: FYA) (formula: C₁₉H₂₅N₆O₇P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			33	19	6	7	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	48	Total	O	0	0
			48	48		
5	B	191	Total	O	0	0
			191	191		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PROTEIN (PHENYLALANYL-TRNA SYNTHETASE)



• Molecule 2: PROTEIN (PHENYLALANYL-TRNA SYNTHETASE)



LEU	A717	G650	E581	D517	T417	I331
ARG	A713	F651	E582	F518	I427	A332
GLY	L652	T583	H584	E519	L428	C337
LEU	Y721	H656	G587	A521	K429	F338
ASP	E723	P657	L588	R522	R430	D339
THR	S724	E658	L589	R523	E438	R344
PRO		I659	F590	F524		K345
		A660	G591	R525	T441	E355
	L727	Q661	E592	D527	P446	A356
	F728	E662	G593	P528	P447	S357
	D729	L663	V594	P529	S448	H556
	L730	E664	G595	R530	H449	R359
	Y731	L665	L596	L531	R450	F360
	Q732	P666	P597	L532	L451	E361
	G733	P667	V598	L533	D452	V364
	P734	V668	A599	L534	D458	D365
	P735	H669	K600	N535	L459	P366
	L736	L670	E601	F536	V460	L367
	P737	F671	R602	L537	E461	Q368
	E738	E672	L603	A538	E462	Q369
	G739	L673	S604	P539	R465	V370
	H740	R674	L608	E540	I466	P371
	K741	L675	L609	X541	Q467	R374
	S742	P676	K610	A542	P473	L377
	L743		L613	A543	D484	S378
	A744	K680	L614	R545	R485	L380
	F745	P681	E614	T546	R486	Q381
	H746	L682	A615	H547	G487	G385
	L747	Q685	L616	L548	D488	A386
	R748		F617	F549	R489	R387
	F749	S688	A618	P550	A490	V388
	R750	R689	G621	G551	P491	A389
	H751		L622	V553	K494	E390
	P752		A623	R554	R497	L393
	K753		F624	V555	L498	E394
	R754	A692	Q629	L556	R499	S397
	L755	A693	A630	K557	E500	P398
	L756	F694	F631	E558	V501	K399
	R757	R695	P632		L505	P400
	D758	D696	L633	D562	G506	P401
	E759	L697	L634	D563	F507	E402
	E760	A698	H635	R564	Q508	A403
	V761	L699	P636	P565	E509	I404
	E762	V700	E566	E567	V510	E409
	E763	V701	G637	L570	T512	R413
	A764	P702	V638	F571	Y513	R414
	V765	A703	S639	E572	S514	L414
	S766	P704		V573	F515	L415
	R767	T705		G574	S516	G416
	V768	P706		R575		
	A769	T707		V576		
	E770	G708				
	A771	E709	V642			
	L772	V710	L643			
	R773	E711	V644			
	A774	A712	R645			
	R775	L713	G646			
	GLY	V714	E647			
	PHE	R715	E648			
	GLY	E716	V649			

4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	174.50Å 174.50Å 140.20Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.50 47.41 – 2.50	Depositor EDS
% Data completeness (in resolution range)	95.7 (50.00-2.50) 95.7 (47.41-2.50)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	15112.19 (at 2.51Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.230 , 0.267 0.229 , 0.264	Depositor DCC
R_{free} test set	4107 reflections (5.03%)	DCC
Wilson B-factor (Å ²)	43.4	Xtriage
Anisotropy	0.241	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 82.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	0.056 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8439	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, FYA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/2180	0.76	0/2957
2	B	0.53	0/6205	0.79	4/8436 (0.0%)
All	All	0.53	0/8385	0.78	4/11393 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	115	GLY	N-CA-C	-5.83	98.54	113.10
2	B	601	GLU	CB-CA-C	-5.59	99.22	110.40
2	B	128	TYR	N-CA-C	5.35	125.44	111.00
2	B	570	LEU	CA-CB-CG	5.28	127.44	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2112	0	2062	164	0
2	B	6054	0	6109	495	0
3	A	1	0	0	0	0
4	A	33	0	24	4	0
5	A	48	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	191	0	0	6	0
All	All	8439	0	8195	626	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 38.

The worst 5 of 626 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:262:GLU:HG2	5:A:1015:HOH:O	1.45	1.16
2:B:600:LYS:HG2	2:B:601:GLU:H	1.04	1.14
2:B:75:VAL:HG11	2:B:108:ILE:HG21	1.31	1.12
2:B:285:THR:HG21	2:B:291:ARG:HE	1.20	1.02
2:B:294:HIS:CD2	2:B:296:GLU:H	1.79	1.00

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/350 (75%)	244 (93%)	14 (5%)	5 (2%)	9	15
2	B	773/785 (98%)	695 (90%)	63 (8%)	15 (2%)	9	15
All	All	1036/1135 (91%)	939 (91%)	77 (7%)	20 (2%)	9	15

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	138	ASN
2	B	488	VAL
2	B	664	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	338	GLY
2	B	708	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	213/277 (77%)	176 (83%)	37 (17%)	2	4
2	B	623/630 (99%)	493 (79%)	130 (21%)	1	2
All	All	836/907 (92%)	669 (80%)	167 (20%)	1	2

5 of 167 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	243	ASN
2	B	361	GLU
2	B	716	GLU
2	B	283	LEU
2	B	307	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	54	HIS
2	B	212	HIS
2	B	656	HIS
2	B	109	GLN
2	B	178	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	FYA	A	1002	-	33,36,36	1.19	3 (9%)	31,52,52	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FYA	A	1002	-	-	0/16/36/36	0/4/4/4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1002	FYA	O3P-C	-4.32	1.27	1.44
4	A	1002	FYA	C8-N7	-2.00	1.30	1.34
4	A	1002	FYA	O4'-C1'	2.52	1.44	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1002	FYA	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	265/350 (75%)	0.07	8 (3%) 51 53	27, 54, 103, 129	0
2	B	775/785 (98%)	0.08	39 (5%) 30 31	21, 59, 108, 128	0
All	All	1040/1135 (91%)	0.08	47 (4%) 34 36	21, 58, 107, 129	0

The worst 5 of 47 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	149	TRP	5.6
2	B	718	ALA	5.0
1	A	350	LEU	4.9
2	B	753	LYS	4.8
2	B	739	GLY	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	MG	A	1001	1/1	0.84	0.26	9.14	42,42,42,42	0
4	FYA	A	1002	33/33	0.97	0.15	-0.01	40,65,73,76	0

6.5 Other polymers [i](#)

There are no such residues in this entry.