



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 15, 2017 – 04:48 AM EDT

PDB ID : 3B8M
Title : Structure of FepE- Bacterial Polysaccharide Co-polymerase
Authors : Tocilj, A.; Matte, A.; Cygler, M.
Deposited on : unknown
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2259 (2.70-2.70)
Clashscore	112137	2590 (2.70-2.70)
Ramachandran outliers	110173	2550 (2.70-2.70)
Sidechain outliers	110143	2550 (2.70-2.70)
RSRZ outliers	101464	2275 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	<div> <div>3%</div> <div>78%</div> <div>11%</div> <div>9%</div> </div>
1	B	280	<div> <div>2%</div> <div>75%</div> <div>15%</div> <div>9%</div> </div>
1	C	280	<div> <div>3%</div> <div>76%</div> <div>13%</div> <div>9%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6159 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ferric enterobactin (Enterochelin) transport.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	255	Total	C	N	O	S	0	0	0
			2048	1304	339	402	3			
1	B	255	Total	C	N	O	S	0	0	0
			2040	1299	337	401	3			
1	C	255	Total	C	N	O	S	0	0	0
			2043	1301	337	402	3			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	52	MET	-	EXPRESSION TAG	UNP Q8XBV8
A	53	GLY	-	EXPRESSION TAG	UNP Q8XBV8
A	54	SER	-	EXPRESSION TAG	UNP Q8XBV8
A	55	SER	-	EXPRESSION TAG	UNP Q8XBV8
A	56	HIS	-	EXPRESSION TAG	UNP Q8XBV8
A	57	HIS	-	EXPRESSION TAG	UNP Q8XBV8
A	58	HIS	-	EXPRESSION TAG	UNP Q8XBV8
A	59	HIS	-	EXPRESSION TAG	UNP Q8XBV8
A	60	HIS	-	EXPRESSION TAG	UNP Q8XBV8
A	61	HIS	-	EXPRESSION TAG	UNP Q8XBV8
A	62	GLY	-	EXPRESSION TAG	UNP Q8XBV8
A	63	SER	-	EXPRESSION TAG	UNP Q8XBV8
B	52	MET	-	EXPRESSION TAG	UNP Q8XBV8
B	53	GLY	-	EXPRESSION TAG	UNP Q8XBV8
B	54	SER	-	EXPRESSION TAG	UNP Q8XBV8
B	55	SER	-	EXPRESSION TAG	UNP Q8XBV8
B	56	HIS	-	EXPRESSION TAG	UNP Q8XBV8
B	57	HIS	-	EXPRESSION TAG	UNP Q8XBV8
B	58	HIS	-	EXPRESSION TAG	UNP Q8XBV8
B	59	HIS	-	EXPRESSION TAG	UNP Q8XBV8
B	60	HIS	-	EXPRESSION TAG	UNP Q8XBV8
B	61	HIS	-	EXPRESSION TAG	UNP Q8XBV8
B	62	GLY	-	EXPRESSION TAG	UNP Q8XBV8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	63	SER	-	EXPRESSION TAG	UNP Q8XBV8
C	52	MET	-	EXPRESSION TAG	UNP Q8XBV8
C	53	GLY	-	EXPRESSION TAG	UNP Q8XBV8
C	54	SER	-	EXPRESSION TAG	UNP Q8XBV8
C	55	SER	-	EXPRESSION TAG	UNP Q8XBV8
C	56	HIS	-	EXPRESSION TAG	UNP Q8XBV8
C	57	HIS	-	EXPRESSION TAG	UNP Q8XBV8
C	58	HIS	-	EXPRESSION TAG	UNP Q8XBV8
C	59	HIS	-	EXPRESSION TAG	UNP Q8XBV8
C	60	HIS	-	EXPRESSION TAG	UNP Q8XBV8
C	61	HIS	-	EXPRESSION TAG	UNP Q8XBV8
C	62	GLY	-	EXPRESSION TAG	UNP Q8XBV8
C	63	SER	-	EXPRESSION TAG	UNP Q8XBV8

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	13	Total O 13 13	0	0
2	B	7	Total O 7 7	0	0
2	C	8	Total O 8 8	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	139.73Å 139.73Å 276.65Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.13 – 2.70 46.11 – 2.70	Depositor EDS
% Data completeness (in resolution range)	92.9 (46.13-2.70) 92.9 (46.11-2.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.36 (at 2.69Å)	Xtriage
Refinement program	REFMAC refmac_5.2.0019	Depositor
R, R_{free}	0.226 , 0.269 0.218 , 0.257	Depositor DCC
R_{free} test set	2072 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	46.0	Xtriage
Anisotropy	0.052	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 38.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6159	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	1/2081 (0.0%)	0.69	0/2814
1	B	0.57	0/2073	0.65	0/2805
1	C	0.60	0/2076	0.67	0/2810
All	All	0.60	1/6230 (0.0%)	0.67	0/8429

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	297	GLU	CG-CD	5.32	1.59	1.51

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2048	0	2064	21	0
1	B	2040	0	2047	18	0
1	C	2043	0	2051	22	0
2	A	13	0	0	0	0
2	B	7	0	0	0	0
2	C	8	0	0	0	0
All	All	6159	0	6162	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:207:VAL:HG12	1:A:313:ILE:HD13	1.41	1.01
1:B:75:GLU:H	1:B:78:GLN:NE2	1.80	0.79
1:B:180:THR:HB	1:B:183:GLU:H	1.56	0.71
1:A:170:THR:O	1:A:171:SER:HB3	1.89	0.70
1:A:167:SER:O	1:A:167:SER:OG	2.08	0.69
1:B:225:ASP:OD2	1:B:301:ARG:NH2	2.34	0.59
1:A:207:VAL:HG12	1:A:313:ILE:CD1	2.26	0.58
1:A:252:ILE:HG22	1:A:274:GLY:HA2	1.86	0.58
1:A:170:THR:O	1:A:171:SER:CB	2.49	0.57
1:A:221:LYS:HE2	1:A:224:GLN:OE1	2.04	0.57
1:A:170:THR:HG22	1:A:171:SER:N	2.20	0.56
1:A:229:MET:HE3	1:A:295:ASN:HD22	1.71	0.55
1:A:75:GLU:H	1:A:78:GLN:HE21	1.54	0.55
1:A:289:THR:HG22	1:A:290:ASP:N	2.20	0.55
1:C:181:SER:HB2	1:C:330:PRO:HG2	1.89	0.55
1:B:221:LYS:NZ	1:B:224:GLN:OE1	2.40	0.54
1:B:106:LEU:HD23	1:B:199:VAL:HG13	1.89	0.54
1:C:72:THR:HG22	1:C:73:PRO:O	2.06	0.54
1:C:162:LYS:O	1:C:164:ASP:N	2.39	0.54
1:C:66:THR:HG1	1:C:177:THR:HG1	1.52	0.54
1:A:99:ASP:HB3	1:A:102:GLU:HB2	1.89	0.53
1:B:75:GLU:OE2	1:C:109:LYS:NZ	2.36	0.53
1:B:181:SER:HB2	1:B:330:PRO:HG2	1.92	0.52
1:C:185:GLN:HG3	1:C:328:SER:O	2.11	0.51
1:B:72:THR:HG22	1:B:322:LYS:O	2.11	0.50
1:C:162:LYS:C	1:C:164:ASP:H	2.14	0.50
1:C:326:SER:HB2	1:C:327:PRO:HD2	1.93	0.49
1:B:252:ILE:HD12	1:B:271:ILE:HD12	1.93	0.49
1:B:178:ALA:HB1	1:B:179:PRO:CD	2.44	0.48
1:C:297:GLU:O	1:C:301:ARG:HG3	2.14	0.48
1:C:128:MET:CE	1:C:128:MET:HA	2.44	0.47
1:C:82:LEU:HD13	1:C:316:VAL:HG11	1.97	0.47
1:C:138:GLU:HB2	1:C:139:LEU:HD22	1.97	0.46
1:B:188:LEU:HD23	1:B:327:PRO:HB3	1.96	0.46
1:C:225:ASP:OD2	1:C:301:ARG:NH1	2.49	0.46
1:A:130:GLN:N	1:A:130:GLN:HE21	2.13	0.46
1:A:225:ASP:OD2	1:A:301:ARG:NH2	2.49	0.46
1:C:185:GLN:HE21	1:C:185:GLN:HB3	1.62	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:164:ASP:O	1:C:166:PRO:HD3	2.15	0.46
1:C:81:GLU:HB3	1:C:316:VAL:HG13	1.97	0.46
1:C:99:ASP:HB3	1:C:102:GLU:HB2	1.98	0.46
1:B:72:THR:CG2	1:B:322:LYS:HB3	2.46	0.45
1:A:68:ALA:O	1:A:327:PRO:HA	2.17	0.45
1:A:170:THR:HG22	1:A:171:SER:H	1.82	0.44
1:C:168:LEU:HD12	1:C:168:LEU:HA	1.84	0.44
1:A:229:MET:HB3	1:A:229:MET:HE2	1.83	0.44
1:B:124:SER:HB3	1:B:127:VAL:HB	2.00	0.44
1:A:298:LEU:HA	1:A:298:LEU:HD12	1.80	0.44
1:C:128:MET:HA	1:C:128:MET:HE2	1.99	0.44
1:B:176:PHE:CE2	1:B:187:VAL:HG11	2.53	0.43
1:B:65:TRP:HD1	1:B:179:PRO:O	2.01	0.42
1:A:160:SER:O	1:A:161:LYS:HB3	2.18	0.42
1:B:255:PRO:HG2	1:C:252:ILE:HD11	2.02	0.41
1:B:294:LEU:HA	1:B:294:LEU:HD23	1.92	0.41
1:C:240:LEU:HA	1:C:240:LEU:HD12	1.84	0.41
1:C:104:PHE:O	1:C:107:PHE:HB3	2.21	0.41
1:A:128:MET:CE	1:A:128:MET:HA	2.51	0.41
1:A:66:THR:HA	1:A:177:THR:HA	2.03	0.41
1:B:252:ILE:HG22	1:B:274:GLY:HA2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/280 (89%)	237 (95%)	10 (4%)	2 (1%)	22	49
1	B	249/280 (89%)	240 (96%)	7 (3%)	2 (1%)	22	49
1	C	249/280 (89%)	241 (97%)	7 (3%)	1 (0%)	38	66
All	All	747/840 (89%)	718 (96%)	24 (3%)	5 (1%)	25	53

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	163	LYS
1	C	163	LYS
1	A	171	SER
1	A	162	LYS
1	B	179	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/251 (91%)	217 (95%)	12 (5%)	27	55
1	B	227/251 (90%)	211 (93%)	16 (7%)	18	40
1	C	228/251 (91%)	213 (93%)	15 (7%)	19	43
All	All	684/753 (91%)	641 (94%)	43 (6%)	21	46

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	79	TRP
1	A	82	LEU
1	A	97	LYS
1	A	130	GLN
1	A	141	LEU
1	A	161	LYS
1	A	171	SER
1	A	198	LEU
1	A	221	LYS
1	A	230	LYS
1	A	298	LEU
1	A	317	ASN
1	B	79	TRP
1	B	81	GLU
1	B	82	LEU
1	B	85	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	92	LEU
1	B	95	ASP
1	B	96	ILE
1	B	119	GLU
1	B	140	ASP
1	B	141	LEU
1	B	180	THR
1	B	198	LEU
1	B	206	ASN
1	B	241	ASN
1	B	289	THR
1	B	308	LEU
1	C	72	THR
1	C	79	TRP
1	C	82	LEU
1	C	88	LYS
1	C	129	ASP
1	C	138	GLU
1	C	160	SER
1	C	165	GLU
1	C	180	THR
1	C	181	SER
1	C	185	GLN
1	C	232	GLN
1	C	256	VAL
1	C	272	SER
1	C	289	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	78	GLN
1	A	105	ASN
1	A	130	GLN
1	A	206	ASN
1	B	78	GLN
1	B	105	ASN
1	C	78	GLN
1	C	105	ASN
1	C	185	GLN
1	C	206	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	255/280 (91%)	-0.01	9 (3%) 44 44	22, 41, 92, 124	0
1	B	255/280 (91%)	-0.08	6 (2%) 59 60	23, 42, 91, 122	0
1	C	255/280 (91%)	0.03	9 (3%) 44 44	24, 41, 95, 124	0
All	All	765/840 (91%)	-0.02	24 (3%) 49 49	22, 41, 92, 124	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	164	ASP	5.5
1	B	164	ASP	5.2
1	B	162	LYS	5.1
1	C	129	ASP	5.0
1	A	161	LYS	4.6
1	B	166	PRO	4.4
1	A	162	LYS	4.3
1	C	164	ASP	4.3
1	C	160	SER	4.1
1	A	163	LYS	3.9
1	A	166	PRO	3.8
1	C	257	TYR	3.2
1	C	165	GLU	3.1
1	C	166	PRO	3.1
1	B	140	ASP	2.9
1	B	160	SER	2.7
1	C	126	TYR	2.6
1	C	161	LYS	2.6
1	A	160	SER	2.6
1	B	163	LYS	2.5
1	A	167	SER	2.4
1	C	139	LEU	2.4
1	A	128	MET	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	159	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.