



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 12:46 pm GMT

PDB ID : 1B9J
Title : OLIGO-PEPTIDE BINDING PROTEIN (OPPA) COMPLEXED WITH KLK
Authors : Tame, J.R.H.; Wilkinson, A.J.
Deposited on : 1999-02-11
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	517	<div> <div>4%</div> <div> <div></div> <div>87%</div> <div>11%</div> </div> </div>
2	B	3	<div> <div>67%</div> <div>33%</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4698 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PROTEIN (OLIGO-PEPTIDE BINDING PROTEIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	517	Total	C	N	O	S	0	2	0
			4170	2670	700	795	5			

- Molecule 2 is a protein called PROTEIN (LYS-LEU-LYS).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	3	Total	C	N	O	0	0	0
			27	18	5	4			

- Molecule 3 is URANYL (VI) ION (three-letter code: IUM) (formula: O₂U).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	U	0	0
			1	1		
3	A	1	Total	U	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	U 1	0	0
3	A	1	Total 1	U 1	0	0
3	A	1	Total 1	U 1	0	0
3	A	1	Total 1	U 1	0	0
3	A	1	Total 1	U 1	0	0
3	A	1	Total 1	U 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	485	Total 491	O 491	0	6
4	B	2	Total 2	O 2	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PROTEIN (OLIGO-PEPTIDE BINDING PROTEIN)



• Molecule 2: PROTEIN (LYS-LEU-LYS)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	110.05Å 75.58Å 70.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 1.80 14.98 – 1.80	Depositor EDS
% Data completeness (in resolution range)	90.9 (15.00-1.80) 90.9 (14.98-1.80)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.03 (at 1.79Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.179 , 0.219 0.169 , 0.208	Depositor DCC
R_{free} test set	2484 reflections (5.22%)	DCC
Wilson B-factor (Å ²)	19.4	Xtriage
Anisotropy	0.197	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 62.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4698	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: IUM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/4290	1.25	30/5850 (0.5%)
2	B	0.59	0/26	1.09	0/30
All	All	0.55	0/4316	1.25	30/5880 (0.5%)

There are no bond length outliers.

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	413	ARG	NE-CZ-NH2	-14.29	113.16	120.30
1	A	201	ARG	NE-CZ-NH1	12.94	126.77	120.30
1	A	413	ARG	NE-CZ-NH1	9.73	125.16	120.30
1	A	462	ARG	NE-CZ-NH1	8.07	124.33	120.30
1	A	410	ASP	CB-CG-OD2	-7.89	111.19	118.30
1	A	235	ARG	NE-CZ-NH2	-7.61	116.49	120.30
1	A	133	ASP	CB-CG-OD2	-6.89	112.10	118.30
1	A	201	ARG	CD-NE-CZ	6.67	132.94	123.60
1	A	485	TYR	CB-CG-CD2	-6.59	117.05	121.00
1	A	466	TYR	CB-CG-CD1	6.42	124.85	121.00
1	A	273	TYR	CB-CG-CD2	-6.41	117.15	121.00
1	A	84	ASP	CB-CG-OD1	6.32	123.99	118.30
1	A	17	ARG	NE-CZ-NH2	-6.30	117.15	120.30
1	A	466	TYR	CB-CG-CD2	-6.23	117.26	121.00
1	A	394	ASN	CB-CG-OD1	6.16	133.93	121.60
1	A	485	TYR	CB-CG-CD1	6.11	124.67	121.00
1	A	410	ASP	CB-CG-OD1	5.89	123.60	118.30
1	A	367	THR	CA-CB-CG2	-5.86	104.20	112.40
1	A	365	TYR	CB-CG-CD2	-5.69	117.59	121.00
1	A	42	ASP	CB-CG-OD1	5.61	123.35	118.30
1	A	404	ARG	NE-CZ-NH1	-5.49	117.55	120.30
1	A	102	ASP	CB-CG-OD2	-5.46	113.39	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	245	TYR	CB-CG-CD1	-5.35	117.79	121.00
1	A	458	ASP	CA-CB-CG	5.32	125.11	113.40
1	A	407	GLY	N-CA-C	5.21	126.12	113.10
1	A	269	TYR	CB-CG-CD2	-5.19	117.89	121.00
1	A	485	TYR	CA-CB-CG	5.18	123.25	113.40
1	A	447	ASP	CB-CG-OD1	5.04	122.83	118.30
1	A	177	ASP	CB-CG-OD1	5.03	122.83	118.30
1	A	223	TYR	CB-CG-CD2	-5.03	117.98	121.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4170	0	4084	24	1
2	B	27	0	39	1	0
3	A	8	0	0	0	1
4	A	491	0	0	1	0
4	B	2	0	0	0	0
All	All	4698	0	4123	24	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:301:ILE:HA	1:A:305:LYS:HG3	1.55	0.86
1:A:229:GLU:HB3	1:A:249:PRO:HD3	1.79	0.64
1:A:418:ALA:HB3	1:A:504:LEU:HD22	1.84	0.59
1:A:460:THR:O	1:A:464:GLU:HG3	2.04	0.57
1:A:63:LYS:NZ	1:A:63:LYS:HB3	2.24	0.53
1:A:43:LEU:HD21	1:A:204:LEU:HD22	1.91	0.52
1:A:453:THR:HG21	1:A:466:TYR:CE1	2.46	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:230:VAL:HG13	1:A:252:LEU:HD11	1.93	0.50
1:A:253:PHE:CD2	1:A:309:GLN:HG2	2.47	0.49
1:A:43:LEU:O	1:A:187:THR:HB	2.13	0.49
1:A:115:TYR:CE1	1:A:428:ASN:HB3	2.47	0.48
1:A:57[B]:SER:HB2	1:A:58:PRO:CD	2.44	0.48
1:A:118:ILE:HD12	1:A:118:ILE:N	2.30	0.47
1:A:382:TRP:HB3	1:A:388:VAL:HG22	1.96	0.47
1:A:118:ILE:HD12	1:A:118:ILE:H	1.81	0.45
1:A:57[B]:SER:HB2	1:A:58:PRO:HD2	1.98	0.45
1:A:215:LYS:HD2	4:A:790:HOH:O	2.19	0.43
1:A:57[A]:SER:HB3	1:A:58:PRO:CD	2.49	0.43
1:A:265:ARG:O	1:A:488:ALA:HA	2.20	0.42
1:A:366:ASN:HA	1:A:395:GLN:O	2.19	0.42
1:A:119:ALA:O	1:A:120:ASN:HB2	2.20	0.41
1:A:401:LEU:HD22	2:B:2:LEU:HD22	2.01	0.41
1:A:489:ARG:NH1	1:A:498:TYR:OH	2.47	0.41
1:A:3:VAL:HA	1:A:4:PRO:HD3	1.90	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:394:ASN:OD1	3:A:521:IUM:U[3_555]	1.96	0.24

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	517/517 (100%)	498 (96%)	18 (4%)	1 (0%)	51	35
2	B	1/3 (33%)	1 (100%)	0	0	100	100
All	All	518/520 (100%)	499 (96%)	18 (4%)	1 (0%)	51	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	225	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	457/455 (100%)	449 (98%)	8 (2%)	64	53
2	B	3/3 (100%)	3 (100%)	0	100	100
All	All	460/458 (100%)	452 (98%)	8 (2%)	66	55

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	LYS
1	A	269	TYR
1	A	306	VAL
1	A	334	LYS
1	A	338	GLN
1	A	413	ARG
1	A	458	ASP
1	A	475	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	195	ASN
1	A	199	ASN
1	A	279	ASN
1	A	280	GLN
1	A	304	ASN
1	A	406	GLN
1	A	517	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are modelled with single atom - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	517/517 (100%)	-0.21	19 (3%) 42 37	12, 20, 39, 61	0
2	B	3/3 (100%)	-0.80	0 100 100	14, 14, 16, 18	0
All	All	520/520 (100%)	-0.21	19 (3%) 42 37	12, 20, 39, 61	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	408	THR	6.2
1	A	252	LEU	5.1
1	A	308	ASN	5.0
1	A	305	LYS	4.1
1	A	356	ASP	3.9
1	A	5	ALA	3.5
1	A	249	PRO	3.4
1	A	310	GLY	3.0
1	A	357	LYS	2.9
1	A	253	PHE	2.8
1	A	258	LYS	2.7
1	A	177	ASP	2.5
1	A	306	VAL	2.4
1	A	300	ASP	2.4
1	A	384	LYS	2.4
1	A	407	GLY	2.4
1	A	262	ASN	2.3
1	A	6	GLY	2.3
1	A	254	GLN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	IUM	A	525	1/3	0.99	0.03	-	23,23,23,23	0
3	IUM	A	526	1/3	0.98	0.04	-	27,27,27,27	1
3	IUM	A	528	1/3	0.97	0.04	-	35,35,35,35	1
3	IUM	A	523	1/3	0.97	0.04	-	27,27,27,27	0
3	IUM	A	522	1/3	0.99	0.04	-	23,23,23,23	0
3	IUM	A	527	1/3	0.99	0.06	-	21,21,21,21	1
3	IUM	A	524	1/3	1.00	0.17	-	42,42,42,42	0
3	IUM	A	521	1/3	1.00	0.03	-	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.