



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 20, 2017 – 07:53 PM EST

PDB ID : 2BBV
Title : THE REFINED THREE-DIMENSIONAL STRUCTURE OF AN INSECT VIRUS AT 2.8 ANGSTROMS RESOLUTION
Authors : Wery, J.-P.; Reddy, V.S.; Hosur, M.V.; Johnson, J.E.
Deposited on : unknown
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| Clashscore | 112137 | 3033 (2.80-2.80) |
| Ramachandran outliers | 110173 | 2983 (2.80-2.80) |
| Sidechain outliers | 110143 | 2985 (2.80-2.80) |
| RNA backbone | 2435 | 1007 (3.10-2.50) |


The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|---|
| 1 | N | 10 | <div> <div>50%</div> <div>40%</div> <div>10%</div> </div> |
| 2 | A | 363 | <div> <div>39%</div> <div>34%</div> <div>11%</div> <div>•</div> <div>15%</div> </div> |
| 2 | B | 363 | <div> <div>40%</div> <div>34%</div> <div>9%</div> <div>•</div> <div>15%</div> </div> |
| 2 | C | 363 | <div> <div>36%</div> <div>38%</div> <div>13%</div> <div>•</div> <div>12%</div> </div> |
| 3 | D | 44 | <div> <div>7%</div> <div>16%</div> <div>9%</div> <div>5%</div> <div>64%</div> </div> |
| 3 | E | 44 | <div> <div>11%</div> <div>16%</div> <div>5%</div> <div>5%</div> <div>64%</div> </div> |

Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|--|
| 3 | F | 44 |  A horizontal bar chart showing the quality of chain F. The bar is divided into four segments: green (11%), yellow (14%), orange (9%), and red (64%). The segments are labeled with their respective percentages. The total length of the bar is 100%. |

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7817 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (5'-R(*UP*CP*UP*UP*AP*UP*AP*UP*CP*U)-3').

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|----|----|----|---|---------|---------|-------|
| 1 | N | 10 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 201 | 92 | 28 | 72 | 9 | | | |

- Molecule 2 is a protein called PROTEIN (BLACK BEETLE VIRUS CAPSID PROTEIN).

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|---------|-------|
| 2 | A | 308 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 2311 | 1478 | 385 | 438 | 10 | | | |
| 2 | B | 308 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 2311 | 1478 | 385 | 438 | 10 | | | |
| 2 | C | 321 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 2406 | 1536 | 402 | 458 | 10 | | | |

- Molecule 3 is a protein called PROTEIN (BLACK BEETLE VIRUS CAPSID PROTEIN).

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|----|----|----|---|---------|---------|-------|
| 3 | D | 16 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 125 | 80 | 22 | 22 | 1 | | | |
| 3 | E | 16 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 125 | 80 | 22 | 22 | 1 | | | |
| 3 | F | 16 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 125 | 80 | 22 | 22 | 1 | | | |

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 4 | B | 1 | Total | Ca | 0 | 0 |
| | | | 1 | 1 | | |
| 4 | A | 2 | Total | Ca | 0 | 0 |
| | | | 2 | 2 | | |


Continued on next page...

Continued from previous page...

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 4 | C | 2 | Total | Ca | 0 | 0 |
| | | | 2 | 2 | | |


- Molecule 5 is water.

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 5 | A | 68 | Total | O | 0 | 0 |
| | | | 68 | 68 | | |
| 5 | D | 1 | Total | O | 0 | 0 |
| | | | 1 | 1 | | |
| 5 | B | 70 | Total | O | 0 | 0 |
| | | | 70 | 70 | | |
| 5 | C | 65 | Total | O | 0 | 0 |
| | | | 65 | 65 | | |
| 5 | F | 4 | Total | O | 0 | 0 |
| | | | 4 | 4 | | |




• Molecule 2: PROTEIN (BLACK BEETLE VIRUS CAPSID PROTEIN)

Chain C: 36% 38% 13% 12%




• Molecule 3: PROTEIN (BLACK BEETLE VIRUS CAPSID PROTEIN)

Chain D: 7% 16% 9% 5% 64%




• Molecule 3: PROTEIN (BLACK BEETLE VIRUS CAPSID PROTEIN)

Chain E: 11% 16% 5% 5% 64%



• Molecule 3: PROTEIN (BLACK BEETLE VIRUS CAPSID PROTEIN)

Chain F: 11% 14% 9% 64%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

| Property | Value | Source |
|--|--|-----------|
| Space group | P 4 ₂ 3 ₂ | Depositor |
| Cell constants a, b, c, α , β , γ | 362.00 Å 362.00 Å 362.00 Å 90.00° 90.00° 90.00° | Depositor |
| Resolution (Å) | 6.00 – 2.80 | Depositor |
| % Data completeness (in resolution range) | 67.5 (6.00-2.80) | Depositor |
| R_{merge} | (Not available) | Depositor |
| R_{sym} | (Not available) | Depositor |
| Refinement program | X-PLOR, PROLSQ | Depositor |
| R, R_{free} | 0.221 , (Not available) | Depositor |
| Estimated twinning fraction | No twinning to report. | Xtriage |
| Total number of atoms | 7817 | wwPDB-VP |
| Average B, all atoms (Å ²) | 12.0 | wwPDB-VP |

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|---------------|-------------|-----------------|
| | | RMSZ | # $ Z > 5$ | RMSZ | # $ Z > 5$ |
| 1 | N | 1.31 | 1/222 (0.5%) | 1.59 | 3/342 (0.9%) |
| 2 | A | 0.89 | 0/2372 | 1.41 | 15/3249 (0.5%) |
| 2 | B | 0.88 | 0/2372 | 1.43 | 15/3249 (0.5%) |
| 2 | C | 0.88 | 1/2469 (0.0%) | 1.41 | 22/3384 (0.7%) |
| 3 | D | 1.01 | 0/126 | 1.57 | 5/167 (3.0%) |
| 3 | E | 1.02 | 0/126 | 1.43 | 2/167 (1.2%) |
| 3 | F | 1.48 | 1/126 (0.8%) | 1.75 | 3/167 (1.8%) |
| All | All | 0.92 | 3/7813 (0.0%) | 1.43 | 65/10725 (0.6%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 2 | B | 0 | 1 |

All (3) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|--------|-------------|----------|
| 3 | F | 364 | ALA | CA-CB | 12.65 | 1.79 | 1.52 |
| 1 | N | 7 | A | N9-C4 | -10.46 | 1.31 | 1.37 |
| 2 | C | 363 | ASN | CA-C | -8.50 | 1.30 | 1.52 |

The worst 5 of 65 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-----------|--------|-------------|----------|
| 3 | F | 364 | ALA | N-CA-CB | -13.14 | 91.71 | 110.10 |
| 2 | A | 352 | ARG | NE-CZ-NH1 | 11.89 | 126.24 | 120.30 |
| 2 | C | 348 | ARG | NE-CZ-NH2 | 10.31 | 125.45 | 120.30 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-----------|------|-------------|----------|
| 2 | B | 87 | ARG | NE-CZ-NH1 | 9.61 | 125.11 | 120.30 |
| 2 | C | 55 | ALA | O-C-N | 8.82 | 136.81 | 122.70 |

There are no chirality outliers.

All (1) planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|-----------|
| 2 | B | 58 | ARG | Sidechain |

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | N | 201 | 0 | 106 | 34 | 0 |
| 2 | A | 2311 | 0 | 2262 | 209 | 0 |
| 2 | B | 2311 | 0 | 2262 | 190 | 0 |
| 2 | C | 2406 | 0 | 2357 | 242 | 0 |
| 3 | D | 125 | 0 | 135 | 18 | 0 |
| 3 | E | 125 | 0 | 135 | 22 | 0 |
| 3 | F | 125 | 0 | 135 | 12 | 0 |
| 4 | A | 2 | 0 | 0 | 0 | 0 |
| 4 | B | 1 | 0 | 0 | 0 | 0 |
| 4 | C | 2 | 0 | 0 | 0 | 0 |
| 5 | A | 68 | 0 | 0 | 7 | 0 |
| 5 | B | 70 | 0 | 0 | 12 | 0 |
| 5 | C | 65 | 0 | 0 | 13 | 0 |
| 5 | D | 1 | 0 | 0 | 0 | 0 |
| 5 | F | 4 | 0 | 0 | 0 | 0 |
| All | All | 7817 | 0 | 7392 | 674 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 45.

The worst 5 of 674 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|------------------|--------------------------|-------------------|
| 3:F:364:ALA:CA | 3:F:364:ALA:CB | 1.79 | 1.58 |
| 2:B:207:THR:CG2 | 2:B:208:PRO:HD3 | 1.36 | 1.52 |
| 2:C:21:GLN:HG3 | 2:C:181:LEU:CD1 | 1.40 | 1.48 |
| 2:C:21:GLN:CG | 2:C:181:LEU:HD11 | 1.54 | 1.36 |
| 2:C:264:PRO:O | 2:C:267:VAL:HG13 | 1.22 | 1.32 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|----------------|-----------|---------|----------|-------------|-----|
| 2 | A | 306/363 (84%) | 280 (92%) | 23 (8%) | 3 (1%) | 18 | 50 |
| 2 | B | 306/363 (84%) | 278 (91%) | 25 (8%) | 3 (1%) | 18 | 50 |
| 2 | C | 317/363 (87%) | 296 (93%) | 18 (6%) | 3 (1%) | 20 | 52 |
| 3 | D | 14/44 (32%) | 12 (86%) | 2 (14%) | 0 | 100 | 100 |
| 3 | E | 14/44 (32%) | 13 (93%) | 1 (7%) | 0 | 100 | 100 |
| 3 | F | 14/44 (32%) | 13 (93%) | 0 | 1 (7%) | 1 | 3 |
| All | All | 971/1221 (80%) | 892 (92%) | 69 (7%) | 10 (1%) | 18 | 50 |

5 of 10 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2 | A | 57 | THR |
| 2 | A | 209 | ALA |
| 2 | B | 81 | GLY |
| 2 | B | 206 | THR |
| 2 | C | 236 | ILE |

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|---------------|-----------|-----------|-------------|---|
| 2 | A | 251/301 (83%) | 196 (78%) | 55 (22%) | 1 | 3 |
| 2 | B | 251/301 (83%) | 202 (80%) | 49 (20%) | 1 | 4 |
| 2 | C | 263/301 (87%) | 202 (77%) | 61 (23%) | 1 | 2 |
| 3 | D | 14/32 (44%) | 5 (36%) | 9 (64%) | 0 | 0 |
| 3 | E | 14/32 (44%) | 9 (64%) | 5 (36%) | 0 | 0 |
| 3 | F | 14/32 (44%) | 10 (71%) | 4 (29%) | 0 | 1 |
| All | All | 807/999 (81%) | 624 (77%) | 183 (23%) | 1 | 3 |

5 of 183 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2 | B | 181 | LEU |
| 2 | B | 310 | SER |
| 2 | C | 319 | LEU |
| 2 | B | 195 | VAL |
| 2 | B | 220 | LEU |

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2 | B | 242 | GLN |
| 2 | B | 326 | ASN |
| 2 | C | 331 | GLN |
| 2 | B | 293 | ASN |
| 2 | A | 246 | ASN |

5.3.3 RNA ⓘ

| Mol | Chain | Analysed | Backbone Outliers | Pucker Outliers |
|-----|-------|------------|-------------------|-----------------|
| 1 | N | 9/10 (90%) | 5 (55%) | 0 |

All (5) RNA backbone outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | N | 2 | C |
| 1 | N | 3 | U |
| 1 | N | 4 | U |
| 1 | N | 5 | A |
| 1 | N | 7 | A |

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.