



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 16, 2017 – 08:40 PM EDT

PDB ID : 3BC9  
Title : Alpha-amylase B in complex with acarbose  
Authors : Tan, T.-C.; Mijts, B.N.; Swaminathan, K.; Patel, B.K.C.; Divne, C.  
Deposited on : unknown  
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1024 (1.38-1.34)
Clashscore	112137	1063 (1.38-1.34)
Ramachandran outliers	110173	1048 (1.38-1.34)
Sidechain outliers	110143	1048 (1.38-1.34)
RSRZ outliers	101464	1025 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	599	<div> <div>5%</div> <div>89%</div> <div>8%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GLC	A	1001	-	-	-	X
3	GLC	A	1002	X	-	-	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GLC	A	802	-	-	-	X
4	ACI	A	803	-	-	-	X
6	BGC	A	809	-	-	-	X
7	ACR	A	901	X	-	-	-
8	CA	A	705	-	-	-	X

## 2 Entry composition [i](#)


There are 10 unique types of molecules in this entry. The entry contains 5566 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha amylase, catalytic region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	585	4822	3071	780	959	12	0	9	0

- Molecule 2 is 6-DEOXY-ALPHA-D-GLUCOSE (three-letter code: G6D) (formula:  $C_6H_{12}O_5$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	10	6	4	0	0

- Molecule 3 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula:  $C_6H_{12}O_6$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			12	6	6		
3	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 4 is 6-AMINO-4-HYDROXYMETHYL-CYCLOHEX-4-ENE-1,2,3-TRIOL (three-letter code: ACI) (formula: C<sub>7</sub>H<sub>13</sub>NO<sub>4</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			12	7	1	4		
4	A	1	Total	C	N	O	0	0
			12	7	1	4		

- Molecule 5 is 4,6-DIDEOXYGLUCOSE (three-letter code: GLD) (formula:  $C_6H_{12}O_4$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			9	6	3		
5	A	1	Total	C	O	0	0
			9	6	3		

- Molecule 6 is BETA-D-GLUCOSE (three-letter code: BGC) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 7 is ALPHA-ACARBOSE (three-letter code: ACR) (formula:  $C_{25}H_{43}NO_{18}$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			44	25	1	18		

- Molecule 8 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	4	Total	Ca	0	0
			4	4		

- Molecule 9 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1	Total	Na	0	0
			1	1		

- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	574	Total	O	0	0
			574	574		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha amylase, catalytic region



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	227.85Å 77.24Å 50.38Å 90.00° 98.63° 90.00°	Depositor
Resolution (Å)	30.00 – 1.35 29.71 – 1.35	Depositor EDS
% Data completeness (in resolution range)	99.2 (30.00-1.35) 99.2 (29.71-1.35)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.96 (at 1.35Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.151 , 0.178 0.160 , 0.181	Depositor DCC
$R_{free}$ test set	1877 reflections (1.01%)	DCC
Wilson B-factor (Å <sup>2</sup> )	13.0	Xtriage
Anisotropy	0.667	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 46.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5566	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, NA, ACI, GLC, GLD, G6D, ACR, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.05	5/4955 (0.1%)	1.04	19/6751 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	588	GLU	CB-CG	-5.91	1.41	1.52
1	A	334	TRP	CZ3-CH2	-5.57	1.31	1.40
1	A	158	GLU	CB-CG	-5.51	1.41	1.52
1	A	194	GLU	CB-CG	-5.23	1.42	1.52
1	A	194	GLU	CG-CD	-5.11	1.44	1.51

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	502	ARG	NE-CZ-NH2	10.12	125.36	120.30
1	A	356	ASP	CB-CG-OD2	8.58	126.03	118.30
1	A	502	ARG	NE-CZ-NH1	-8.33	116.13	120.30
1	A	283	ASP	CB-CG-OD2	7.88	125.39	118.30
1	A	467	TYR	CB-CG-CD1	7.84	125.70	121.00
1	A	54	ASP	CB-CG-OD1	7.34	124.91	118.30
1	A	467	TYR	CB-CG-CD2	-7.03	116.78	121.00
1	A	473	ARG	NE-CZ-NH2	-6.83	116.88	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	283	ASP	CB-CG-OD1	-6.15	112.77	118.30
1	A	319	ASP	CB-CG-OD1	-6.12	112.79	118.30
1	A	412	PHE	CB-CG-CD2	6.00	125.00	120.80
1	A	54	ASP	CB-CG-OD2	-5.81	113.07	118.30
1	A	311	ASP	CB-CG-OD1	5.56	123.31	118.30
1	A	462	ARG	NE-CZ-NH2	-5.48	117.56	120.30
1	A	541	ASP	CB-CG-OD2	5.45	123.20	118.30
1	A	420	TYR	CB-CG-CD2	5.42	124.25	121.00
1	A	46	LYS	CD-CE-NZ	5.41	124.15	111.70
1	A	31	LYS	CD-CE-NZ	5.31	123.92	111.70
1	A	196	ASP	CB-CG-OD2	5.04	122.83	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	314	TYR	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4822	0	4443	26	0
2	A	10	0	1	0	0
3	A	57	0	50	4	0
4	A	24	0	22	0	0
5	A	18	0	18	0	0
6	A	12	0	11	0	0
7	A	44	0	43	0	0
8	A	4	0	0	0	0
9	A	1	0	0	0	0
10	A	574	0	0	5	1
All	All	5566	0	4588	27	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:52:LEU:HD23	1:A:61[B]:ILE:HD12	1.48	0.95
1:A:105:GLU:OE1	1:A:569:LYS:HE2	1.89	0.72
1:A:599:GLU:O	10:A:1558:HOH:O	2.09	0.70
1:A:371:SER:HB3	10:A:1402:HOH:O	1.99	0.63
1:A:307:THR:H	3:A:1001:GLC:H62	1.67	0.59
1:A:52:LEU:CD2	1:A:61[B]:ILE:HD12	2.28	0.59
1:A:119:GLN:O	1:A:119:GLN:HG3	2.05	0.56
1:A:383:VAL:CG1	1:A:388:ASP:HB3	2.36	0.55
1:A:340:ASN:OD1	1:A:371:SER:HB2	2.05	0.55
1:A:371:SER:OG	1:A:373[A]:ARG:HB3	2.07	0.54
1:A:75:ILE:HD13	1:A:88:LEU:HD21	1.92	0.52
1:A:307:THR:H	3:A:1001:GLC:C6	2.25	0.50
1:A:81:GLU:OE2	10:A:1471:HOH:O	2.19	0.49
1:A:36:ILE:HG21	1:A:88:LEU:HD22	1.96	0.48
1:A:515:ASN:HD21	1:A:521:SER:H	1.61	0.47
1:A:147:ASN:HD21	1:A:202:ARG:HH12	1.61	0.47
1:A:386[A]:VAL:HG13	10:A:1135:HOH:O	2.14	0.47
3:A:1001:GLC:O4	3:A:1001:GLC:O6	2.22	0.47
1:A:147:ASN:ND2	1:A:202:ARG:HH12	2.14	0.46
1:A:544:SER:OG	1:A:546:ASN:OD1	2.17	0.46
1:A:589:ASP:O	1:A:592[A]:TRP:CZ3	2.70	0.45
1:A:515:ASN:ND2	1:A:521:SER:H	2.14	0.45
1:A:239:TYR:CD1	3:A:1002:GLC:H2	2.53	0.43
1:A:333:ASP:OD1	10:A:1461:HOH:O	2.22	0.43
1:A:390:LYS:HD2	1:A:432[B]:SER:OG	2.19	0.43
1:A:386[A]:VAL:HG23	1:A:387:ASP:N	2.34	0.43
1:A:36:ILE:HG21	1:A:88:LEU:CD2	2.49	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:A:1384:HOH:O	10:A:1405:HOH:O[4_555]	2.07	0.13

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	592/599 (99%)	577 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	513/517 (99%)	508 (99%)	5 (1%)	80	51

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	119	GLN
1	A	289	ASP
1	A	371	SER
1	A	473	ARG
1	A	564	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	ASN
1	A	85	ASN
1	A	127	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	147	ASN
1	A	248	ASN
1	A	515	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 5 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GLC	A	1001	-	12,12,12	1.02	0	17,17,17	2.73	5 (29%)
3	GLC	A	1002	-	12,12,12	0.50	0	17,17,17	2.22	7 (41%)
2	G6D	A	801	3	9,10,11	1.10	1 (11%)	13,14,16	1.35	3 (23%)
3	GLC	A	802	2,4	11,11,12	1.12	0	13,15,17	2.29	8 (61%)
4	ACI	A	803	3,5	10,12,12	1.65	2 (20%)	11,17,17	1.50	2 (18%)
5	GLD	A	804	3,4	9,9,10	1.36	1 (11%)	10,12,14	1.38	1 (10%)
3	GLC	A	805	5,4	11,11,12	1.00	0	13,15,17	1.70	4 (30%)
4	ACI	A	806	3,5	10,12,12	2.49	6 (60%)	11,17,17	2.78	4 (36%)
5	GLD	A	807	3,4	9,9,10	1.08	0	10,12,14	2.62	3 (30%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GLC	A	808	5,6	11,11,12	1.10	1 (9%)	13,15,17	1.27	1 (7%)
6	BGC	A	809	3	12,12,12	1.17	0	17,17,17	1.43	3 (17%)
7	ACR	A	901	-	45,47,47	0.70	0	59,70,70	1.31	9 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	A	1001	-	-	0/2/22/22	0/1/1/1
3	GLC	A	1002	-	1/1/5/5	0/2/22/22	0/1/1/1
2	G6D	A	801	3	-	0/0/17/20	0/1/1/1
3	GLC	A	802	2,4	-	0/2/19/22	0/1/1/1
4	ACI	A	803	3,5	-	0/2/22/22	0/1/1/1
5	GLD	A	804	3,4	-	0/0/13/16	0/1/1/1
3	GLC	A	805	5,4	-	0/2/19/22	0/1/1/1
4	ACI	A	806	3,5	-	0/2/22/22	0/1/1/1
5	GLD	A	807	3,4	-	0/0/13/16	0/1/1/1
3	GLC	A	808	5,6	-	0/2/19/22	0/1/1/1
6	BGC	A	809	3	-	0/2/22/22	0/1/1/1
7	ACR	A	901	-	1/1/20/22	0/18/98/98	0/4/4/4

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	804	GLD	O5-C1	-3.38	1.38	1.43
4	A	806	ACI	C3-C4	-2.96	1.48	1.52
2	A	801	G6D	O2-C2	-2.30	1.38	1.43
4	A	806	ACI	C1-C7	2.15	1.55	1.49
3	A	808	GLC	C2-C3	2.16	1.55	1.52
4	A	806	ACI	O6-C6	2.63	1.50	1.41
4	A	803	ACI	C7-C5	2.80	1.36	1.32
4	A	806	ACI	O2-C2	2.89	1.49	1.43
4	A	803	ACI	O4-C4	3.20	1.48	1.42
4	A	806	ACI	C1-N1	3.68	1.59	1.47
4	A	806	ACI	C7-C5	3.68	1.38	1.32

All (50) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1001	GLC	C4-C3-C2	-6.84	98.78	110.84
4	A	806	ACI	O6-C6-C5	-6.39	96.90	112.63
5	A	807	GLD	C3-C4-C5	-6.22	101.64	111.27
3	A	1001	GLC	C6-C5-C4	-6.18	98.55	113.00
5	A	807	GLD	O5-C5-C4	-4.04	105.16	109.38
6	A	809	BGC	C3-C4-C5	-3.38	104.26	110.22
3	A	808	GLC	C2-C3-C4	-3.07	105.52	110.88
3	A	805	GLC	C2-C3-C4	-3.04	105.58	110.88
3	A	805	GLC	O5-C1-C2	-3.01	106.08	110.79
3	A	802	GLC	O2-C2-C3	-2.97	104.33	110.17
3	A	802	GLC	O4-C4-C3	-2.94	103.95	110.36
3	A	802	GLC	C3-C4-C5	-2.90	105.11	110.22
3	A	1002	GLC	C4-C3-C2	-2.87	105.77	110.84
6	A	809	BGC	C1-O5-C5	-2.87	108.21	113.39
3	A	1001	GLC	O3-C3-C4	-2.82	104.23	110.36
7	A	901	ACR	O5C-C1C-C2C	-2.64	105.21	110.30
3	A	802	GLC	O5-C1-C2	-2.60	106.71	110.79
7	A	901	ACR	O2D-C2D-C3D	-2.57	104.76	110.36
4	A	806	ACI	C2-C3-C4	-2.52	106.78	110.46
4	A	803	ACI	C7-C1-N1	-2.47	106.29	110.83
7	A	901	ACR	C7A-C1A-N4B	-2.37	107.45	110.88
5	A	807	GLD	C6-C5-C4	-2.34	109.64	113.39
2	A	801	G6D	O2-C2-C3	-2.27	105.71	110.17
3	A	802	GLC	O3-C3-C2	-2.26	105.91	110.02
7	A	901	ACR	O6A-C6A-C5A	-2.25	107.10	112.63
2	A	801	G6D	O3-C3-C2	-2.24	105.94	110.02
6	A	809	BGC	C1-C2-C3	-2.20	106.68	110.65
7	A	901	ACR	C2A-C3A-C4A	-2.17	107.29	110.46
3	A	805	GLC	O4-C4-C3	-2.15	105.67	110.36
3	A	802	GLC	O3-C3-C4	-2.11	105.77	110.36
7	A	901	ACR	C1B-C2B-C3B	-2.09	106.09	109.98
3	A	1002	GLC	O5-C1-C2	2.10	113.53	110.04
4	A	803	ACI	O4-C4-C3	2.10	114.61	110.28
3	A	802	GLC	O2-C2-C1	2.17	113.59	109.18
3	A	1001	GLC	O2-C2-C1	2.24	114.40	109.75
7	A	901	ACR	O3A-C3A-C4A	2.24	113.46	109.48
3	A	1002	GLC	C1-O5-C5	2.39	117.70	113.39
3	A	1002	GLC	O5-C5-C6	2.48	112.36	106.41
2	A	801	G6D	C1-C2-C3	2.59	112.94	109.65
7	A	901	ACR	C2D-C3D-C4D	2.72	115.25	109.61
3	A	805	GLC	C1-O5-C5	2.89	116.14	112.17
3	A	1002	GLC	O5-C5-C4	3.06	115.30	109.66
7	A	901	ACR	O2D-C2D-C1D	3.22	116.44	109.75

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	806	ACI	C2-C1-N1	3.24	117.98	111.40
5	A	804	GLD	O5-C5-C6	3.72	111.75	106.17
3	A	802	GLC	C1-C2-C3	3.75	114.41	109.65
3	A	1001	GLC	O5-C5-C4	4.30	117.58	109.66
4	A	806	ACI	C7-C1-N1	4.48	119.03	110.83
3	A	1002	GLC	O1-C1-O5	4.55	123.67	110.20
3	A	1002	GLC	O1-C1-C2	4.69	122.65	109.42

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
7	A	901	ACR	C1D
3	A	1002	GLC	C1

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1001	GLC	3	0
3	A	1002	GLC	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	585/599 (97%)	0.11	29 (4%) 30 34	10, 17, 28, 39	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	454	SER	4.8
1	A	96	TYR	4.5
1	A	453	GLY	4.4
1	A	97	ASP	4.3
1	A	452	GLU	4.2
1	A	599	GLU	4.0
1	A	140	THR	3.8
1	A	119	GLN	3.4
1	A	577	GLU	3.2
1	A	592[A]	TRP	3.0
1	A	95	ILE	3.0
1	A	118	ASP	3.0
1	A	137	GLU	2.9
1	A	513	ASP	2.9
1	A	537	LEU	2.7
1	A	58	ASP	2.7
1	A	231	LEU	2.7
1	A	558	ASP	2.6
1	A	169	LEU	2.6
1	A	141	GLU	2.4
1	A	467	TYR	2.3
1	A	489	GLU	2.2
1	A	535	LEU	2.2
1	A	229	ALA	2.2
1	A	57	THR	2.2
1	A	497	LEU	2.2
1	A	470	ILE	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	466	ALA	2.1
1	A	471	LEU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	GLC	A	1001	12/12	0.85	0.30	12.94	26,40,42,46	0
4	ACI	A	803	12/12	0.90	0.18	6.45	20,22,29,38	0
6	BGC	A	809	12/12	0.88	0.26	4.59	26,38,43,47	0
3	GLC	A	802	11/12	0.91	0.17	3.87	25,26,30,35	0
8	CA	A	705	1/1	1.00	0.17	3.23	30,30,30,30	0
7	ACR	A	901	44/44	0.92	0.12	1.16	14,23,48,52	0
4	ACI	A	806	12/12	0.93	0.09	0.25	14,18,24,30	0
3	GLC	A	805	11/12	0.97	0.06	-0.29	15,17,21,24	0
8	CA	A	702	1/1	1.00	0.07	-0.51	10,10,10,10	0
8	CA	A	703	1/1	1.00	0.04	-0.99	14,14,14,14	0
5	GLD	A	807	9/10	0.98	0.05	-1.02	13,14,17,19	0
8	CA	A	701	1/1	1.00	0.03	-1.68	11,11,11,11	0
9	NA	A	704	1/1	1.00	0.03	-1.73	10,10,10,10	0
3	GLC	A	808	11/12	0.96	0.09	-	17,20,24,29	0
3	GLC	A	1002	12/12	0.64	0.24	-	72,74,74,75	0
2	G6D	A	801	10/11	0.89	0.28	-	26,35,41,41	0
5	GLD	A	804	9/10	0.95	0.08	-	21,22,28,28	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.