



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 10, 2018 – 06:56 PM EST

PDB ID : 6BDH
Title : Crystal structure of human CYP3A4 bound to an inhibitor
Authors : Sevrioukova, I.
Deposited on : 2017-10-23
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1062 (2.26-2.26)
Clashscore	112137	1178 (2.26-2.26)
Ramachandran outliers	110173	1145 (2.26-2.26)
Sidechain outliers	110143	1146 (2.26-2.26)
RSRZ outliers	101464	1066 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	487	<div> <div>11%</div> <div>68%</div> <div>20%</div> <div>10%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	DEV	A	602	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3656 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450 3A4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	439	Total	C	N	O	S	0	7	0
			3546	2314	574	631	27			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	LEU	deletion	UNP P08684
A	?	-	ILE	deletion	UNP P08684
A	?	-	PRO	deletion	UNP P08684
A	?	-	ASP	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	ALA	deletion	UNP P08684
A	?	-	MET	deletion	UNP P08684
A	?	-	GLU	deletion	UNP P08684
A	?	-	THR	deletion	UNP P08684
A	?	-	TRP	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	ALA	deletion	UNP P08684
A	?	-	VAL	deletion	UNP P08684
A	?	-	SER	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	VAL	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	?	-	LEU	deletion	UNP P08684
A	504	HIS	-	expression tag	UNP P08684
A	505	HIS	-	expression tag	UNP P08684
A	506	HIS	-	expression tag	UNP P08684
A	507	HIS	-	expression tag	UNP P08684

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



- [illegible]

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			37	29	4	3	1		


- 

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	30	Total	O	0	0
			30	30		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cytochrome P450 3A4



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	76.37Å 99.88Å 125.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.19 – 2.25 38.19 – 2.25	Depositor EDS
% Data completeness (in resolution range)	95.7 (38.19-2.25) 95.7 (38.19-2.25)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.74 (at 2.24Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575)	Depositor
R, R_{free}	0.206 , 0.280 0.223 , 0.288	Depositor DCC
R_{free} test set	1138 reflections (5.12%)	DCC
Wilson B-factor (Å ²)	61.0	Xtriage
Anisotropy	0.066	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 78.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3656	wwPDB-VP
Average B, all atoms (Å ²)	102.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, DEV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.43	0/3652	0.60	0/4938

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3546	0	3647	78	0
2	A	43	0	30	3	0
3	A	37	0	0	0	0
4	A	30	0	0	1	0
All	All	3656	0	3677	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:257:LYS:H	1:A:258:GLU:HB3	1.14	1.05
1:A:257:LYS:N	1:A:258:GLU:HB3	1.80	0.96
1:A:257:LYS:HB3	1:A:258:GLU:HA	1.45	0.95
1:A:258:GLU:O	1:A:258:GLU:HG2	1.74	0.84
1:A:275:MET:HE2	1:A:290:LEU:HD21	1.63	0.79
1:A:253:VAL:HB	1:A:296:VAL:HG22	1.67	0.75
1:A:249:LEU:HA	1:A:252:SER:HB3	1.72	0.71
1:A:257:LYS:HB3	1:A:258:GLU:CA	2.22	0.69
1:A:421:LYS:HA	1:A:424:LYS:HG2	1.75	0.68
1:A:199:PRO:C	1:A:200:GLN:HG3	2.12	0.68
1:A:139:SER:HB3	1:A:446:ARG:HH22	1.59	0.67
1:A:226:PHE:HA	1:A:228:PHE:CE2	2.34	0.63
1:A:128:ARG:NH2	1:A:289:ALA:O	2.32	0.62
1:A:251:LYS:O	1:A:255:ARG:N	2.30	0.62
1:A:252:SER:HA	1:A:255:ARG:HB3	1.81	0.62
1:A:270:ASP:O	1:A:274:LEU:HG	2.00	0.61
1:A:166:THR:O	1:A:168:LYS:N	2.26	0.61
1:A:200:GLN:O	1:A:201:ASP:O	2.19	0.60
1:A:101:VAL:HG21	1:A:381[A]:VAL:HG11	1.84	0.60
1:A:458:ARG:NH2	1:A:461:GLN:OE1	2.31	0.59
1:A:87:PRO:HG3	1:A:431:ILE:HD11	1.83	0.59
1:A:194:ASP:O	1:A:196:LEU:N	2.34	0.59
1:A:235:VAL:HG13	1:A:236:LEU:HD23	1.85	0.59
1:A:257:LYS:CB	1:A:258:GLU:HA	2.27	0.58
2:A:601:HEM:HBC2	2:A:601:HEM:HHD	1.87	0.56
1:A:132:LEU:HD11	1:A:278:SER:OG	2.04	0.56
1:A:58:CYS:HB3	1:A:399:TYR:CD2	2.40	0.56
1:A:477:LEU:HD13	1:A:483:LEU:HD11	1.87	0.55
1:A:202:PRO:HB3	1:A:245:VAL:CG1	2.36	0.55
1:A:172:LEU:HB3	1:A:176:PHE:CE2	2.43	0.53
1:A:202:PRO:HB3	1:A:245:VAL:HG11	1.91	0.52
1:A:276:ILE:HA	1:A:279:GLN:HB3	1.90	0.52
1:A:202:PRO:HG2	1:A:203:PHE:H	1.75	0.51
1:A:273:GLN:HG3	1:A:277:ASP:OD2	2.09	0.51
1:A:200:GLN:HB3	1:A:248:PHE:HE2	1.76	0.51
1:A:478:SER:C	1:A:480:GLY:H	2.14	0.51
1:A:446:ARG:HD3	4:A:703:HOH:O	2.11	0.50
1:A:232:ILE:O	1:A:235:VAL:HG12	2.12	0.49
1:A:356:LEU:O	1:A:360:VAL:HG23	2.12	0.49
1:A:436:GLY:HA3	2:A:601:HEM:HBA1	1.95	0.49
1:A:202:PRO:CB	1:A:245:VAL:HG11	2.43	0.48
1:A:126:TRP:CZ2	1:A:440:ARG:HG3	2.49	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:217:ASP:N	1:A:218:PRO:HD3	2.27	0.48
1:A:247:ASN:HA	1:A:250:ARG:HB2	1.95	0.48
1:A:420:SER:O	1:A:424:LYS:N	2.43	0.48
1:A:257:LYS:HD2	1:A:257:LYS:HA	1.74	0.47
1:A:172:LEU:HB3	1:A:176:PHE:CD2	2.50	0.47
1:A:243:ARG:CZ	1:A:243:ARG:HB2	2.45	0.46
1:A:196:LEU:HA	1:A:196:LEU:HD12	1.84	0.46
1:A:71:VAL:HG13	1:A:82:LEU:HD11	1.97	0.46
1:A:257:LYS:H	1:A:258:GLU:CB	2.06	0.45
1:A:132:LEU:HD21	1:A:290:LEU:HD22	1.97	0.45
1:A:29:SER:OG	1:A:29:SER:O	2.33	0.45
1:A:334:GLU:OE2	1:A:353:MET:HB3	2.16	0.44
1:A:169:PRO:HG3	1:A:468:CYS:SG	2.57	0.44
1:A:181:MET:HE2	1:A:307:TYR:CD2	2.52	0.44
1:A:76:ASP:HB3	1:A:224:THR:CG2	2.47	0.44
1:A:117:ALA:HB1	1:A:301:ILE:HG13	2.00	0.43
1:A:199:PRO:O	1:A:200:GLN:HG3	2.17	0.43
1:A:91:LYS:HG3	1:A:430:TYR:CZ	2.54	0.43
1:A:335:ILE:HG21	1:A:461:GLN:HG3	2.01	0.43
1:A:110:PRO:HG3	1:A:233:LEU:HB3	2.00	0.42
1:A:236:LEU:C	1:A:238:ILE:H	2.23	0.42
1:A:106:ARG:HG3	1:A:393:VAL:HG21	2.01	0.42
1:A:370:ALA:O	1:A:371:MET:HB2	2.19	0.42
1:A:442:CYS:HB2	2:A:601:HEM:NA	2.34	0.42
1:A:421:LYS:HG3	1:A:424:LYS:HE3	2.02	0.41
1:A:200:GLN:HB3	1:A:248:PHE:CE2	2.54	0.41
1:A:294:GLU:O	1:A:298:GLN:HG2	2.19	0.41
1:A:483:LEU:HA	1:A:483:LEU:HD12	1.71	0.41
1:A:428:ASP:HB3	1:A:431:ILE:HG12	2.02	0.41
1:A:458:ARG:HD2	1:A:458:ARG:HA	1.78	0.41
1:A:401:LEU:HB3	1:A:431:ILE:HG23	2.02	0.40
1:A:188:SER:HB3	1:A:271:PHE:HD2	1.86	0.40
1:A:471:THR:HG22	1:A:473:ILE:HG13	2.01	0.40
1:A:128:ARG:HH21	1:A:289:ALA:C	2.25	0.40
1:A:232:ILE:O	1:A:236:LEU:HG	2.22	0.40
1:A:219:PHE:CE2	1:A:240:VAL:HG12	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	438/487 (90%)	400 (91%)	33 (8%)	5 (1%)	17 12

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	202	PRO
1	A	167	GLY
1	A	201	ASP
1	A	218	PRO
1	A	479	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	406/443 (92%)	396 (98%)	10 (2%)	53 63

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	127	LYS
1	A	184	ILE
1	A	192	ASN
1	A	219	PHE
1	A	224	THR
1	A	232	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	257	LYS
1	A	258	GLU
1	A	458	ARG
1	A	482	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	601	1,3	28,50,50	1.99	9 (32%)	17,82,82	2.07	6 (35%)
3	DEV	A	602	2	38,39,39	1.89	12 (31%)	50,51,51	1.75	8 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	601	1,3	-	0/6/54/54	0/0/8/8
3	DEV	A	602	2	-	0/32/32/32	0/3/3/3

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	HEM	C3C-C2C	-4.17	1.34	1.40
2	A	601	HEM	C3B-C2B	-3.40	1.35	1.40
3	A	602	DEV	C31-C32	-2.76	1.44	1.51
2	A	601	HEM	C4A-NA	2.02	1.40	1.36
2	A	601	HEM	CMB-C2B	2.07	1.56	1.51
3	A	602	DEV	C16-C15	2.12	1.42	1.39
2	A	601	HEM	C4C-NC	2.19	1.39	1.36
3	A	602	DEV	C13-N14	2.21	1.47	1.45
2	A	601	HEM	CMC-C2C	2.23	1.56	1.51
3	A	602	DEV	C24-C25	2.37	1.56	1.51
3	A	602	DEV	C15-N14	2.46	1.44	1.39
3	A	602	DEV	C10-S11	2.51	1.87	1.81
3	A	602	DEV	O05-C06	2.72	1.40	1.34
2	A	601	HEM	C1C-NC	3.01	1.40	1.36
3	A	602	DEV	C09-N08	3.11	1.52	1.46
3	A	602	DEV	C13-C21	3.15	1.61	1.52
3	A	602	DEV	C12-S11	3.25	1.89	1.81
2	A	601	HEM	C3C-CAC	3.80	1.55	1.47
3	A	602	DEV	C21-N23	4.41	1.42	1.33
2	A	601	HEM	C3B-CAB	4.56	1.56	1.47
3	A	602	DEV	C24-N23	4.91	1.56	1.46

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	HEM	CBD-CAD-C3D	-4.84	103.23	112.47
2	A	601	HEM	CMA-C3A-C4A	-3.53	123.04	128.46
3	A	602	DEV	O07-C06-N08	-3.27	119.25	124.87
2	A	601	HEM	CMD-C2D-C1D	-2.59	124.48	128.46
3	A	602	DEV	C01-C02-C03	-2.51	104.39	111.16
3	A	602	DEV	O05-C06-O07	-2.40	120.88	125.56
2	A	601	HEM	CBA-CAA-C2A	-2.17	108.34	112.48
2	A	601	HEM	CMA-C3A-C2A	2.25	129.19	124.94
2	A	601	HEM	CMB-C2B-C3B	2.57	129.66	124.89
3	A	602	DEV	C13-C21-N23	2.95	122.53	116.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	DEV	C25-C24-N23	3.04	119.66	112.96
3	A	602	DEV	O05-C06-N08	3.20	115.79	110.06
3	A	602	DEV	C24-N23-C21	3.35	126.93	122.39
3	A	602	DEV	C12-S11-C10	7.11	114.22	102.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	HEM	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	439/487 (90%)	0.60	52 (11%) 5 5	47, 96, 163, 227	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	253	VAL	6.8
1	A	226	PHE	6.5
1	A	275	MET	5.9
1	A	255	ARG	5.3
1	A	278	SER	5.2
1	A	290	LEU	5.2
1	A	245	VAL	5.1
1	A	203	PHE	5.0
1	A	256	MET	4.8
1	A	251	LYS	4.8
1	A	221	LEU	4.4
1	A	257	LYS	4.3
1	A	248	PHE	4.0
1	A	240	VAL	3.7
1	A	191	VAL	3.7
1	A	241	PHE	3.6
1	A	165	GLU	3.4
1	A	277	ASP	3.2
1	A	271	PHE	3.1
1	A	252	SER	3.0
1	A	310	THR	3.0
1	A	198	ASN	2.9
1	A	309	THR	2.9
1	A	371	MET	2.8
1	A	295	LEU	2.6
1	A	276	ILE	2.6
1	A	289	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	222	SER	2.5
1	A	270	ASP	2.5
1	A	274	LEU	2.5
1	A	199	PRO	2.5
1	A	244	GLU	2.5
1	A	192	ASN	2.5
1	A	197	ASN	2.5
1	A	444	GLY	2.4
1	A	169	PRO	2.4
1	A	229	LEU	2.3
1	A	189	PHE	2.3
1	A	479	LEU	2.3
1	A	137	PHE	2.3
1	A	442	CYS	2.2
1	A	202	PRO	2.2
1	A	217	ASP	2.2
1	A	483	LEU	2.2
1	A	109	GLY	2.2
1	A	67	LYS	2.1
1	A	279	GLN	2.1
1	A	493	VAL	2.1
1	A	341	ASN	2.0
1	A	314	LEU	2.0
1	A	482	LEU	2.0
1	A	254	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	DEV	A	602	37/37	0.86	0.37	2.02	59,128,155,156	0
2	HEM	A	601	43/43	0.99	0.23	0.78	48,63,71,76	0

6.5 Other polymers ⓘ

There are no such residues in this entry.