



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 06:16 pm GMT

PDB ID : 2BFO
Title : LEISHMANIA MAJOR PTERIDINE REDUCTASE 1 IN COMPLEX WITH NADPH
Authors : Schuettelkopf, A.W.; Hunter, W.N.
Deposited on : 2004-12-10
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2542 (2.60-2.60)
Clashscore	112137	2895 (2.60-2.60)
Ramachandran outliers	110173	2848 (2.60-2.60)
Sidechain outliers	110143	2848 (2.60-2.60)
RSRZ outliers	101464	2550 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	288	<div> <div>2%</div> <div> <div></div> <div>76%</div> <div>16%</div> <div>• 7%</div> </div> </div>
1	B	288	<div> <div>4%</div> <div> <div></div> <div>70%</div> <div>21%</div> <div>• 8%</div> </div> </div>
1	C	288	<div> <div>6%</div> <div> <div></div> <div>74%</div> <div>14%</div> <div>• 11%</div> </div> </div>
1	D	288	<div> <div>6%</div> <div> <div></div> <div>74%</div> <div>15%</div> <div>11%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	1291	-	-	-	X

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 8381 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PTERIDINE REDUCTASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	0	0
			1991	1255	354	371	11			
1	B	266	Total	C	N	O	S	0	0	0
			1987	1253	354	369	11			
1	C	255	Total	C	N	O	S	0	0	0
			1903	1197	341	355	10			
1	D	256	Total	C	N	O	S	0	0	0
			1906	1201	342	353	10			

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	118	Total	O	0	0
			118	118		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	105	Total 105	O 105	0	0
4	C	80	Total 80	O 80	0	0
4	D	75	Total 75	O 75	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PTERIDINE REDUCTASE 1




• Molecule 1: PTERIDINE REDUCTASE 1



• Molecule 1: PTERIDINE REDUCTASE 1



• Molecule 1: PTERIDINE REDUCTASE 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	94.31Å 103.49Å 137.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.50 – 2.60 24.55 – 2.59	Depositor EDS
% Data completeness (in resolution range)	100.0 (24.50-2.60) 94.8 (24.55-2.59)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.64 (at 2.60Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.226 , 0.269 0.229 , 0.273	Depositor DCC
R_{free} test set	2055 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	28.4	Xtriage
Anisotropy	0.172	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	8381	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.76 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.7564e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.63	0/2030	0.75	2/2765 (0.1%)
1	B	0.62	0/2026	0.75	3/2760 (0.1%)
1	C	0.55	0/1936	0.70	1/2630 (0.0%)
1	D	0.55	0/1940	0.67	0/2638
All	All	0.59	0/7932	0.72	6/10793 (0.1%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	142	ASP	CB-CG-OD2	6.22	123.90	118.30
1	B	65	ASP	CB-CG-OD2	6.14	123.83	118.30
1	C	142	ASP	CB-CG-OD2	5.77	123.49	118.30
1	A	181	ASP	CB-CG-OD2	5.53	123.27	118.30
1	A	104	ASP	CB-CG-OD2	5.04	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1991	0	1999	33	0
1	B	1987	0	1998	44	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1903	0	1918	32	0
1	D	1906	0	1928	32	0
2	A	48	0	26	1	0
2	B	48	0	26	0	0
2	C	48	0	26	1	0
2	D	48	0	26	1	0
3	A	12	0	18	1	0
3	B	4	0	6	1	0
3	C	4	0	6	0	0
3	D	4	0	6	1	0
4	A	118	0	0	1	0
4	B	105	0	0	8	0
4	C	80	0	0	1	0
4	D	75	0	0	4	0
All	All	8381	0	7983	120	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 120 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:121:ASN:HB3	4:B:2013:HOH:O	1.68	0.93
1:B:12:THR:HG23	4:B:2032:HOH:O	1.72	0.90
1:B:278:LYS:HB2	1:D:275:THR:HG21	1.51	0.89
1:A:275:THR:HG21	1:C:278:LYS:HB2	1.55	0.88
1:B:275:THR:HG21	1:D:278:LYS:HB2	1.55	0.87

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/288 (91%)	247 (95%)	13 (5%)	1 (0%)	38	63
1	B	260/288 (90%)	245 (94%)	13 (5%)	2 (1%)	22	44
1	C	247/288 (86%)	234 (95%)	11 (4%)	2 (1%)	22	44
1	D	248/288 (86%)	236 (95%)	12 (5%)	0	100	100
All	All	1016/1152 (88%)	962 (95%)	49 (5%)	5 (0%)	32	58

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	230	VAL
1	B	73	PRO
1	A	180	VAL
1	B	180	VAL
1	C	180	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	207/223 (93%)	204 (99%)	3 (1%)	71	89
1	B	207/223 (93%)	204 (99%)	3 (1%)	71	89
1	C	198/223 (89%)	195 (98%)	3 (2%)	70	88
1	D	198/223 (89%)	196 (99%)	2 (1%)	80	93
All	All	810/892 (91%)	799 (99%)	11 (1%)	71	89

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	168	LYS
1	B	242	ARG
1	C	242	ARG
1	B	133	ARG
1	C	193	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	63	GLN
1	D	216	GLN
1	B	216	GLN
1	A	169	HIS
1	C	216	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NDP	A	1289	-	43,52,52	1.33	4 (9%)	49,80,80	1.65	2 (4%)
3	EDO	A	1290	-	3,3,3	0.84	0	2,2,2	0.43	0
3	EDO	A	1291	-	3,3,3	0.25	0	2,2,2	0.79	0
3	EDO	A	1292	-	3,3,3	0.45	0	2,2,2	0.18	0
2	NDP	B	1289	-	43,52,52	1.31	5 (11%)	49,80,80	1.62	3 (6%)
3	EDO	B	1290	-	3,3,3	0.37	0	2,2,2	0.13	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NDP	C	1289	-	43,52,52	1.44	4 (9%)	49,80,80	1.62	2 (4%)
3	EDO	C	1290	-	3,3,3	0.48	0	2,2,2	0.27	0
2	NDP	D	1289	-	43,52,52	1.40	3 (6%)	49,80,80	1.91	2 (4%)
3	EDO	D	1290	-	3,3,3	0.40	0	2,2,2	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	1289	-	-	0/30/77/77	0/5/5/5
3	EDO	A	1290	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1291	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1292	-	-	0/1/1/1	0/0/0/0
2	NDP	B	1289	-	-	0/30/77/77	0/5/5/5
3	EDO	B	1290	-	-	0/1/1/1	0/0/0/0
2	NDP	C	1289	-	-	0/30/77/77	0/5/5/5
3	EDO	C	1290	-	-	0/1/1/1	0/0/0/0
2	NDP	D	1289	-	-	0/30/77/77	0/5/5/5
3	EDO	D	1290	-	-	0/1/1/1	0/0/0/0

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1289	NDP	C2A-N1A	2.05	1.37	1.33
2	B	1289	NDP	O4B-C1B	2.15	1.44	1.41
2	A	1289	NDP	C2A-N1A	2.28	1.38	1.33
2	A	1289	NDP	C2A-N3A	2.33	1.36	1.32
2	B	1289	NDP	C6N-C5N	2.40	1.37	1.33

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1289	NDP	N3A-C2A-N1A	-11.67	118.69	128.86
2	A	1289	NDP	N3A-C2A-N1A	-9.54	120.55	128.86
2	B	1289	NDP	N3A-C2A-N1A	-9.51	120.58	128.86
2	C	1289	NDP	N3A-C2A-N1A	-9.21	120.83	128.86
2	C	1289	NDP	C4B-O4B-C1B	-4.17	105.33	109.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1289	NDP	1	0
3	A	1291	EDO	1	0
3	B	1290	EDO	1	0
2	C	1289	NDP	1	0
2	D	1289	NDP	1	0
3	D	1290	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	267/288 (92%)	-0.03	6 (2%) 62 56	22, 34, 49, 67	1 (0%)
1	B	266/288 (92%)	0.11	11 (4%) 38 30	18, 34, 52, 70	1 (0%)
1	C	255/288 (88%)	-0.04	17 (6%) 19 13	22, 34, 56, 82	3 (1%)
1	D	256/288 (88%)	0.08	16 (6%) 21 15	22, 34, 56, 86	2 (0%)
All	All	1044/1152 (90%)	0.03	50 (4%) 31 24	18, 34, 54, 86	7 (0%)

The worst 5 of 50 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	81	ALA	6.5
1	B	121	ASN	5.5
1	A	132	ASP	4.4
1	D	72	ALA	4.3
1	B	120	ARG	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	EDO	A	1291	4/4	0.92	0.26	8.82	35,37,38,44	0
3	EDO	A	1290	4/4	0.89	0.28	1.72	17,19,23,24	0
3	EDO	C	1290	4/4	0.96	0.15	0.34	35,36,37,37	0
3	EDO	D	1290	4/4	0.98	0.13	0.17	39,41,43,43	0
2	NDP	B	1289	48/48	0.95	0.15	-0.52	26,36,40,42	0
2	NDP	D	1289	48/48	0.95	0.12	-0.66	27,36,39,42	0
2	NDP	A	1289	48/48	0.95	0.15	-0.69	26,36,40,42	0
2	NDP	C	1289	48/48	0.95	0.12	-0.96	27,36,39,42	0
3	EDO	A	1292	4/4	0.97	0.11	-1.93	24,25,26,30	0
3	EDO	B	1290	4/4	0.98	0.10	-2.69	20,21,22,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.