



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 15, 2017 – 05:06 PM EDT

PDB ID : 3BGE  
Title : Crystal structure of the C-terminal fragment of AAA+ATPase from Haemophilus influenzae  
Authors : Ramagopal, U.A.; Patskovsky, Y.; Bonanno, J.B.; Meyer, A.J.; Toro, R.; Freeman, J.; Adams, J.; Koss, J.; Maletic, M.; Gheyi, T.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : unknown  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1923 (1.86-1.86)
Clashscore	112137	2083 (1.86-1.86)
Ramachandran outliers	110173	2060 (1.86-1.86)
Sidechain outliers	110143	2060 (1.86-1.86)
RSRZ outliers	101464	1932 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	201	
1	B	201	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2855 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Predicted ATPase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	163	Total	C	N	O	S	0	3	0
			1300	829	226	240	5			
1	B	163	Total	C	N	O	S	0	5	0
			1320	843	230	240	7			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	248	MET	-	EXPRESSION TAG	UNP Q4QL22
A	249	SER	-	EXPRESSION TAG	UNP Q4QL22
A	250	LEU	-	EXPRESSION TAG	UNP Q4QL22
A	340	SER	ALA	ENGINEERED	UNP Q4QL22
A	441	GLU	-	EXPRESSION TAG	UNP Q4QL22
A	442	GLY	-	EXPRESSION TAG	UNP Q4QL22
A	443	HIS	-	EXPRESSION TAG	UNP Q4QL22
A	444	HIS	-	EXPRESSION TAG	UNP Q4QL22
A	445	HIS	-	EXPRESSION TAG	UNP Q4QL22
A	446	HIS	-	EXPRESSION TAG	UNP Q4QL22
A	447	HIS	-	EXPRESSION TAG	UNP Q4QL22
A	448	HIS	-	EXPRESSION TAG	UNP Q4QL22
B	248	MET	-	EXPRESSION TAG	UNP Q4QL22
B	249	SER	-	EXPRESSION TAG	UNP Q4QL22
B	250	LEU	-	EXPRESSION TAG	UNP Q4QL22
B	340	SER	ALA	ENGINEERED	UNP Q4QL22
B	441	GLU	-	EXPRESSION TAG	UNP Q4QL22
B	442	GLY	-	EXPRESSION TAG	UNP Q4QL22
B	443	HIS	-	EXPRESSION TAG	UNP Q4QL22
B	444	HIS	-	EXPRESSION TAG	UNP Q4QL22
B	445	HIS	-	EXPRESSION TAG	UNP Q4QL22
B	446	HIS	-	EXPRESSION TAG	UNP Q4QL22
B	447	HIS	-	EXPRESSION TAG	UNP Q4QL22
B	448	HIS	-	EXPRESSION TAG	UNP Q4QL22

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	128	Total	O	0	0
			128	128		
3	B	92	Total	O	0	0
			92	92		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Predicted ATPase



#### • Molecule 1: Predicted ATPase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.08 Å 93.08 Å 83.41 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.12 – 1.85 21.94 – 1.85	Depositor EDS
% Data completeness (in resolution range)	96.4 (23.12-1.85) 96.4 (21.94-1.85)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.36 (at 1.85 Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.187 , 0.240 0.187 , 0.238	Depositor DCC
$R_{free}$ test set	1468 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	20.2	Xtriage
Anisotropy	0.002	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 60.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.031 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2855	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.63	0/1337	0.67	0/1818
1	B	0.64	0/1363	0.72	1/1849 (0.1%)
All	All	0.63	0/2700	0.70	1/3667 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	271	ASP	CB-CG-OD2	5.53	123.28	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1300	0	1281	29	0
1	B	1320	0	1315	29	0
2	A	5	0	0	0	0
2	B	10	0	0	0	0
3	A	128	0	0	2	0
3	B	92	0	0	2	0
All	All	2855	0	2596	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:319[B]:CYS:SG	1:A:327:GLU:HG3	2.00	1.02
1:B:295:LEU:HD22	1:B:310[B]:MET:SD	2.10	0.92
1:A:321:THR:HG23	1:B:321:THR:OG1	1.82	0.78
1:A:433:LEU:O	1:A:434:ARG:HB2	1.84	0.78
1:B:253[A]:ARG:HH11	1:B:253[A]:ARG:CG	1.98	0.76
1:B:295:LEU:CD2	1:B:310[B]:MET:SD	2.76	0.73
1:B:319[B]:CYS:SG	1:B:327:GLU:HG3	2.29	0.72
1:A:319[B]:CYS:SG	1:A:327:GLU:CG	2.77	0.71
1:B:258:ILE:HG12	1:B:293:ARG:HD3	1.70	0.71
1:B:253[A]:ARG:HH11	1:B:253[A]:ARG:HG3	1.55	0.69
1:A:295:LEU:HD22	1:A:310[B]:MET:SD	2.33	0.69
1:A:289:TYR:CE1	1:A:293:ARG:NH2	2.64	0.66
1:A:346:ASN:O	1:A:350[A]:THR:HG22	1.96	0.64
1:A:321:THR:HG23	1:B:321:THR:CG2	2.29	0.63
1:A:274:LEU:HD11	1:A:337:ILE:HD11	1.84	0.59
1:B:253[A]:ARG:NH1	1:B:253[A]:ARG:CG	2.66	0.57
1:A:289:TYR:CZ	1:A:293:ARG:NH2	2.73	0.57
1:B:253[B]:ARG:NH1	3:B:514:HOH:O	2.38	0.57
1:A:358:GLN:NE2	1:A:414:TYR:OH	2.38	0.56
1:B:427:LYS:O	1:B:431:GLU:HG3	2.05	0.55
1:A:346:ASN:ND2	1:A:350[B]:THR:HG23	2.22	0.54
1:B:253[B]:ARG:NH2	1:B:253[B]:ARG:HB2	2.23	0.54
1:A:321:THR:CG2	1:B:321:THR:HG23	2.39	0.53
1:A:433:LEU:O	1:A:434:ARG:CB	2.55	0.52
1:A:278:ALA:O	1:A:282:THR:HG23	2.10	0.52
1:A:321:THR:CG2	1:B:321:THR:CG2	2.89	0.51
1:A:419:ASN:HD21	1:A:427:LYS:NZ	2.09	0.50
1:B:319[B]:CYS:SG	1:B:327:GLU:CG	2.98	0.50
1:A:295:LEU:CD2	1:A:310[B]:MET:SD	3.00	0.49
1:A:321:THR:OG1	1:B:321:THR:HG23	2.12	0.49
1:A:434:ARG:HH11	1:A:434:ARG:HG2	1.78	0.48
1:A:321:THR:HG23	1:B:321:THR:HG23	1.95	0.48
1:A:308:ARG:HD3	3:A:535:HOH:O	2.13	0.48
1:A:346:ASN:ND2	1:A:350[B]:THR:CG2	2.78	0.46
1:B:422[B]:MET:HE3	1:B:422[B]:MET:HA	1.96	0.46
1:A:434:ARG:HH11	1:A:434:ARG:CG	2.29	0.46
1:B:311:GLN:NE2	3:B:532:HOH:O	2.47	0.46

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:379:MET:HG3	1:B:380:LYS:HG3	1.98	0.46
1:B:307:PRO:CD	1:B:422[B]:MET:HG3	2.46	0.45
1:A:429:LYS:O	1:A:433:LEU:HD13	2.17	0.45
1:B:274:LEU:HD11	1:B:337:ILE:HD11	1.99	0.44
1:B:255:TYR:O	1:B:258:ILE:HG22	2.18	0.43
1:B:307:PRO:HD3	1:B:422[B]:MET:HG3	2.01	0.43
1:A:282:THR:HG23	3:A:513:HOH:O	2.19	0.43
1:B:268:SER:HA	1:B:340:SER:O	2.18	0.42
1:A:425:GLN:NE2	1:A:425:GLN:H	2.18	0.42
1:B:380:LYS:HE2	1:B:380:LYS:HB2	1.90	0.42
1:A:324:GLY:HA2	1:B:288:LEU:HD12	2.01	0.41
1:B:258:ILE:HD12	1:B:258:ILE:HA	1.93	0.41
1:B:294:LEU:HD11	1:B:336:ILE:HG13	2.02	0.41
1:A:434:ARG:NH1	1:A:434:ARG:CG	2.82	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	162/201 (81%)	159 (98%)	3 (2%)	0	100	100
1	B	164/201 (82%)	160 (98%)	4 (2%)	0	100	100
All	All	326/402 (81%)	319 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/160 (82%)	131 (99%)	1 (1%)	85	80
1	B	136/160 (85%)	128 (94%)	8 (6%)	23	7
All	All	268/320 (84%)	259 (97%)	9 (3%)	45	22

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	432	ARG
1	B	253[A]	ARG
1	B	253[B]	ARG
1	B	266	ARG
1	B	330	ARG
1	B	378	LEU
1	B	380	LYS
1	B	420	ARG
1	B	434	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	346	ASN
1	A	358	GLN
1	A	419	ASN
1	A	425	GLN
1	B	357	GLN
1	B	358	GLN
1	B	419	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	1	-	4,4,4	0.23	0	6,6,6	0.29	0
2	SO4	B	2	-	4,4,4	0.29	0	6,6,6	0.46	0
2	SO4	B	3	-	4,4,4	0.15	0	6,6,6	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	1	-	-	0/0/0/0	0/0/0/0
2	SO4	B	2	-	-	0/0/0/0	0/0/0/0
2	SO4	B	3	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	163/201 (81%)	-0.03	12 (7%) 15 15	11, 20, 43, 60	0
1	B	163/201 (81%)	0.05	10 (6%) 22 22	11, 20, 46, 61	0
All	All	326/402 (81%)	0.01	22 (6%) 19 18	11, 20, 47, 61	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	253[A]	ARG	5.2
1	B	433	LEU	4.5
1	A	378	LEU	3.6
1	B	432	ARG	3.3
1	A	252	ASP	3.2
1	A	253	ARG	2.9
1	A	432	ARG	2.8
1	A	434	ARG	2.8
1	B	434	ARG	2.7
1	A	433	LEU	2.4
1	B	431	GLU	2.4
1	B	373	ASN	2.4
1	A	251	GLY	2.4
1	B	251	GLY	2.4
1	A	431	GLU	2.3
1	B	378	LEU	2.3
1	B	255	TYR	2.3
1	A	361	ASP	2.3
1	A	428	GLU	2.2
1	A	377	ASN	2.2
1	B	377	ASN	2.1
1	A	425	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	B	3	5/5	0.92	0.14	0.11	43,50,50,51	5
2	SO4	A	1	5/5	0.99	0.07	-0.93	24,27,28,28	0
2	SO4	B	2	5/5	0.99	0.07	-1.17	25,25,29,32	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.