



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 06:05 pm GMT

PDB ID : 4BKE  
Title : Recombinant human serum albumin with palmitic acid. Synthetic cationic antimicrobial peptides bind with their hydrophobic parts to drug site II of human serum albumin  
Authors : Sivertsen, A.; Isaksson, J.; Leiros, H.-K.S.; Svenson, J.; Svendsen, J.-S.; Brandsdal, B.-O.  
Deposited on : 2013-04-24  
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1522 (2.38-2.34)
Clashscore	112137	1626 (2.38-2.34)
Ramachandran outliers	110173	1605 (2.38-2.34)
Sidechain outliers	110143	1606 (2.38-2.34)
RSRZ outliers	101464	1528 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	609	<div> <div>4%</div> <div>77%</div> <div>16%</div> <div>• •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PLM	A	1001	-	-	-	X
2	PLM	A	1002	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PLM	A	1003	-	-	-	X
2	PLM	A	1004	-	-	-	X
2	PLM	A	1005	-	-	-	X
2	PLM	A	1007	-	-	-	X

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4793 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SERUM ALBUMIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	582	Total	C	N	O	S	183	2	0
			4648	2933	786	888	41			

- Molecule 2 is PALMITIC ACID (three-letter code: PLM) (formula:  $C_{16}H_{32}O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	11	2		
2	A	1	Total	C	O	0	0
			18	16	2		
2	A	1	Total	C	O	0	0
			17	15	2		
2	A	1	Total	C	O	0	0
			18	16	2		
2	A	1	Total	C	O	0	0
			17	15	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			18	16	2		
2	A	1	Total	C	O	2	0
			18	16	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	26	Total	O	0	0
			26	26		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: SERUM ALBUMIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	189.37Å 39.00Å 96.30Å 90.00° 105.48° 90.00°	Depositor
Resolution (Å)	20.00 – 2.35 19.95 – 2.35	Depositor EDS
% Data completeness (in resolution range)	96.6 (20.00-2.35) 96.6 (19.95-2.35)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.12 (at 2.35Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.198 , 0.254 0.195 , 0.249	Depositor DCC
$R_{free}$ test set	2006 reflections (7.76%)	DCC
Wilson B-factor (Å <sup>2</sup> )	57.7	Xtriage
Anisotropy	0.093	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 47.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4793	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.04% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PLM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	2.34	46/4738 (1.0%)	1.57	64/6390 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (46) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	396	GLU	CD-OE1	-67.34	0.51	1.25
1	A	244	GLU	CD-OE1	-44.52	0.76	1.25
1	A	396	GLU	CD-OE2	40.55	1.70	1.25
1	A	137	LYS	CD-CE	-37.42	0.57	1.51
1	A	396	GLU	CG-CD	-35.03	0.99	1.51
1	A	277	GLU	CG-CD	-34.80	0.99	1.51
1	A	205	LYS	CD-CE	-33.47	0.67	1.51
1	A	204	GLN	CB-CG	-32.34	0.65	1.52
1	A	170	GLN	CG-CD	-28.35	0.85	1.51
1	A	294	GLU	CG-CD	28.27	1.94	1.51
1	A	500	LYS	CE-NZ	-26.94	0.81	1.49
1	A	184	GLU	CD-OE1	26.65	1.54	1.25
1	A	41	LYS	CE-NZ	-26.19	0.83	1.49
1	A	6	GLU	CD-OE1	26.13	1.54	1.25
1	A	519	LYS	CG-CD	-25.88	0.64	1.52
1	A	372	LYS	CA-CB	-21.41	1.06	1.53
1	A	466	LYS	CD-CE	-21.34	0.97	1.51
1	A	244	GLU	CD-OE2	20.32	1.48	1.25

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	538	LYS	CE-NZ	-19.38	1.00	1.49
1	A	560	LYS	CB-CG	-18.71	1.02	1.52
1	A	184	GLU	CD-OE2	-18.54	1.05	1.25
1	A	389	LYS	CG-CD	-18.13	0.90	1.52
1	A	570	GLU	CA-CB	-16.86	1.16	1.53
1	A	444	LYS	CB-CG	16.29	1.96	1.52
1	A	459	GLN	CB-CG	-15.28	1.11	1.52
1	A	109	ASN	CB-CG	-15.27	1.16	1.51
1	A	240	LYS	CG-CD	-15.08	1.01	1.52
1	A	276	LYS	CB-CG	-14.87	1.12	1.52
1	A	536	LYS	CG-CD	-13.11	1.07	1.52
1	A	524	LYS	CG-CD	-11.37	1.13	1.52
1	A	313	LYS	CA-CB	-10.60	1.30	1.53
1	A	57	GLU	CB-CG	-10.08	1.33	1.52
1	A	351	LYS	CD-CE	-9.77	1.26	1.51
1	A	82	GLU	CB-CG	9.61	1.70	1.52
1	A	170	GLN	CB-CG	9.04	1.76	1.52
1	A	545	LYS	CE-NZ	-8.51	1.27	1.49
1	A	518	GLU	CB-CG	-8.41	1.36	1.52
1	A	111	ASN	CB-CG	-8.15	1.32	1.51
1	A	390	GLN	CB-CG	-7.97	1.31	1.52
1	A	402	LYS	CE-NZ	-7.93	1.29	1.49
1	A	466	LYS	CG-CD	-7.13	1.28	1.52
1	A	57	GLU	CG-CD	7.05	1.62	1.51
1	A	498	VAL	CB-CG1	-6.50	1.39	1.52
1	A	557	LYS	CB-CG	-5.39	1.38	1.52
1	A	311	GLU	CB-CG	5.37	1.62	1.52
1	A	3	HIS	CA-CB	5.13	1.65	1.53

All (64) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	244	GLU	OE1-CD-OE2	42.28	174.04	123.30
1	A	396	GLU	OE1-CD-OE2	39.66	170.90	123.30
1	A	519	LYS	CG-CD-CE	-27.54	29.29	111.90
1	A	6	GLU	OE1-CD-OE2	-22.72	96.04	123.30
1	A	277	GLU	CG-CD-OE2	-21.07	76.17	118.30
1	A	396	GLU	CG-CD-OE2	-20.84	76.61	118.30
1	A	519	LYS	CB-CG-CD	-20.71	57.75	111.60
1	A	244	GLU	CG-CD-OE2	-20.66	76.98	118.30
1	A	521	ARG	CB-CG-CD	-19.80	60.13	111.60
1	A	277	GLU	CG-CD-OE1	19.76	157.82	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	538	LYS	CD-CE-NZ	-18.46	69.25	111.70
1	A	277	GLU	CB-CG-CD	-18.22	65.02	114.20
1	A	536	LYS	CG-CD-CE	18.07	166.12	111.90
1	A	570	GLU	CB-CA-C	17.95	146.31	110.40
1	A	570	GLU	N-CA-CB	-16.94	80.10	110.60
1	A	317	LYS	CB-CG-CD	-16.73	68.11	111.60
1	A	444	LYS	CA-CB-CG	-16.61	76.86	113.40
1	A	109	ASN	CB-CG-OD1	-13.67	94.26	121.60
1	A	205	LYS	CG-CD-CE	13.55	152.54	111.90
1	A	402	LYS	CD-CE-NZ	13.47	142.67	111.70
1	A	389	LYS	CB-CG-CD	13.41	146.46	111.60
1	A	109	ASN	CA-CB-CG	-12.86	85.11	113.40
1	A	137	LYS	CD-CE-NZ	-12.52	82.92	111.70
1	A	536	LYS	CB-CG-CD	12.33	143.66	111.60
1	A	137	LYS	CG-CD-CE	-12.25	75.14	111.90
1	A	498	VAL	CG1-CB-CG2	12.11	130.28	110.90
1	A	317	LYS	CG-CD-CE	-11.73	76.70	111.90
1	A	3	HIS	CB-CA-C	-11.46	87.47	110.40
1	A	570	GLU	CA-CB-CG	-10.96	89.29	113.40
1	A	109	ASN	CB-CG-ND2	10.72	142.42	116.70
1	A	372	LYS	CB-CA-C	10.23	130.87	110.40
1	A	372	LYS	CA-CB-CG	9.96	135.32	113.40
1	A	111	ASN	N-CA-CB	9.58	127.85	110.60
1	A	184	GLU	CG-CD-OE2	9.39	137.09	118.30
1	A	372	LYS	N-CA-CB	-9.37	93.74	110.60
1	A	560	LYS	CB-CG-CD	-9.27	87.51	111.60
1	A	170	GLN	CA-CB-CG	-9.26	93.03	113.40
1	A	276	LYS	CB-CG-CD	-9.07	88.03	111.60
1	A	536	LYS	CD-CE-NZ	-8.87	91.30	111.70
1	A	481	LEU	N-CA-CB	-8.34	93.73	110.40
1	A	466	LYS	CD-CE-NZ	-8.30	92.60	111.70
1	A	276	LYS	CA-CB-CG	-8.19	95.37	113.40
1	A	294	GLU	CB-CG-CD	-8.01	92.59	114.20
1	A	378	LYS	CB-CG-CD	7.95	132.26	111.60
1	A	240	LYS	CB-CG-CD	7.62	131.41	111.60
1	A	524	LYS	CB-CG-CD	7.49	131.08	111.60
1	A	60	GLU	CA-CB-CG	7.41	129.69	113.40
1	A	244	GLU	CG-CD-OE1	-7.32	103.66	118.30
1	A	466	LYS	CG-CD-CE	-7.14	90.47	111.90
1	A	436	LYS	CB-CG-CD	6.93	129.62	111.60
1	A	396	GLU	CB-CG-CD	6.78	132.49	114.20
1	A	184	GLU	OE1-CD-OE2	-6.71	115.25	123.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	466	LYS	CB-CG-CD	6.63	128.84	111.60
1	A	500	LYS	CD-CE-NZ	6.38	126.38	111.70
1	A	390	GLN	CA-CB-CG	6.36	127.39	113.40
1	A	311	GLU	CA-CB-CG	-6.15	99.88	113.40
1	A	389	LYS	CG-CD-CE	6.12	130.24	111.90
1	A	481	LEU	CA-CB-CG	-5.66	102.29	115.30
1	A	351	LYS	CG-CD-CE	5.40	128.10	111.90
1	A	311	GLU	CB-CG-CD	-5.37	99.71	114.20
1	A	184	GLU	CG-CD-OE1	-5.34	107.62	118.30
1	A	556	GLU	CB-CG-CD	5.17	128.16	114.20
1	A	301	ASP	CB-CG-OD2	5.16	122.94	118.30
1	A	257	ARG	NE-CZ-NH2	-5.07	117.77	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	6	GLU	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4648	0	4561	35	0
2	A	119	0	194	3	0
3	A	26	0	0	4	0
All	All	4793	0	4755	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:29[B]:GLN:HG3	3:A:2003:HOH:O	1.87	0.73

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:383:GLU:HG3	1:A:384:PRO:HD3	1.75	0.68
1:A:562:ASP:OD1	1:A:562:ASP:N	2.25	0.67
1:A:241:VAL:HG22	1:A:256:ASP:HB3	1.77	0.65
1:A:272:SER:HB3	1:A:275:LEU:HD22	1.79	0.65
1:A:460:LEU:HD11	2:A:1004:PLM:HC1	1.77	0.64
1:A:108:ASP:HB3	1:A:148:TYR:CE2	2.39	0.58
1:A:519:LYS:O	1:A:523:ILE:HG12	2.07	0.54
1:A:216:VAL:CG2	1:A:235:VAL:HG21	2.38	0.53
1:A:408:LEU:HD13	1:A:427:SER:HB2	1.91	0.53
1:A:195:LYS:HG3	1:A:455:VAL:HG11	1.90	0.52
1:A:426:VAL:HG11	1:A:460:LEU:HD13	1.92	0.51
1:A:81:ARG:NH2	1:A:89:ASP:OD1	2.42	0.51
1:A:415:VAL:HG12	1:A:418:VAL:HG23	1.93	0.51
1:A:312:SER:HB3	1:A:315:VAL:HG23	1.92	0.51
1:A:7:VAL:HG22	3:A:2001:HOH:O	2.12	0.50
1:A:216:VAL:HG22	1:A:235:VAL:HG21	1.94	0.50
1:A:108:ASP:HB3	1:A:148:TYR:HE2	1.76	0.50
1:A:383:GLU:HG3	1:A:384:PRO:CD	2.42	0.49
1:A:408:LEU:HD13	1:A:427:SER:CB	2.42	0.49
1:A:101:CYS:O	1:A:105:HIS:HD2	1.96	0.49
1:A:576:VAL:HG12	1:A:580:GLN:HE21	1.78	0.48
1:A:507:PHE:CD1	2:A:1005:PLM:HB2	2.50	0.47
1:A:29[B]:GLN:HB2	1:A:147:PRO:HA	1.97	0.46
1:A:107:ASP:OD1	1:A:108:ASP:O	2.34	0.45
1:A:224:PRO:HD2	1:A:296:ASP:HB3	1.99	0.45
1:A:344:VAL:HG22	1:A:450:GLU:OE2	2.17	0.45
1:A:348:ARG:HG2	1:A:482:VAL:HG12	1.99	0.44
1:A:268:GLN:HG2	3:A:2015:HOH:O	2.17	0.44
1:A:119:GLU:HG3	3:A:2010:HOH:O	2.17	0.43
2:A:1003:PLM:H81	2:A:1004:PLM:H22	2.01	0.43
1:A:302:LEU:HA	1:A:303:PRO:HD3	1.88	0.42
1:A:307:ALA:HA	1:A:311:GLU:HB2	2.01	0.42
1:A:156:PHE:HZ	1:A:284:LEU:O	2.04	0.41
1:A:417:GLN:HB2	1:A:469:VAL:HG13	2.02	0.41
1:A:348:ARG:CG	1:A:482:VAL:HG12	2.50	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	582/609 (96%)	559 (96%)	21 (4%)	2 (0%)	44	53

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	561	ALA
1	A	322	ALA

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	511/533 (96%)	484 (95%)	27 (5%)	26	32

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	VAL
1	A	14	LEU
1	A	74	LEU
1	A	97	GLU
1	A	203	LEU
1	A	225	LYS
1	A	238	LEU
1	A	275	LEU
1	A	283	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	313	LYS
1	A	334	TYR
1	A	380	LEU
1	A	383	GLU
1	A	389	LYS
1	A	408	LEU
1	A	419	SER
1	A	430	LEU
1	A	451	ASP
1	A	457	LEU
1	A	496	THR
1	A	498	VAL
1	A	521	ARG
1	A	545	LYS
1	A	560	LYS
1	A	562	ASP
1	A	566	THR
1	A	573	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	105	HIS
1	A	242	HIS
1	A	267	ASN
1	A	318	ASN
1	A	397	GLN
1	A	580	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PLM	A	1001	-	9,12,17	0.31	0	8,12,17	0.49	0
2	PLM	A	1002	-	14,17,17	0.29	0	13,17,17	0.55	0
2	PLM	A	1003	-	13,16,17	0.24	0	12,16,17	0.65	0
2	PLM	A	1004	-	14,17,17	0.22	0	13,17,17	0.73	0
2	PLM	A	1005	-	13,16,17	0.24	0	12,16,17	0.59	0
2	PLM	A	1006	-	14,17,17	0.18	0	13,17,17	0.82	0
2	PLM	A	1007	-	14,17,17	3.60	1 (7%)	13,17,17	1.47	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLM	A	1001	-	-	0/8/10/15	0/0/0/0
2	PLM	A	1002	-	-	0/13/15/15	0/0/0/0
2	PLM	A	1003	-	-	0/12/14/15	0/0/0/0
2	PLM	A	1004	-	-	0/13/15/15	0/0/0/0
2	PLM	A	1005	-	-	0/12/14/15	0/0/0/0
2	PLM	A	1006	-	-	0/13/15/15	0/0/0/0
2	PLM	A	1007	-	-	0/13/15/15	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1007	PLM	CF-CE	13.44	2.47	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1007	PLM	CF-CE-CD	-4.85	67.97	115.40

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1003	PLM	1	0
2	A	1004	PLM	2	0
2	A	1005	PLM	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	582/609 (95%)	0.15	25 (4%)	36 48	37, 58, 82, 94	75 (12%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	HIS	4.4
1	A	566	THR	4.3
1	A	176	ALA	3.9
1	A	561	ALA	3.0
1	A	307	ALA	2.8
1	A	330	PHE	2.6
1	A	528	ALA	2.6
1	A	94	GLN	2.5
1	A	309	PHE	2.5
1	A	515	THR	2.5
1	A	371	ALA	2.3
1	A	70	PHE	2.3
1	A	148	TYR	2.2
1	A	30	TYR	2.2
1	A	559	CYS	2.2
1	A	26	ALA	2.2
1	A	546	ALA	2.1
1	A	532	LEU	2.1
1	A	110	PRO	2.1
1	A	565	GLU	2.1
1	A	29[A]	GLN	2.1
1	A	331	LEU	2.1
1	A	457	LEU	2.0
1	A	108	ASP	2.0
1	A	74	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PLM	A	1001	13/18	0.70	0.35	8.74	72,74,77,77	0
2	PLM	A	1007	18/18	0.63	0.50	6.50	73,79,88,89	2
2	PLM	A	1003	17/18	0.92	0.32	5.11	49,56,59,59	0
2	PLM	A	1002	18/18	0.88	0.27	2.60	55,59,65,65	0
2	PLM	A	1005	17/18	0.85	0.31	2.49	59,62,68,68	0
2	PLM	A	1004	18/18	0.88	0.28	2.43	56,59,61,61	0
2	PLM	A	1006	18/18	0.84	0.24	1.72	56,59,72,72	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.