



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 09:28 pm GMT

PDB ID : 2BLC  
Title : SP21 DOUBLE MUTANT P. VIVAX DIHYDROFOLATE REDUCTASE IN COMPLEX WITH DES-CHLOROPYRIMETHAMINE  
Authors : Kongsaree, P.; Khongsuk, P.; Leartsakulpanich, U.; Chitnumsub, P.; Tarnchompoo, B.; Walkinshaw, M.D.; Yuthavong, Y.  
Deposited on : 2005-03-03  
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1062 (2.26-2.26)
Clashscore	112137	1178 (2.26-2.26)
Ramachandran outliers	110173	1145 (2.26-2.26)
Sidechain outliers	110143	1146 (2.26-2.26)
RSRZ outliers	101464	1066 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	238	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2045 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DIHYDROFOLATE REDUCTASE-THYMIDYLATE SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	215	1760	1137	291	322	10	0	0	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	3	ASN	ASP	CONFLICT	UNP Q5U9H1
A	213	GLU	GLN	CONFLICT	UNP Q5U9H1

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula:  $C_{21}H_{30}N_7O_{17}P_3$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	48	21	7	17	3	0	0

- Molecule 3 is 6-ETHYL-5-PHENYLPYRIMIDINE-2,4-DIAMINE (three-letter code: CP7) (formula:  $C_{12}H_{14}N_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			16	12	4		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula:  $C_6H_{13}NO_4S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	209	Total 209	O 209	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: DIHYDROFOLATE REDUCTASE-THYMIDYLATE SYNTHASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	132.00Å 56.31Å 45.67Å 90.00° 107.50° 90.00°	Depositor
Resolution (Å)	35.45 – 2.25 35.45 – 2.25	Depositor EDS
% Data completeness (in resolution range)	97.3 (35.45-2.25) 97.4 (35.45-2.25)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.42 (at 2.24Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.215 , 0.280 0.221 , 0.285	Depositor DCC
$R_{free}$ test set	1515 reflections (10.15%)	DCC
Wilson B-factor (Å <sup>2</sup> )	46.3	Xtriage
Anisotropy	0.436	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 58.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.019 for -h-2*k,l	Xtriage
$F_o, F_c$ correlation	0.86	EDS
Total number of atoms	2045	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.44% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, MES, CP7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.62	0/1798	0.73	0/2424

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1760	0	1778	89	10
2	A	48	0	26	6	0
3	A	16	0	14	2	0
4	A	12	0	13	1	0
5	A	209	0	0	29	3
All	All	2045	0	1831	90	12

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:224:ASN:O	5:A:2195:HOH:O	1.77	1.02
1:A:213:GLU:HB3	5:A:2179:HOH:O	1.70	0.90
1:A:220:SER:N	5:A:2191:HOH:O	2.05	0.88
1:A:106:LYS:N	1:A:106:LYS:HE2	1.90	0.86
1:A:220:SER:CA	5:A:2191:HOH:O	2.26	0.82
1:A:157:ASP:HA	1:A:160:LEU:HD12	1.61	0.81
1:A:220:SER:HA	5:A:2191:HOH:O	1.80	0.80
1:A:141:THR:HG22	1:A:144:ASP:CG	2.04	0.77
1:A:224:ASN:HD21	1:A:227:GLY:H	1.32	0.76
1:A:27:LYS:HB2	5:A:2030:HOH:O	1.85	0.76
1:A:219:VAL:C	5:A:2191:HOH:O	2.24	0.74
1:A:174:GLY:HA3	2:A:1239:NDP:H5N	1.71	0.72
1:A:73:LYS:HE2	1:A:77:GLU:OE1	1.90	0.71
1:A:224:ASN:ND2	1:A:227:GLY:H	1.88	0.71
1:A:224:ASN:HD21	1:A:227:GLY:N	1.89	0.70
1:A:141:THR:HG22	1:A:144:ASP:OD1	1.92	0.70
1:A:2:GLU:HA	5:A:2003:HOH:O	1.92	0.69
1:A:201:PRO:HD2	5:A:2169:HOH:O	1.94	0.66
1:A:106:LYS:NZ	5:A:2096:HOH:O	1.99	0.66
1:A:3:ASN:HD21	1:A:237:LYS:NZ	1.95	0.65
1:A:121:ILE:O	1:A:126:LYS:HD3	1.96	0.65
1:A:153:ASP:HB3	5:A:2116:HOH:O	1.95	0.64
1:A:66:GLU:HG2	5:A:2077:HOH:O	1.97	0.64
1:A:134:VAL:HG22	1:A:150:PHE:HB2	1.77	0.64
1:A:155:ILE:HG13	1:A:181:GLU:HG2	1.79	0.64
1:A:218:SER:OG	5:A:2189:HOH:O	2.15	0.63
1:A:78:ARG:HH11	1:A:78:ARG:HG3	1.65	0.62
1:A:234:VAL:HG21	5:A:2061:HOH:O	2.01	0.61
1:A:219:VAL:HG13	1:A:233:LEU:CD2	2.31	0.60
1:A:224:ASN:HD21	1:A:227:GLY:CA	2.14	0.60
1:A:195:ARG:HB2	1:A:231:ASP:OD2	2.01	0.59
1:A:139:THR:O	5:A:2119:HOH:O	2.17	0.59
1:A:3:ASN:HD21	1:A:237:LYS:HZ3	1.49	0.59
1:A:39:LEU:O	2:A:1239:NDP:H2N	2.03	0.59
1:A:133:ASN:HD22	1:A:133:ASN:N	2.00	0.58
1:A:34:ARG:NH1	1:A:211:GLU:OE1	2.37	0.57
1:A:148:LYS:HE3	5:A:2055:HOH:O	2.04	0.57
1:A:27:LYS:CB	5:A:2030:HOH:O	2.47	0.56
1:A:106:LYS:N	1:A:106:LYS:CE	2.66	0.55
1:A:130:ASN:ND2	5:A:2108:HOH:O	2.40	0.55
1:A:79:TYR:HA	1:A:82:MET:CE	2.36	0.55
1:A:224:ASN:HD22	1:A:224:ASN:C	2.10	0.55

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125:TYR:HE2	5:A:2103:HOH:O	1.89	0.54
1:A:121:ILE:HG12	4:A:1241:MES:H61	1.91	0.53
1:A:58:ARG:NH1	5:A:2065:HOH:O	2.40	0.53
1:A:8:PHE:HB3	1:A:167:TYR:OH	2.09	0.53
1:A:230:LEU:HD23	1:A:230:LEU:N	2.25	0.52
1:A:59:SER:HB3	5:A:2068:HOH:O	2.09	0.52
1:A:174:GLY:HA3	2:A:1239:NDP:C5N	2.39	0.51
1:A:213:GLU:O	1:A:238:VAL:HG23	2.10	0.51
1:A:208:GLU:HB3	5:A:2176:HOH:O	2.12	0.50
1:A:133:ASN:HB2	1:A:149:VAL:HG12	1.92	0.50
1:A:129:PRO:O	1:A:130:ASN:HB2	2.12	0.50
1:A:28:ASN:O	1:A:30:PRO:HD3	2.12	0.50
2:A:1239:NDP:H42N	3:A:1240:CP7:C12	2.42	0.49
1:A:234:VAL:CG2	5:A:2061:HOH:O	2.59	0.49
1:A:176:ALA:HB3	2:A:1239:NDP:O2N	2.12	0.48
1:A:79:TYR:HA	1:A:82:MET:HE3	1.95	0.48
1:A:210:ASP:O	1:A:213:GLU:HG2	2.13	0.48
1:A:180:ARG:NH2	5:A:2147:HOH:O	2.46	0.47
1:A:213:GLU:HG3	1:A:214:PHE:CD1	2.49	0.47
1:A:223:TYR:O	1:A:229:THR:HA	2.14	0.47
1:A:113:MET:HA	1:A:174:GLY:O	2.15	0.47
1:A:168:TYR:CD2	1:A:169:LYS:HD3	2.49	0.47
1:A:78:ARG:CG	1:A:78:ARG:HH11	2.29	0.46
1:A:123:LYS:C	1:A:125:TYR:H	2.18	0.46
1:A:2:GLU:N	5:A:2004:HOH:O	2.47	0.46
1:A:53:ASP:OD2	3:A:1240:CP7:N6	2.49	0.46
1:A:70:GLU:HG3	5:A:2083:HOH:O	2.16	0.46
1:A:118:TRP:CH2	1:A:126:LYS:HB3	2.51	0.45
1:A:79:TYR:HA	1:A:82:MET:HE2	1.99	0.45
1:A:54:MET:HE2	5:A:2054:HOH:O	2.15	0.45
1:A:218:SER:HB2	1:A:234:VAL:HB	1.99	0.44
1:A:142:LYS:HD3	1:A:142:LYS:HA	1.70	0.44
1:A:155:ILE:HG13	1:A:181:GLU:CG	2.47	0.44
1:A:69:TYR:OH	1:A:166:LYS:HG3	2.18	0.43
1:A:8:PHE:HB3	1:A:167:TYR:HH	1.84	0.43
1:A:124:GLN:HG2	1:A:125:TYR:CD1	2.54	0.43
1:A:115:ARG:HD3	1:A:119:GLU:OE1	2.18	0.42
1:A:138:LYS:HB2	1:A:138:LYS:HE3	1.82	0.42
1:A:204:VAL:HG21	2:A:1239:NDP:H4D	2.02	0.42
1:A:69:TYR:O	1:A:72:LEU:N	2.51	0.42
1:A:40:GLY:HA2	1:A:46:PRO:HD3	2.01	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:ASN:HA	1:A:203:ASP:OD2	2.20	0.41
1:A:224:ASN:HD22	1:A:225:SER:N	2.17	0.41
1:A:28:ASN:OD1	5:A:2031:HOH:O	2.22	0.41
1:A:81:ARG:O	1:A:83:GLU:N	2.54	0.41
1:A:34:ARG:HB2	5:A:2174:HOH:O	2.20	0.41
1:A:32:SER:HA	1:A:33:PRO:HD3	1.96	0.40
1:A:133:ASN:ND2	1:A:133:ASN:N	2.68	0.40

All (12) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:29:GLU:OE2	1:A:221:GLU:N[4_545]	1.07	1.13
1:A:32:SER:OG	1:A:223:TYR:OH[4_545]	1.13	1.07
1:A:32:SER:CB	1:A:223:TYR:OH[4_545]	1.81	0.39
1:A:27:LYS:NZ	1:A:146:LYS:O[4_546]	1.82	0.38
1:A:32:SER:OG	1:A:223:TYR:CZ[4_545]	1.83	0.37
5:A:2104:HOH:O	5:A:2118:HOH:O[4_556]	1.86	0.34
1:A:26:THR:O	1:A:219:VAL:O[4_545]	1.89	0.31
5:A:2002:HOH:O	5:A:2002:HOH:O[2_555]	1.90	0.30
1:A:29:GLU:OE2	1:A:221:GLU:CA[4_545]	2.06	0.14
1:A:205:PHE:CB	5:A:2193:HOH:O[4_545]	2.10	0.10
1:A:29:GLU:OE2	1:A:220:SER:C[4_545]	2.11	0.09
1:A:32:SER:OG	1:A:223:TYR:CE2[4_545]	2.13	0.07

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	211/238 (89%)	191 (90%)	17 (8%)	3 (1%)	<b>13</b> <b>8</b>

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	49	CYS
1	A	82	MET
1	A	124	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/215 (93%)	186 (94%)	13 (6%)	20	19

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	ARG
1	A	78	ARG
1	A	106	LYS
1	A	133	ASN
1	A	138	LYS
1	A	142	LYS
1	A	156	ASP
1	A	157	ASP
1	A	169	LYS
1	A	184	SER
1	A	197	ASN
1	A	224	ASN
1	A	237	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	41	ASN
1	A	177	GLN
1	A	197	ASN
1	A	224	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NDP	A	1239	-	43,52,52	1.88	11 (25%)	49,80,80	1.83	13 (26%)
3	CP7	A	1240	-	17,17,17	2.24	7 (41%)	20,23,23	1.35	1 (5%)
4	MES	A	1241	-	12,12,12	1.67	4 (33%)	14,16,16	1.55	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	1239	-	-	0/30/77/77	0/5/5/5
3	CP7	A	1240	-	-	0/6/6/6	0/2/2/2
4	MES	A	1241	-	-	0/6/14/14	0/1/1/1

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1239	NDP	C4N-C5N	-4.71	1.38	1.49
2	A	1239	NDP	P2B-O2B	-3.33	1.53	1.59
2	A	1239	NDP	C3B-C2B	-3.02	1.46	1.53
2	A	1239	NDP	C8A-N7A	-2.44	1.30	1.34
2	A	1239	NDP	PA-O1A	-2.35	1.42	1.50
4	A	1241	MES	C5-N4	2.00	1.52	1.47
2	A	1239	NDP	C2D-C1D	2.02	1.59	1.53
2	A	1239	NDP	C5D-C4D	2.10	1.58	1.51
3	A	1240	CP7	C11-C12	2.11	1.42	1.38
4	A	1241	MES	C7-N4	2.13	1.52	1.47
3	A	1240	CP7	C9-C8	2.19	1.43	1.38
3	A	1240	CP7	C8-C7	2.21	1.44	1.39
2	A	1239	NDP	O4D-C1D	2.31	1.47	1.42
4	A	1241	MES	C8-S	2.43	1.81	1.77
4	A	1241	MES	C3-N4	3.25	1.55	1.47
3	A	1240	CP7	C4-C3	3.26	1.47	1.43
2	A	1239	NDP	C6N-C5N	3.31	1.39	1.33
3	A	1240	CP7	C2-N14	3.33	1.40	1.34
2	A	1239	NDP	O4B-C1B	3.54	1.46	1.41
3	A	1240	CP7	C3-N1	4.05	1.41	1.35
2	A	1239	NDP	C2N-C3N	4.12	1.46	1.34
3	A	1240	CP7	C5-N6	4.45	1.42	1.34

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1239	NDP	C4B-O4B-C1B	-5.79	103.61	109.77
2	A	1239	NDP	C3N-C2N-N1N	-4.30	116.84	123.08
2	A	1239	NDP	N3A-C2A-N1A	-4.17	125.22	128.86
2	A	1239	NDP	C1D-N1N-C2N	-3.86	114.53	121.09
2	A	1239	NDP	C3B-C2B-C1B	-2.68	97.52	102.75
4	A	1241	MES	O3S-S-C8	-2.65	102.80	106.06
2	A	1239	NDP	C3D-C2D-C1D	-2.60	96.44	101.43
2	A	1239	NDP	O7N-C7N-N7N	-2.32	117.28	122.92
2	A	1239	NDP	O5B-C5B-C4B	-2.11	101.50	109.00
2	A	1239	NDP	O2A-PA-O1A	2.10	123.17	112.28
2	A	1239	NDP	O4B-C1B-C2B	2.14	110.34	106.59
2	A	1239	NDP	C2D-C3D-C4D	2.22	106.94	102.62
2	A	1239	NDP	O3B-C3B-C2B	2.40	118.00	111.18
3	A	1240	CP7	C4-C3-N1	2.70	124.01	122.49
2	A	1239	NDP	O3B-C3B-C4B	2.73	119.07	111.09
4	A	1241	MES	O1S-S-C8	4.52	110.67	106.79

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1239	NDP	6	0
3	A	1240	CP7	2	0
4	A	1241	MES	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	215/238 (90%)	1.88	87 (40%) <b>0</b> <b>0</b>	27, 47, 62, 73	0

All (87) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	200	TYR	7.2
1	A	232	PHE	6.0
1	A	31	PHE	5.4
1	A	26	THR	5.3
1	A	22	THR	5.0
1	A	202	CYS	5.0
1	A	107	LEU	5.0
1	A	49	CYS	5.0
1	A	228	THR	4.8
1	A	74	TRP	4.8
1	A	19	VAL	4.7
1	A	38	GLY	4.7
1	A	227	GLY	4.6
1	A	23	SER	4.2
1	A	225	SER	4.2
1	A	67	SER	3.9
1	A	125	TYR	3.9
1	A	21	PRO	3.8
1	A	196	ILE	3.7
1	A	209	PHE	3.6
1	A	204	VAL	3.5
1	A	219	VAL	3.5
1	A	59	SER	3.5
1	A	166	LYS	3.4
1	A	214	PHE	3.2
1	A	197	ASN	3.2
1	A	33	PRO	3.2

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	223	TYR	3.1
1	A	222	VAL	3.1
1	A	66	GLU	3.1
1	A	52	VAL	3.1
1	A	27	LYS	3.0
1	A	20	ALA	3.0
1	A	193	PHE	2.9
1	A	178	VAL	2.9
1	A	106	LYS	2.9
1	A	199	ALA	2.9
1	A	24	GLU	2.8
1	A	139	THR	2.8
1	A	36	PHE	2.8
1	A	28	ASN	2.8
1	A	198	GLY	2.7
1	A	171	PHE	2.7
1	A	210	ASP	2.7
1	A	206	PHE	2.7
1	A	44	THR	2.7
1	A	224	ASN	2.6
1	A	34	ARG	2.6
1	A	18	LYS	2.6
1	A	205	PHE	2.6
1	A	47	TRP	2.6
1	A	57	PHE	2.6
1	A	63	TYR	2.6
1	A	234	VAL	2.5
1	A	235	TYR	2.5
1	A	188	ILE	2.5
1	A	108	GLN	2.5
1	A	17	CYS	2.5
1	A	30	PRO	2.5
1	A	29	GLU	2.5
1	A	220	SER	2.4
1	A	195	ARG	2.4
1	A	50	ASN	2.4
1	A	15	ALA	2.4
1	A	175	GLY	2.4
1	A	176	ALA	2.4
1	A	45	LEU	2.3
1	A	212	SER	2.3
1	A	151	ILE	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	146	LYS	2.2
1	A	16	CYS	2.2
1	A	168	TYR	2.2
1	A	116	SER	2.2
1	A	217	THR	2.2
1	A	32	SER	2.2
1	A	70	GLU	2.2
1	A	113	MET	2.2
1	A	140	LEU	2.1
1	A	43	GLY	2.1
1	A	182	CYS	2.1
1	A	172	ILE	2.1
1	A	130	ASN	2.1
1	A	233	LEU	2.1
1	A	12	ALA	2.1
1	A	60	VAL	2.0
1	A	192	TYR	2.0
1	A	58	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	CP7	A	1240	16/16	0.80	0.29	1.94	26,34,42,45	0
2	NDP	A	1239	48/48	0.85	0.24	0.03	34,47,58,61	0
4	MES	A	1241	12/12	0.84	0.24	-0.27	45,54,62,62	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.