



Full wwPDB X-ray Structure Validation Report

Feb 13, 2017 – 06:31 pm GMT

PDB ID : 3BLY
Title : Pyranose 2-oxidase from Trametes multicolor, E542K/L537W
Authors : Tan, T.C.; Divne, C.
Deposited on : 2007-12-11
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	5047 (1.90-1.90)
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	623	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4943 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pyranose oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	573	4524	2859	776	865	24	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	537	TRP	LEU	ENGINEERED	UNP Q7ZA32
A	542	LYS	GLU	ENGINEERED	UNP Q7ZA32


- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES)

(formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	12	6	1	4	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	354	354	354	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pyranose oxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	103.43Å 103.43Å 118.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.40 – 1.90 47.40 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.40-1.90) 100.0 (47.40-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.92 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.149 , 0.181 0.156 , 0.192	Depositor DCC
R_{free} test set	504 reflections (0.99%)	DCC
Wilson B-factor (Å ²)	18.4	Xtrriage
Anisotropy	0.042	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 61.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4943	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.02	3/4642 (0.1%)	0.95	14/6312 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	63	ARG	CG-CD	-8.31	1.31	1.51
1	A	285	ARG	CB-CG	-5.50	1.37	1.52
1	A	294	GLU	CG-CD	5.40	1.60	1.51

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	218	ARG	NE-CZ-NH1	11.91	126.26	120.30
1	A	218	ARG	NE-CZ-NH2	-10.22	115.19	120.30
1	A	417	MET	CG-SD-CE	-8.49	86.61	100.20
1	A	81	ASP	CB-CG-OD1	-8.37	110.76	118.30
1	A	235	ASP	CB-CG-OD2	-7.36	111.67	118.30
1	A	211	ASP	CB-CG-OD1	6.75	124.37	118.30
1	A	81	ASP	CB-CG-OD2	6.11	123.80	118.30
1	A	63	ARG	NE-CZ-NH2	-6.06	117.27	120.30
1	A	63	ARG	NE-CZ-NH1	5.56	123.08	120.30
1	A	235	ASP	CB-CG-OD1	5.45	123.21	118.30
1	A	124	ASP	CB-CG-OD2	-5.28	113.55	118.30
1	A	227	THR	CA-CB-CG2	-5.25	105.05	112.40
1	A	265	ARG	NE-CZ-NH1	5.18	122.89	120.30
1	A	493	ASP	CB-CG-OD1	5.05	122.85	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4524	0	4370	34	1
2	A	53	0	30	0	0
3	A	12	0	12	0	0
4	A	354	0	0	5	0
All	All	4943	0	4412	34	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:393:VAL:H	1:A:417:MET:HE2	1.05	1.21
1:A:341:ASN:HD21	1:A:343:ALA:HB3	1.24	0.98
1:A:110:GLN:HE21	1:A:167:HIS:HD1	1.04	0.98
1:A:393:VAL:N	1:A:417:MET:HE2	1.87	0.90
1:A:393:VAL:H	1:A:417:MET:CE	1.88	0.84
1:A:341:ASN:ND2	1:A:343:ALA:HB3	1.93	0.82
1:A:450:HIS:HD2	4:A:849:HOH:O	1.75	0.69
1:A:97:GLN:HG3	1:A:250:PHE:CE2	2.28	0.69
1:A:81:ASP:C	1:A:81:ASP:OD1	2.31	0.65
1:A:82:SER:O	4:A:1019:HOH:O	2.15	0.65
1:A:218:ARG:HG3	1:A:430:ASP:OD2	2.02	0.60
1:A:404:HIS:HE1	4:A:929:HOH:O	1.87	0.58
1:A:110:GLN:NE2	1:A:167:HIS:HD1	1.89	0.56
1:A:342:PRO:O	1:A:343:ALA:C	2.44	0.56
1:A:450:HIS:CE1	1:A:472:ARG:HH11	2.26	0.54
1:A:100:ILE:HG23	1:A:101:ASP:N	2.21	0.54
1:A:46:LYS:N	4:A:882:HOH:O	2.40	0.54
1:A:459:VAL:CG1	1:A:460:GLN:N	2.71	0.54
1:A:299:HIS:CE1	1:A:310:GLU:OE1	2.61	0.54
1:A:618:PHE:C	1:A:618:PHE:CD1	2.82	0.52
1:A:341:ASN:HD22	1:A:341:ASN:C	2.12	0.52
1:A:100:ILE:CG2	1:A:101:ASP:N	2.76	0.49
1:A:450:HIS:HE1	1:A:472:ARG:HH11	1.61	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:341:ASN:HD21	1:A:343:ALA:CB	2.12	0.47
1:A:216:SER:HB3	1:A:219:HIS:HB3	1.96	0.46
1:A:50:VAL:HG13	1:A:313:ALA:HB2	1.97	0.46
1:A:218:ARG:HD2	4:A:842:HOH:O	2.15	0.46
1:A:63:ARG:HG3	1:A:259:VAL:O	2.17	0.45
1:A:284:GLU:O	1:A:328:LEU:HD12	2.19	0.43
1:A:47:TYR:O	1:A:313:ALA:HA	2.19	0.42
1:A:81:ASP:N	1:A:81:ASP:OD1	2.51	0.42
1:A:91:LYS:HE3	1:A:253:TRP:CD1	2.55	0.42
1:A:570:SER:HB3	1:A:580:LEU:O	2.20	0.42
1:A:158:THR:HG22	1:A:160:VAL:HG22	2.02	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:LEU:CD2	1:A:121:LEU:CD2[8_666]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	571/623 (92%)	556 (97%)	15 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	501/542 (92%)	486 (97%)	15 (3%)	46 37

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	ARG
1	A	168	TRP
1	A	206	PHE
1	A	231	LYS
1	A	341	ASN
1	A	348	LEU
1	A	385	THR
1	A	413	LYS
1	A	450	HIS
1	A	459	VAL
1	A	490	LYS
1	A	496	ASN
1	A	505	ARG
1	A	516	GLU
1	A	618	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	299	HIS
1	A	341	ASN
1	A	404	HIS
1	A	450	HIS
1	A	461	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FAD	A	801	1	51,58,58	1.19	6 (11%)	54,89,89	3.26	18 (33%)
3	MES	A	802	-	12,12,12	1.97	1 (8%)	14,16,16	3.69	6 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	801	1	-	0/28/50/50	0/6/6/6
3	MES	A	802	-	-	0/6/14/14	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	802	MES	C8-S	-6.63	1.67	1.77
2	A	801	FAD	C3B-C4B	-2.33	1.46	1.53
2	A	801	FAD	C2B-C1B	-2.08	1.50	1.53
2	A	801	FAD	C7M-C7	2.11	1.55	1.51
2	A	801	FAD	C6-C5X	2.43	1.45	1.41
2	A	801	FAD	C4X-N5	2.63	1.37	1.33
2	A	801	FAD	C4-C4X	2.96	1.47	1.41

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	MES	O1S-S-C8	-10.89	97.44	106.79
2	A	801	FAD	N3A-C2A-N1A	-9.78	120.34	128.86
2	A	801	FAD	C4X-C4-N3	-9.27	110.29	123.48
3	A	802	MES	O2S-S-C8	-2.95	104.25	106.79
2	A	801	FAD	C4B-O4B-C1B	-2.78	106.81	109.77
2	A	801	FAD	C9A-C5X-N5	-2.55	118.44	122.24
2	A	801	FAD	O5'-P-O1P	-2.28	100.06	109.25
3	A	802	MES	O3S-S-C8	-2.26	103.28	106.06
2	A	801	FAD	C4X-C10-N10	-2.25	118.95	120.52
2	A	801	FAD	C8M-C8-C9	-2.08	115.13	120.34
2	A	801	FAD	C7-C6-C5X	-2.05	117.92	121.08
2	A	801	FAD	O2A-PA-O1A	2.22	123.77	112.28
2	A	801	FAD	O2B-C2B-C1B	2.43	119.22	111.61
2	A	801	FAD	C2A-N1A-C6A	2.47	123.09	118.77
2	A	801	FAD	C4-C4X-C10	2.52	122.00	119.96
2	A	801	FAD	O2B-C2B-C3B	2.63	120.25	111.83
2	A	801	FAD	O3B-C3B-C4B	2.85	119.41	111.09
3	A	802	MES	O3S-S-O1S	2.94	118.11	111.37
2	A	801	FAD	O4B-C4B-C3B	2.94	111.02	105.17
2	A	801	FAD	C1'-N10-C9A	2.97	121.06	118.35
3	A	802	MES	C7-N4-C5	4.24	122.13	111.26
3	A	802	MES	C5-N4-C3	4.55	119.17	108.87
2	A	801	FAD	C4X-N5-C5X	4.91	121.95	116.76
2	A	801	FAD	C4-N3-C2	15.26	128.50	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	573/623 (91%)	-0.37	15 (2%) 56 60	6, 12, 22, 33	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	343	ALA	8.8
1	A	618	PHE	5.4
1	A	342	PRO	5.1
1	A	341	ASN	4.6
1	A	344	ASN	4.6
1	A	268	THR	3.7
1	A	269	ASP	3.3
1	A	385	THR	3.1
1	A	388	GLU	2.7
1	A	272	GLU	2.5
1	A	340	PRO	2.4
1	A	270	ALA	2.4
1	A	389	LEU	2.4
1	A	271	PRO	2.2
1	A	617	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	FAD	A	801	53/53	0.99	0.06	-0.74	11,14,17,20	0
3	MES	A	802	12/12	0.99	0.06	-0.96	17,18,20,21	0

6.5 Other polymers

There are no such residues in this entry.