



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 09:16 am GMT

PDB ID : 4BM7
Title : Crystal Structure of IgG Fc F241A mutant with native glycosylation
Authors : Yu, X.; Baruah, K.; Harvey, D.J.; Vasiljevic, S.; Alonzi, D.S.; Song, B.; Higgins, M.K.; Bowden, T.A.; Crispin, M.; Scanlan, C.N.
Deposited on : 2013-05-07
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2004 (1.96-1.96)
Clashscore	112137	2136 (1.96-1.96)
Ramachandran outliers	110173	2117 (1.96-1.96)
Sidechain outliers	110143	2117 (1.96-1.96)
RSRZ outliers	101464	2018 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	233	<div> <div>11%</div> <div> <div></div> <div>85%</div> <div>•</div> <div>11%</div> </div> </div>
1	B	233	<div> <div>14%</div> <div> <div></div> <div>73%</div> <div>• •</div> <div>21%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	1448	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3543 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IG GAMMA-1 CHAIN C REGION.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	207	Total	C	N	O	S	0	2	0
			1666	1060	280	318	8			
1	B	183	Total	C	N	O	S	0	5	0
			1494	952	252	282	8			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	222	GLY	-	EXPRESSION TAG	UNP P01857
A	447	THR	-	EXPRESSION TAG	UNP P01857
A	448	LYS	-	EXPRESSION TAG	UNP P01857
A	449	HIS	-	EXPRESSION TAG	UNP P01857
A	450	HIS	-	EXPRESSION TAG	UNP P01857
A	451	HIS	-	EXPRESSION TAG	UNP P01857
A	452	HIS	-	EXPRESSION TAG	UNP P01857
A	453	HIS	-	EXPRESSION TAG	UNP P01857
A	454	HIS	-	EXPRESSION TAG	UNP P01857
A	241	ALA	PHE	ENGINEERED MUTATION	UNP P01857
A	356	GLU	ASP	VARIANT	UNP P01857
A	358	MET	LEU	VARIANT	UNP P01857
B	222	GLY	-	EXPRESSION TAG	UNP P01857
B	447	THR	-	EXPRESSION TAG	UNP P01857
B	448	LYS	-	EXPRESSION TAG	UNP P01857
B	449	HIS	-	EXPRESSION TAG	UNP P01857
B	450	HIS	-	EXPRESSION TAG	UNP P01857
B	451	HIS	-	EXPRESSION TAG	UNP P01857
B	452	HIS	-	EXPRESSION TAG	UNP P01857
B	453	HIS	-	EXPRESSION TAG	UNP P01857
B	454	HIS	-	EXPRESSION TAG	UNP P01857
B	241	ALA	PHE	ENGINEERED MUTATION	UNP P01857
B	356	GLU	ASP	VARIANT	UNP P01857
B	358	MET	LEU	VARIANT	UNP P01857

- Molecule 2 is a polymer of unknown type called SUGAR (7-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	7	Total	C	N	O	0	0
			85	48	3	34		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	169	Total	O	0	0
			169	169		
4	B	128	Total	O	0	0
			128	128		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: IG GAMMA-1 CHAIN C REGION



• Molecule 1: IG GAMMA-1 CHAIN C REGION



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.87Å 73.38Å 135.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.92 – 1.95 49.87 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.6 (49.92-1.95) 98.7 (49.87-1.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.198 , 0.234 0.205 , 0.244	Depositor DCC
R_{free} test set	1831 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	28.2	Xtriage
Anisotropy	0.314	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 58.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3543	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.27 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.3534e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, FUL, BMA, NAG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/1714	0.67	3/2334 (0.1%)
1	B	0.48	0/1543	0.64	0/2093
All	All	0.48	0/3257	0.65	3/4427 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	255	ARG	NE-CZ-NH1	5.74	123.17	120.30
1	A	255	ARG	NE-CZ-NH2	-5.67	117.47	120.30
1	A	441	LEU	CA-CB-CG	5.13	127.10	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1666	0	1641	7	0
1	B	1494	0	1482	9	0
2	A	85	0	73	0	0
3	A	1	0	0	0	0
4	A	169	0	0	3	0
4	B	128	0	0	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3543	0	3196	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:430:GLU:OE1	4:A:2149:HOH:O	2.01	0.78
1:B:311[A]:GLN:OE1	4:B:2025:HOH:O	2.14	0.65
1:A:256:THR:HG23	4:A:2014:HOH:O	2.05	0.56
1:B:344:ARG:NH1	4:B:2040:HOH:O	2.31	0.55
1:A:387:PRO:HB3	4:A:2106:HOH:O	2.11	0.49
1:B:263:VAL:O	1:B:301:ARG:HB2	2.18	0.44
1:B:256:THR:HG23	4:B:2013:HOH:O	2.17	0.44
1:B:343:PRO:HA	1:B:373:TYR:O	2.19	0.43
1:B:311[A]:GLN:HG2	4:B:2042:HOH:O	2.19	0.43
1:A:406:LEU:HD12	1:A:406:LEU:C	2.39	0.43
1:A:292:ARG:HG2	1:A:302:VAL:HG22	2.01	0.43
1:B:406:LEU:HD12	1:B:406:LEU:C	2.39	0.42
1:B:439:LYS:HE2	4:B:2048:HOH:O	2.20	0.42
1:B:293:GLU:HG2	1:B:301:ARG:O	2.20	0.41
1:A:292:ARG:CG	1:A:302:VAL:HG22	2.51	0.41
1:A:386:GLN:HA	1:A:387:PRO:HD3	1.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	207/233 (89%)	205 (99%)	2 (1%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	180/233 (77%)	177 (98%)	3 (2%)	0	100	100
All	All	387/466 (83%)	382 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/214 (91%)	193 (100%)	1 (0%)	91	90
1	B	176/214 (82%)	171 (97%)	5 (3%)	49	37
All	All	370/428 (86%)	364 (98%)	6 (2%)	71	63

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	256	THR
1	B	293	GLU
1	B	295	GLN
1	B	301	ARG
1	B	336[A]	ILE
1	B	336[B]	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

7 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	1444	1,2	14,14,15	0.52	0	15,19,21	0.97	0
2	NAG	A	1445	2	14,14,15	0.59	0	15,19,21	0.86	0
2	BMA	A	1446	2	11,11,12	0.37	0	13,15,17	0.76	0
2	MAN	A	1447	2	11,11,12	0.52	0	13,15,17	0.85	0
2	NAG	A	1448	2	14,14,15	0.57	0	15,19,21	0.60	0
2	MAN	A	1449	2	11,11,12	0.52	0	13,15,17	0.89	0
2	FUL	A	1450	2	9,10,11	0.33	0	13,14,16	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1444	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	1445	2	-	0/6/23/26	0/1/1/1
2	BMA	A	1446	2	-	0/2/19/22	0/1/1/1
2	MAN	A	1447	2	-	0/2/19/22	0/1/1/1
2	NAG	A	1448	2	-	0/6/23/26	0/1/1/1
2	MAN	A	1449	2	-	0/2/19/22	0/1/1/1
2	FUL	A	1450	2	-	0/0/17/20	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	207/233 (88%)	0.67	25 (12%) 5 8	18, 37, 85, 110	0
1	B	183/233 (78%)	0.85	33 (18%) 2 2	19, 36, 97, 156	0
All	All	390/466 (83%)	0.75	58 (14%) 3 4	18, 37, 94, 156	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	323	VAL	7.4
1	B	239	SER	6.4
1	A	268	HIS	6.2
1	A	266	VAL	6.2
1	A	269	GLU	5.7
1	A	300	TYR	5.6
1	B	301	ARG	5.3
1	B	240	VAL	5.1
1	B	278	TYR	4.9
1	B	303	VAL	4.7
1	B	262	VAL	4.6
1	A	240	VAL	4.4
1	B	275	PHE	4.2
1	B	285	HIS	4.1
1	B	322	LYS	4.0
1	A	264	VAL	3.8
1	B	333	GLU	3.8
1	B	279	VAL	3.8
1	B	286	ASN	3.6
1	B	284	VAL	3.6
1	B	289	THR	3.5
1	B	321	CYS	3.5
1	B	281	GLY	3.3
1	A	328	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	276	ASN	3.2
1	B	243	PHE	3.2
1	A	298	SER	3.1
1	A	355	ARG	3.1
1	B	241	ALA	3.0
1	A	324	SER	3.0
1	A	238	PRO	2.9
1	B	293	GLU	2.9
1	A	239	SER	2.8
1	A	296	TYR	2.8
1	B	295	GLN	2.7
1	B	302	VAL	2.7
1	B	292	ARG	2.6
1	B	385	GLY	2.6
1	B	444	SER	2.6
1	A	332	ILE	2.5
1	B	263	VAL	2.5
1	A	284	VAL	2.5
1	B	238	PRO	2.4
1	A	292	ARG	2.4
1	A	282	VAL	2.4
1	B	334	LYS	2.3
1	A	263	VAL	2.2
1	B	277	TRP	2.2
1	B	294	GLU	2.2
1	A	326	LYS	2.2
1	B	283	GLU	2.2
1	A	237	GLY	2.2
1	A	270	ASP	2.2
1	A	331	PRO	2.1
1	A	333	GLU	2.1
1	B	291	PRO	2.1
1	A	272	GLU	2.0
1	A	384	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAG	A	1448	14/15	0.76	0.27	2.35	53,62,68,71	0
2	NAG	A	1445	14/15	0.82	0.28	-	64,68,71,74	0
2	BMA	A	1446	11/12	0.80	0.33	-	62,66,68,69	0
2	MAN	A	1449	11/12	0.68	0.35	-	68,74,78,81	0
2	FUL	A	1450	10/11	0.68	0.32	-	95,97,100,101	0
2	MAN	A	1447	11/12	0.88	0.23	-	67,68,71,73	0
2	NAG	A	1444	14/15	0.72	0.45	-	62,74,84,90	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	CL	A	1451	1/1	1.00	0.10	-2.44	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.