



wwPDB X-ray Structure Validation Summary Report

Feb 14, 2017 – 12:08 pm GMT

PDB ID : 1BO5
Title : CRYSTAL STRUCTURE OF THE COMPLEX BETWEEN ESCHERICHIA COLI GLYCEROL KINASE AND THE ALLOSTERIC REGULATOR FRUCTOSE 1,6-BISPHOSPHATE.
Authors : Ormo, M.; Bystrom, C.E.; Remington, S.J.
Deposited on : 1998-08-10
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	1009 (3.20-3.20)
Ramachandran outliers	110173	1118 (3.22-3.18)
Sidechain outliers	110143	1117 (3.22-3.18)
RSRZ outliers	101464	1020 (3.22-3.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	O	501	 23% 50% 22% . .
1	Z	501	 21% 52% 22% . .

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 7818 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (GLYCEROL KINASE).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	O	498	3878	2453	671	735	19	0	0	0
1	Z	498	3888	2456	674	739	19	0	0	0

- Molecule 2 is FRUCTOSE-1,6-DIPHOSPHATE (three-letter code: FBP) (formula: $C_6H_{14}O_{12}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
2	O	1	40	12	24	4	0	1

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	O	1	Total	C	O	0	0
			6	3	3		
3	Z	1	Total	C	O	0	0
			6	3	3		

L441	G128
G442	L130
A443	V131
A444	N132
Y445	D133
L446	P134
A447	G135
G448	F136
L449	S137
A450	G138
V451	D139
G452	D201
F453	K140
W454	V141
Q455	K142
W456	W143
L457	E205
D458	L145
E459	L207
L460	H147
Q461	V148
E462	E149
K463	G150
A464	S151
Y465	R152
L466	R154
E467	A155
R468	E156
E469	V157
F470	R218
R471	G158
P472	S220
E475	S221
T476	L160
T477	L161
E478	F162
R479	F163
N480	G164
Y481	Y224
R482	G225
Y483	T164
A484	V165
G485	W166
W486	D166
A489	L169
V490	L170
R491	W171
A492	K172
A493	M173
M494	T174
A495	Q175
W496	G176
E497	R177
E498	V178
H499	H179
ASP	V180
GLU	T181
	D182
	Y183
	T184
	M185
	A186
	S187
	R188
	G252
	T189
	Q253
	K232
	G233
	G234
	T235
	R236
	I237
	P238
	I239
	S240
	G241
	I242
	A243
	G244
	D245
	M308
	A309
	G310
	Q246
	A248
	A249
	L250
	P251
	G252
	R188
	T189
	L316
	L254
	C255
	V256
	K257
	E258
	G259
	W260
	A261
	K262
	M263
	T264
	Y265
	T267
	G268
	E205
	I144
	F270
	L207
	M271
	L272
	M273
	M274
	E277
	V280
	K281
	S282
	E283
	L286
	L287
	T288
	T289
	L290
	A291
	G292
	C293
	P294
	T295
	G296
	E297
	V298
	N299
	Y300
	A301
	L302
	E303
	G304
	A305
	V306
	F307
	M308
	A309
	G310
	A311
	S312
	I313
	Q314
	W315
	L316
	B317
	D318
	E319
	K320
	K321
	L322
	T323
	M324
	D325
	A326
	T327
	R328
	D329
	S329
	E330
	Y331
	F332
	C269
	A333
	T334
	Q336
	A337
	V336
	Q337
	M338
	T339
	N340
	G341
	V342
	Y343
	V344
	V345
	P346
	A347
	F348
	T349
	G350
	L351
	A291
	G352
	A353
	P354
	Y355
	W356
	D357
	P358
	Y359
	A360
	R361
	G362
	A363
	I364
	F365
	T368
	V371
	N372
	A373
	S374
	H375
	I376
	I377
	R378
	A379
	T428
	R429
	V430
	E431
	R432
	P433
	E434
	V435
	R436
	E437
	V438
	T439
	A440

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	169.41Å 169.41Å 204.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 3.20 19.97 – 3.00	Depositor EDS
% Data completeness (in resolution range)	94.0 (20.00-3.20) 90.3 (19.97-3.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 2.98Å)	Xtrriage
Refinement program	TNT V. 5-F	Depositor
R, R_{free}	0.211 , (Not available) 0.190 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	44.4	Xtrriage
Anisotropy	0.024	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 92.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7818	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FBP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	O	1.17	35/3958 (0.9%)	1.51	55/5373 (1.0%)
1	Z	1.18	32/3968 (0.8%)	1.53	55/5387 (1.0%)
All	All	1.18	67/7926 (0.8%)	1.52	110/10760 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	O	1	1

The worst 5 of 67 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	Z	51	GLU	CD-OE1	8.97	1.35	1.25
1	Z	92	GLU	CD-OE1	8.22	1.34	1.25
1	Z	382	GLU	CD-OE2	8.19	1.34	1.25
1	O	51	GLU	CD-OE1	8.05	1.34	1.25
1	Z	212	GLU	CD-OE2	7.96	1.34	1.25

The worst 5 of 110 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	Z	22	ASP	CB-CG-OD1	8.79	126.21	118.30
1	O	201	ASP	CB-CG-OD1	-8.59	110.57	118.30
1	Z	10	ASP	CB-CG-OD1	-8.38	110.76	118.30
1	Z	409	ASP	CB-CG-OD2	-8.35	110.79	118.30
1	O	24	ASP	CB-CG-OD1	-7.97	111.12	118.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	O	21	MET	CA

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	O	351	LEU	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	O	3878	0	3774	472	0
1	Z	3888	0	3778	463	0
2	O	40	0	20	3	0
3	O	6	0	8	2	0
3	Z	6	0	8	0	0
All	All	7818	0	7588	935	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 61.

The worst 5 of 935 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:O:83:ARG:HH11	1:O:83:ARG:HG3	1.04	1.11
1:O:145:LEU:HD11	1:O:213:MET:HE1	1.30	1.11
1:Z:438:VAL:HA	1:Z:441:LEU:HD12	1.30	1.06
1:Z:71:SER:HB2	1:Z:235:THR:HG21	1.32	1.05
1:O:255:CYS:HB3	1:O:260:MET:HB3	1.39	1.03

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	O	496/501 (99%)	406 (82%)	74 (15%)	16 (3%)	5	30
1	Z	496/501 (99%)	414 (84%)	67 (14%)	15 (3%)	5	32
All	All	992/1002 (99%)	820 (83%)	141 (14%)	31 (3%)	5	31

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	O	99	ASN
1	O	121	GLU
1	O	151	SER
1	Z	149	GLU
1	O	72	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	O	396/412 (96%)	301 (76%)	95 (24%)	1	3
1	Z	398/412 (97%)	295 (74%)	103 (26%)	0	2
All	All	794/824 (96%)	596 (75%)	198 (25%)	1	2

5 of 198 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	O	466	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	Z	74	ILE
1	Z	455	GLN
1	O	477	THR
1	Z	29	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	Z	11	GLN
1	Z	179	HIS
1	Z	461	GLN
1	Z	147	HIS
1	Z	228	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FBP	O	502[A]	-	18,20,20	1.00	0	23,32,32	1.38	2 (8%)
2	FBP	O	502[B]	-	18,20,20	0.92	0	23,32,32	1.16	2 (8%)
3	GOL	O	601	-	5,5,5	0.38	0	5,5,5	0.48	0
3	GOL	Z	601	-	5,5,5	0.30	0	5,5,5	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FBP	O	502[A]	-	-	0/13/32/32	0/1/1/1
2	FBP	O	502[B]	-	-	0/13/32/32	0/1/1/1
3	GOL	O	601	-	-	0/4/4/4	0/0/0/0
3	GOL	Z	601	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	O	502[B]	FBP	O3P-P1-O2P	2.81	118.94	107.61
2	O	502[B]	FBP	O6P-P2-O5P	2.91	119.34	107.61
2	O	502[A]	FBP	O6P-P2-O5P	2.96	119.54	107.61
2	O	502[A]	FBP	O3P-P1-O2P	4.61	126.22	107.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	O	502[A]	FBP	2	0
2	O	502[B]	FBP	1	0
3	O	601	GOL	2	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	O	498/501 (99%)	-0.85	0 100 100	12, 28, 62, 90	0
1	Z	498/501 (99%)	-0.89	1 (0%) 94 93	10, 26, 60, 89	0
All	All	996/1002 (99%)	-0.87	1 (0%) 95 95	10, 27, 61, 90	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Z	325	ASP	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	FBP	O	502[B]	20/20	0.88	0.26	1.80	49,49,49,49	20
2	FBP	O	502[A]	20/20	0.88	0.26	1.70	49,49,49,49	20

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	Z	601	6/6	0.98	0.14	1.24	10,11,15,34	0
3	GOL	O	601	6/6	0.97	0.14	0.97	10,18,32,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.