



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:17 am GMT

PDB ID : 2BOV
Title : MOLECULAR RECOGNITION OF AN ADP-RIBOSYLATING CLOSTRIDIUM BOTULINUM C3 EXOENZYME BY RALA GTPASE
Authors : Holbourn, K.P.; Sutton, J.M.; Evans, H.R.; Shone, C.C.; Acharya, K.R.
Deposited on : 2005-04-14
Resolution : 2.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	1026 (2.68-2.64)
Ramachandran outliers	110173	1010 (2.68-2.64)
Sidechain outliers	110143	1010 (2.68-2.64)
RSRZ outliers	101464	3511 (2.70-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	206	<div> <div>2%</div> <div>66%</div> <div>17%</div> <div>•</div> <div>16%</div> </div>
2	B	251	<div> <div>2%</div> <div>56%</div> <div>25%</div> <div>•</div> <div>17%</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3314 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called RAS-RELATED PROTEIN RAL-A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	174	Total	C	N	O	S	11	0	1
			1389	870	235	278	6			

- Molecule 2 is a protein called MONO-ADP-RIBOSYLTRANSFERASE C3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	208	Total	C	N	O	S	32	0	0
			1635	1038	279	311	7			

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Mg 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	131	Total 131	O 131	0	0
5	B	130	Total 130	O 130	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RAS-RELATED PROTEIN RAL-A



• Molecule 2: MONO-ADP-RIBOSYLTRANSFERASE C3



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.63Å 90.84Å 100.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.66 43.94 – 2.66	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-2.66) 97.2 (43.94-2.66)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.94 (at 2.65Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.206 , 0.274 0.199 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	21.1	Xtriage
Anisotropy	0.292	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3314	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/1409	0.56	0/1895
2	B	0.34	0/1664	0.57	0/2234
All	All	0.35	0/3073	0.57	0/4129

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1389	0	1349	26	1
2	B	1635	0	1658	48	1
3	A	28	0	12	0	0
4	A	1	0	0	0	0
5	A	131	0	0	2	0
5	B	130	0	0	7	0
All	All	3314	0	3019	73	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1131:ASP:HB3	2:B:1132:PRO:HD2	1.70	0.73
1:A:88:GLY:HA3	1:A:175:ILE:HD11	1.71	0.73
1:A:33:GLN:HG3	1:A:161:ARG:NH1	2.04	0.72
1:A:72:GLN:H	1:A:72:GLN:NE2	1.88	0.71
2:B:1224:HIS:HD2	2:B:1226:ASP:OD1	1.75	0.69
1:A:33:GLN:HG3	1:A:161:ARG:HH11	1.59	0.66
1:A:70:ALA:HB1	1:A:72:GLN:HE22	1.60	0.64
2:B:1145:ASN:HB3	5:B:2065:HOH:O	2.00	0.61
2:B:1093:ASN:HB3	2:B:1096:VAL:HG22	1.82	0.61
2:B:1173:ILE:HB	2:B:1217:LEU:HB2	1.84	0.60
2:B:1209:PHE:CZ	2:B:1212:GLN:HB3	2.37	0.59
2:B:1180:VAL:HG11	2:B:1212:GLN:HG3	1.85	0.59
2:B:1150:ILE:HG12	2:B:1237:ILE:HG13	1.84	0.58
2:B:1250:PRO:O	2:B:1251:LYS:HB3	2.05	0.57
2:B:1127:PHE:HE2	2:B:1191:LYS:HE2	1.70	0.56
1:A:91:CYS:HB2	1:A:122:PHE:CZ	2.41	0.56
2:B:1070:LYS:O	2:B:1074:GLU:HG3	2.06	0.55
2:B:1132:PRO:HD3	2:B:1187:PRO:HG3	1.89	0.55
2:B:1150:ILE:HD13	2:B:1235:LYS:C	2.27	0.54
1:A:112:LEU:HD22	1:A:117:ASP:O	2.06	0.54
2:B:1251:LYS:HG2	2:B:1251:LYS:O	2.06	0.54
2:B:1126:LEU:CD2	2:B:1215:MET:HG2	2.37	0.54
1:A:12:LEU:HD11	1:A:61:GLU:O	2.09	0.53
2:B:1126:LEU:HD22	2:B:1215:MET:HG2	1.90	0.53
2:B:1140:GLN:HB2	5:B:2061:HOH:O	2.09	0.53
2:B:1063:TYR:HA	2:B:1066:TYR:CD2	2.44	0.52
1:A:169:PHE:O	1:A:173:ARG:HG3	2.09	0.52
2:B:1183:PHE:O	2:B:1185:GLY:N	2.39	0.52
2:B:1249:ASN:N	2:B:1250:PRO:HD3	2.26	0.50
2:B:1051:ASN:ND2	2:B:1054:GLN:H	2.09	0.50
1:A:47:LYS:HE3	1:A:49:ASP:OD2	2.12	0.50
2:B:1128:ARG:HA	2:B:1183:PHE:CZ	2.47	0.49
1:A:161:ARG:HG3	1:A:161:ARG:HH11	1.76	0.49
2:B:1102:SER:HA	2:B:1105:ILE:HG22	1.95	0.48
2:B:1051:ASN:HD21	2:B:1053:ASP:HB2	1.79	0.47
1:A:90:LEU:HG	1:A:171:LEU:HD22	1.96	0.47
1:A:126:GLY:O	1:A:155:GLU:HA	2.15	0.47
1:A:52:ARG:HA	1:A:64:ILE:O	2.15	0.47
2:B:1208:ALA:O	2:B:1209:PHE:C	2.54	0.47
2:B:1244:MET:HE3	5:B:2120:HOH:O	2.14	0.47
2:B:1063:TYR:HA	2:B:1066:TYR:CE2	2.50	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1165:LYS:HE3	5:B:2069:HOH:O	2.16	0.46
2:B:1149:THR:HA	2:B:1235:LYS:HB3	1.97	0.46
2:B:1172:TYR:CZ	2:B:1219:ARG:HD3	2.51	0.46
2:B:1233:ASP:HB2	5:B:2113:HOH:O	2.16	0.45
2:B:1113:LYS:HB2	5:B:2021:HOH:O	2.16	0.45
1:A:50:SER:HA	1:A:66:ILE:O	2.17	0.45
2:B:1181:SER:C	2:B:1183:PHE:N	2.70	0.45
2:B:1189:ILE:HG13	2:B:1236:GLN:NE2	2.32	0.44
2:B:1131:ASP:CG	2:B:1186:ARG:HH11	2.22	0.44
1:A:127:ASN:O	1:A:128:LYS:HB2	2.18	0.43
2:B:1085:GLU:HG3	5:B:2031:HOH:O	2.18	0.43
2:B:1125:MET:SD	2:B:1193:LYS:HG2	2.59	0.43
1:A:33:GLN:O	1:A:37:ASP:HA	2.19	0.43
1:A:166:LYS:HE2	5:A:2118:HOH:O	2.18	0.43
2:B:1248:ILE:C	2:B:1250:PRO:HD3	2.40	0.43
2:B:1076:ILE:O	2:B:1080:THR:HG23	2.19	0.43
2:B:1087:ASN:O	2:B:1091:ARG:HG3	2.19	0.42
2:B:1111:LEU:HD23	2:B:1172:TYR:CE2	2.54	0.42
2:B:1127:PHE:CE2	2:B:1191:LYS:HG2	2.53	0.42
1:A:90:LEU:HD23	1:A:123:LEU:HB2	2.00	0.42
1:A:176:ARG:HG3	5:A:2001:HOH:O	2.18	0.42
1:A:161:ARG:NH1	1:A:161:ARG:HG3	2.35	0.42
1:A:115:LYS:O	1:A:116:GLU:C	2.58	0.41
2:B:1248:ILE:HG22	2:B:1250:PRO:HD3	2.02	0.41
2:B:1098:ASN:C	2:B:1098:ASN:HD22	2.24	0.41
2:B:1181:SER:C	2:B:1183:PHE:H	2.23	0.41
1:A:139:VAL:HG21	2:B:1077:VAL:HG11	2.03	0.40
2:B:1094:LYS:HD3	2:B:1170:TYR:CE2	2.57	0.40
2:B:1115:PHE:CD1	2:B:1218:PRO:HG3	2.56	0.40
1:A:91:CYS:CB	1:A:122:PHE:CZ	3.05	0.40
1:A:79:ARG:HD3	1:A:80:ASP:OD1	2.20	0.40
1:A:119:ASN:O	1:A:178:ARG:NH1	2.54	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:ARG:NH1	2:B:1100:PHE:O[3_645]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	172/206 (84%)	159 (92%)	12 (7%)	1 (1%)	28	43
2	B	206/251 (82%)	193 (94%)	11 (5%)	2 (1%)	18	28
All	All	378/457 (83%)	352 (93%)	23 (6%)	3 (1%)	22	34

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	1209	PHE
2	B	1212	GLN
1	A	182	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	150/178 (84%)	146 (97%)	4 (3%)	50	71
2	B	177/213 (83%)	170 (96%)	7 (4%)	36	55
All	All	327/391 (84%)	316 (97%)	11 (3%)	42	61

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	72	GLN
1	A	79	ARG
1	A	119	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	123	LEU
2	B	1089	LYS
2	B	1098	ASN
2	B	1111	LEU
2	B	1143	LEU
2	B	1164	ASN
2	B	1220	HIS
2	B	1233	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	GLN
1	A	127	ASN
1	A	163	ASN
2	B	1051	ASN
2	B	1060	ASN
2	B	1093	ASN
2	B	1098	ASN
2	B	1140	GLN
2	B	1224	HIS
2	B	1236	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	GDP	A	1184	4	25,30,30	1.55	4 (16%)	26,47,47	1.99	5 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDP	A	1184	4	-	0/12/32/32	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1184	GDP	O3'-C3'	2.25	1.48	1.43
3	A	1184	GDP	C2-N1	2.46	1.39	1.35
3	A	1184	GDP	PB-O3A	3.88	1.66	1.60
3	A	1184	GDP	C6-N1	4.86	1.41	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1184	GDP	C5-C6-N1	-6.95	113.58	123.48
3	A	1184	GDP	N3-C2-N1	-2.95	123.15	127.46
3	A	1184	GDP	O3'-C3'-C4'	-2.30	104.37	111.09
3	A	1184	GDP	O2'-C2'-C1'	2.10	118.18	111.61
3	A	1184	GDP	C6-N1-C2	4.82	123.00	116.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	174/206 (84%)	-0.20	2 (1%) 80 80	10, 18, 43, 70	3 (1%)
2	B	206/251 (82%)	-0.03	4 (1%) 67 66	8, 20, 43, 63	6 (2%)
All	All	380/457 (83%)	-0.11	6 (1%) 72 72	8, 19, 44, 70	9 (2%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1181	SER	5.0
1	A	117	ASP	3.2
2	B	1251	LYS	3.1
2	B	1180	VAL	3.1
2	B	1182	GLN	2.6
1	A	183	SER	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GDP	A	1184	28/28	0.99	0.15	-0.18	13,20,24,27	0
4	MG	A	1185	1/1	0.92	0.13	-0.67	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.