



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 02:59 am GMT

PDB ID : 2BOY
Title : CRYSTAL STRUCTURE OF 3-CHLOROCATECHOL 1,2-DIOXYGENASE
FROM RHODOCOCCLUS OPACUS 1CP
Authors : Ferraroni, M.; Solyanikova, I.P.; Kolomytseva, M.P.; Scozzafava, A.; Golovleva,
L.A.; Briganti, F.
Deposited on : 2005-04-15
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	254	<div> <div>3%</div> <div>86%</div> <div>11%</div> <div>.</div> </div>
1	B	254	<div> <div>4%</div> <div>85%</div> <div>13%</div> <div>..</div> </div>
1	C	254	<div> <div>2%</div> <div>87%</div> <div>9%</div> <div>..</div> </div>
1	D	254	<div> <div>4%</div> <div>85%</div> <div>11%</div> <div>..</div> </div>
1	E	254	<div> <div>4%</div> <div>80%</div> <div>16%</div> <div>...</div> </div>
1	F	254	<div> <div>2%</div> <div>84%</div> <div>11%</div> <div>..</div> </div>
1	G	254	<div> <div>3%</div> <div>80%</div> <div>15%</div> <div>...</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	H	254	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	BHO	B	1256[B]	-	-	X	-
3	BHO	E	1256[A]	-	-	X	-
3	BHO	F	1256[B]	-	-	X	-
3	BHO	H	1256[B]	-	-	X	-
4	LPP	A	1257	-	-	-	X
4	LPP	B	1257	-	-	-	X
4	LPP	C	1257	-	-	-	X
4	LPP	D	1257	-	-	-	X
4	LPP	E	1257	-	-	-	X
4	LPP	F	1257	-	-	-	X
4	LPP	G	1257	-	-	-	X
4	LPP	H	1257	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 17938 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-CHLOROCATECHOL 1,2-DIOXYGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	253	Total	C	N	O	S	0	4	0
			1990	1279	324	383	4			
1	B	252	Total	C	N	O	S	0	2	0
			1967	1266	319	378	4			
1	C	251	Total	C	N	O	S	0	4	0
			1973	1267	322	380	4			
1	D	251	Total	C	N	O	S	0	5	0
			1975	1268	323	380	4			
1	E	252	Total	C	N	O	S	0	4	2
			1973	1269	320	380	4			
1	F	248	Total	C	N	O	S	0	5	0
			1952	1252	318	378	4			
1	G	251	Total	C	N	O	S	0	1	1
			1952	1259	316	373	4			
1	H	248	Total	C	N	O	S	0	1	0
			1924	1240	314	366	4			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	44	LYS	GLN	CONFLICT	UNP Q8G9L3
B	44	LYS	GLN	CONFLICT	UNP Q8G9L3
C	44	LYS	GLN	CONFLICT	UNP Q8G9L3
D	44	LYS	GLN	CONFLICT	UNP Q8G9L3
E	44	LYS	GLN	CONFLICT	UNP Q8G9L3
F	44	LYS	GLN	CONFLICT	UNP Q8G9L3
G	44	LYS	GLN	CONFLICT	UNP Q8G9L3
H	44	LYS	GLN	CONFLICT	UNP Q8G9L3

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	1	Total Fe 1 1	0	0
2	D	1	Total Fe 1 1	0	0
2	E	1	Total Fe 1 1	0	0
2	H	1	Total Fe 1 1	0	0
2	B	1	Total Fe 1 1	0	0
2	C	1	Total Fe 1 1	0	0
2	A	1	Total Fe 1 1	0	0
2	F	1	Total Fe 1 1	0	0

- Molecule 3 is BENZHYDROXAMIC ACID (three-letter code: BHO) (formula: $C_7H_7NO_2$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 10 7 1 2	0	0
3	B	1	Total C N O 20 14 2 4	0	1
3	C	1	Total C N O 20 14 2 4	0	1
3	D	1	Total C N O 14 8 2 4	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	E	1	Total	C	N	O	0	1
			20	14	2	4		
3	F	1	Total	C	N	O	0	1
			20	14	2	4		
3	G	1	Total	C	N	O	0	0
			10	7	1	2		
3	H	1	Total	C	N	O	0	1
			20	14	2	4		

- Molecule 4 is 2-(HEXADECANOYLOXY)-1-[(PHOSPHONOOXY)METHYL]ETHYL HEXADECANOATE (three-letter code: LPP) (formula: C₃₅H₆₉O₈P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O		0	0
			33	29	4			
4	B	1	Total	C	O		0	0
			33	29	4			
4	C	1	Total	C	O		0	0
			34	30	4			
4	D	1	Total	C	O	P	0	0
			36	30	5	1		
4	E	1	Total	C	O		0	0
			34	31	3			
4	F	1	Total	C	O		0	0
			34	31	3			
4	G	1	Total	C	O		0	0
			34	31	3			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	H	1	Total	C	O	0	0
			36	32	4		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	G	1	Total	Mg	0	0
			1	1		
5	D	2	Total	Mg	0	0
			2	2		
5	E	1	Total	Mg	0	0
			1	1		
5	H	2	Total	Mg	0	0
			2	2		
5	B	3	Total	Mg	0	0
			3	3		
5	C	4	Total	Mg	0	0
			4	4		
5	A	2	Total	Mg	0	0
			2	2		
5	F	1	Total	Mg	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	267	Total	O	0	0
			267	267		
6	B	241	Total	O	0	0
			241	241		
6	C	203	Total	O	0	0
			203	203		
6	D	241	Total	O	0	0
			241	241		
6	E	233	Total	O	0	0
			233	233		
6	F	230	Total	O	0	0
			230	230		
6	G	180	Total	O	0	0
			180	180		
6	H	205	Total	O	0	0
			205	205		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: 3-CHLOROCATECHOL 1,2-DIOXYGENASE




• Molecule 1: 3-CHLOROCATECHOL 1,2-DIOXYGENASE




• Molecule 1: 3-CHLOROCATECHOL 1,2-DIOXYGENASE






• Molecule 1: 3-CHLOROCATECHOL 1,2-DIOXYGENASE









• Molecule 1: 3-CHLOROCATECHOL 1,2-DIOXYGENASE





• Molecule 1: 3-CHLOROCATECHOL 1,2-DIOXYGENASE



• Molecule 1: 3-CHLOROCATECHOL 1,2-DIOXYGENASE



• Molecule 1: 3-CHLOROCATECHOL 1,2-DIOXYGENASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	83.18Å 86.61Å 93.45Å 85.37° 66.53° 76.94°	Depositor
Resolution (Å)	84.51 – 1.90 28.12 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.2 (84.51-1.90) 92.8 (28.12-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.35 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.176 , 0.219 0.177 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	24.0	Xtriage
Anisotropy	0.301	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.001 for -h,-k,-h+l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	17938	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, FE, BHO, LPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	0/2061	0.89	7/2806 (0.2%)
1	B	0.79	0/2027	0.87	6/2763 (0.2%)
1	C	0.80	0/2043	0.87	5/2785 (0.2%)
1	D	0.77	0/2050	0.86	4/2792 (0.1%)
1	E	0.78	0/2044	0.90	13/2783 (0.5%)
1	F	0.76	0/2026	0.88	10/2760 (0.4%)
1	G	0.69	0/2007	0.88	8/2734 (0.3%)
1	H	0.75	0/1978	0.89	10/2696 (0.4%)
All	All	0.77	0/16236	0.88	63/22119 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

The worst 5 of 63 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	95	ARG	NE-CZ-NH2	-11.84	114.38	120.30
1	E	122	LEU	CA-CB-CG	9.49	137.14	115.30
1	E	95	ARG	NE-CZ-NH1	8.55	124.58	120.30
1	G	95	ARG	NE-CZ-NH2	-8.11	116.25	120.30
1	E	128	ASP	CB-CG-OD2	7.95	125.46	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	212	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1990	0	1906	27	0
1	B	1967	0	1862	29	0
1	C	1973	0	1868	27	0
1	D	1975	0	1885	27	0
1	E	1973	0	1889	43	0
1	F	1952	0	1847	25	0
1	G	1952	0	1860	31	0
1	H	1924	0	1828	48	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	A	10	0	6	3	0
3	B	20	0	14	9	0
3	C	20	0	11	3	0
3	D	14	0	1	2	0
3	E	20	0	12	8	0
3	F	20	0	11	4	0
3	G	10	0	6	3	0
3	H	20	0	13	13	0
4	A	33	0	47	0	0
4	B	33	0	47	1	0
4	C	34	0	49	0	0
4	D	36	0	51	0	0
4	E	34	0	55	0	0
4	F	34	0	55	2	0
4	G	34	0	55	1	0
4	H	36	0	57	1	0
5	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	3	0	0	0	0
5	C	4	0	0	0	0
5	D	2	0	0	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
5	G	1	0	0	0	0
5	H	2	0	0	0	0
6	A	267	0	0	3	0
6	B	241	0	0	7	0
6	C	203	0	0	2	0
6	D	241	0	0	7	0
6	E	233	0	0	8	0
6	F	230	0	0	6	0
6	G	180	0	0	4	0
6	H	205	0	0	8	0
All	All	17938	0	15435	265	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 265 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:167[B]:TYR:OH	3:H:1256[B]:BHO:H6	1.06	1.19
1:C:167[B]:TYR:CE1	1:C:191:HIS:CE1	2.42	1.08
1:A:212:ASN:H	1:A:212:ASN:ND2	1.46	1.05
1:F:112:GLN:HG2	6:F:2136:HOH:O	1.56	1.04
1:C:167[B]:TYR:HE1	1:C:191:HIS:CE1	1.75	1.03

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/254 (100%)	248 (97%)	6 (2%)	1 (0%)	38	26
1	B	250/254 (98%)	247 (99%)	3 (1%)	0	100	100
1	C	251/254 (99%)	247 (98%)	4 (2%)	0	100	100
1	D	252/254 (99%)	248 (98%)	4 (2%)	0	100	100
1	E	252/254 (99%)	246 (98%)	5 (2%)	1 (0%)	38	26
1	F	249/254 (98%)	244 (98%)	5 (2%)	0	100	100
1	G	248/254 (98%)	243 (98%)	4 (2%)	1 (0%)	38	26
1	H	245/254 (96%)	239 (98%)	6 (2%)	0	100	100
All	All	2002/2032 (98%)	1962 (98%)	37 (2%)	3 (0%)	51	45

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	152	ASP
1	G	152	ASP
1	A	237	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	208/209 (100%)	199 (96%)	9 (4%)	33	22
1	B	200/209 (96%)	192 (96%)	8 (4%)	36	25
1	C	203/209 (97%)	194 (96%)	9 (4%)	33	22
1	D	206/209 (99%)	197 (96%)	9 (4%)	33	22
1	E	206/209 (99%)	188 (91%)	18 (9%)	12	4
1	F	203/209 (97%)	190 (94%)	13 (6%)	20	10
1	G	199/209 (95%)	181 (91%)	18 (9%)	11	4
1	H	196/209 (94%)	181 (92%)	15 (8%)	15	6
All	All	1621/1672 (97%)	1522 (94%)	99 (6%)	22	11

5 of 99 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	123	ASP
1	F	85[A]	GLU
1	H	199	GLU
1	E	224	VAL
1	E	251	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	57	HIS
1	E	185	HIS
1	H	67	ASN
1	E	67	ASN
1	E	75	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 46 ligands modelled in this entry, 24 are monoatomic - leaving 22 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BHO	A	1256	2	10,10,10	3.32	2 (20%)	12,12,12	2.27	3 (25%)
4	LPP	A	1257	-	32,32,43	1.29	2 (6%)	34,34,48	1.47	6 (17%)
3	BHO	B	1256[A]	2	10,10,10	3.35	2 (20%)	12,12,12	1.38	2 (16%)
3	BHO	B	1256[B]	2	10,10,10	3.50	2 (20%)	12,12,12	1.08	2 (16%)
4	LPP	B	1257	-	32,32,43	1.24	2 (6%)	34,34,48	1.41	6 (17%)
3	BHO	C	1256[A]	2	10,10,10	3.49	2 (20%)	12,12,12	1.07	1 (8%)
3	BHO	C	1256[B]	1,2	10,10,10	3.11	2 (20%)	12,12,12	1.18	1 (8%)
4	LPP	C	1257	-	33,33,43	1.24	2 (6%)	35,35,48	1.45	6 (17%)
3	BHO	D	1256[A]	2	10,10,10	3.49	2 (20%)	12,12,12	1.28	1 (8%)
3	BHO	D	1256[B]	2	10,10,10	3.23	2 (20%)	12,12,12	1.75	3 (25%)
4	LPP	D	1257	-	33,35,43	1.29	2 (6%)	35,37,48	1.25	2 (5%)
3	BHO	E	1256[A]	2	10,10,10	3.19	2 (20%)	12,12,12	1.10	0
3	BHO	E	1256[B]	1,2	10,10,10	3.51	3 (30%)	12,12,12	1.19	0
4	LPP	E	1257	-	32,32,43	1.07	2 (6%)	31,31,48	0.85	1 (3%)
3	BHO	F	1256[A]	2	10,10,10	3.50	2 (20%)	12,12,12	1.37	2 (16%)
3	BHO	F	1256[B]	1,2	10,10,10	3.07	2 (20%)	12,12,12	1.10	0
4	LPP	F	1257	-	32,32,43	1.03	2 (6%)	31,31,48	0.85	0
3	BHO	G	1256	2	10,10,10	3.36	2 (20%)	12,12,12	1.67	4 (33%)
4	LPP	G	1257	-	32,32,43	1.06	2 (6%)	31,31,48	0.81	0
3	BHO	H	1256[A]	2	10,10,10	3.28	2 (20%)	12,12,12	1.09	0
3	BHO	H	1256[B]	1,2	10,10,10	3.32	2 (20%)	12,12,12	1.36	2 (16%)
4	LPP	H	1257	-	34,34,43	1.04	2 (5%)	33,33,48	0.86	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BHO	A	1256	2	-	2/6/6/6	0/1/1/1
4	LPP	A	1257	-	-	2/33/33/45	0/0/0/0
3	BHO	B	1256[A]	2	-	2/6/6/6	0/1/1/1
3	BHO	B	1256[B]	2	-	2/6/6/6	0/1/1/1
4	LPP	B	1257	-	-	0/33/33/45	0/0/0/0
3	BHO	C	1256[A]	2	-	0/6/6/6	0/1/1/1
3	BHO	C	1256[B]	1,2	-	2/6/6/6	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	LPP	C	1257	-	-	1/34/34/45	0/0/0/0
3	BHO	D	1256[A]	2	-	0/6/6/6	0/1/1/1
3	BHO	D	1256[B]	2	-	2/6/6/6	0/1/1/1
4	LPP	D	1257	-	-	0/36/37/45	0/0/0/0
3	BHO	E	1256[A]	2	-	2/6/6/6	0/1/1/1
3	BHO	E	1256[B]	1,2	-	2/6/6/6	0/1/1/1
4	LPP	E	1257	-	-	0/29/29/45	0/0/0/0
3	BHO	F	1256[A]	2	-	2/6/6/6	0/1/1/1
3	BHO	F	1256[B]	1,2	-	2/6/6/6	0/1/1/1
4	LPP	F	1257	-	-	0/29/29/45	0/0/0/0
3	BHO	G	1256	2	-	2/6/6/6	0/1/1/1
4	LPP	G	1257	-	-	0/29/29/45	0/0/0/0
3	BHO	H	1256[A]	2	-	0/6/6/6	0/1/1/1
3	BHO	H	1256[B]	1,2	-	2/6/6/6	0/1/1/1
4	LPP	H	1257	-	-	0/31/31/45	0/0/0/0

The worst 5 of 45 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1256[B]	BHO	O2-N	-9.27	1.23	1.39
3	F	1256[A]	BHO	O2-N	-9.21	1.23	1.39
3	C	1256[A]	BHO	O2-N	-9.00	1.24	1.39
3	D	1256[A]	BHO	O2-N	-8.83	1.24	1.39
3	A	1256	BHO	O2-N	-8.70	1.24	1.39

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1256	BHO	O1-C-N	-3.07	116.91	122.96
4	B	1257	LPP	C7-O9-C11	-3.07	114.01	117.90
3	D	1256[B]	BHO	O1-C-C1	-3.00	115.61	120.94
3	D	1256[B]	BHO	C3-C2-C1	-2.52	117.38	120.35
3	D	1256[A]	BHO	C3-C2-C1	-2.52	117.38	120.35

There are no chirality outliers.

5 of 25 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	1256[B]	BHO	C1-C-N-O2
3	E	1256[B]	BHO	O1-C-N-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	F	1256[A]	BHO	C1-C-N-O2
3	F	1256[A]	BHO	O1-C-N-O2
3	G	1256	BHO	C1-C-N-O2

There are no ring outliers.

16 monomers are involved in 50 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1256	BHO	3	0
3	B	1256[A]	BHO	3	0
3	B	1256[B]	BHO	6	0
4	B	1257	LPP	1	0
3	C	1256[A]	BHO	1	0
3	C	1256[B]	BHO	2	0
3	D	1256[B]	BHO	2	0
3	E	1256[A]	BHO	6	0
3	E	1256[B]	BHO	2	0
3	F	1256[B]	BHO	4	0
4	F	1257	LPP	2	0
3	G	1256	BHO	3	0
4	G	1257	LPP	1	0
3	H	1256[A]	BHO	3	0
3	H	1256[B]	BHO	10	0
4	H	1257	LPP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	253/254 (99%)	-0.19	8 (3%)	48	51	16, 22, 38, 57	0
1	B	252/254 (99%)	-0.14	9 (3%)	43	47	17, 23, 36, 55	0
1	C	251/254 (98%)	-0.24	6 (2%)	59	63	17, 25, 38, 53	0
1	D	251/254 (98%)	-0.16	10 (3%)	39	43	17, 25, 40, 52	0
1	E	252/254 (99%)	-0.13	9 (3%)	43	47	18, 25, 42, 53	0
1	F	248/254 (97%)	-0.22	5 (2%)	65	69	16, 24, 43, 53	0
1	G	251/254 (98%)	-0.02	8 (3%)	48	51	19, 30, 47, 66	0
1	H	248/254 (97%)	-0.09	5 (2%)	65	69	18, 28, 43, 53	0
All	All	2006/2032 (98%)	-0.15	60 (2%)	51	54	16, 25, 42, 66	0

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	237	ALA	5.7
1	B	237	ALA	5.6
1	B	254	VAL	5.5
1	G	1	MET	5.3
1	E	237	ALA	5.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	LPP	B	1257	33/44	0.65	0.26	8.01	35,43,64,64	0
4	LPP	E	1257	34/44	0.62	0.29	7.23	32,44,63,64	0
4	LPP	C	1257	34/44	0.60	0.29	7.18	40,45,64,65	0
4	LPP	D	1257	36/44	0.52	0.32	6.79	37,46,69,76	0
4	LPP	F	1257	34/44	0.72	0.24	6.62	33,41,58,60	0
4	LPP	H	1257	36/44	0.65	0.28	6.01	37,45,63,64	0
4	LPP	A	1257	33/44	0.68	0.24	5.04	38,42,58,58	0
4	LPP	G	1257	34/44	0.73	0.22	4.96	35,44,58,61	0
3	BHO	F	1256[A]	10/10	0.95	0.14	1.16	20,26,27,29	10
3	BHO	F	1256[B]	10/10	0.95	0.14	1.12	24,27,29,29	10
3	BHO	E	1256[B]	10/10	0.96	0.12	1.09	16,21,23,23	10
3	BHO	E	1256[A]	10/10	0.96	0.12	1.09	21,22,25,27	10
5	MG	H	1259	1/1	0.95	0.14	1.03	40,40,40,40	0
3	BHO	G	1256	10/10	0.91	0.15	0.91	24,31,41,42	0
3	BHO	D	1256[B]	10/10	0.94	0.13	0.71	25,27,32,39	4
3	BHO	D	1256[A]	10/10	0.94	0.13	0.71	25,26,27,27	4
3	BHO	C	1256[B]	10/10	0.96	0.13	0.60	20,22,24,25	10
3	BHO	C	1256[A]	10/10	0.96	0.13	0.56	17,26,27,28	10
3	BHO	H	1256[A]	10/10	0.95	0.12	0.22	22,28,29,30	10
3	BHO	H	1256[B]	10/10	0.95	0.12	0.19	28,29,30,31	10
3	BHO	B	1256[B]	10/10	0.98	0.14	0.18	26,28,30,30	10
3	BHO	B	1256[A]	10/10	0.98	0.14	0.14	18,20,24,25	10
3	BHO	A	1256	10/10	0.96	0.12	0.05	23,27,37,37	0
2	FE	B	1255	1/1	0.99	0.10	-0.94	21,21,21,21	0
2	FE	H	1255	1/1	0.99	0.07	-1.13	22,22,22,22	0
2	FE	G	1255	1/1	1.00	0.07	-1.42	23,23,23,23	0
2	FE	D	1255	1/1	1.00	0.06	-1.88	23,23,23,23	0
2	FE	F	1255	1/1	1.00	0.06	-2.13	19,19,19,19	0
2	FE	E	1255	1/1	1.00	0.04	-2.25	20,20,20,20	0
2	FE	C	1255	1/1	1.00	0.06	-2.67	21,21,21,21	0
5	MG	D	1259	1/1	0.96	0.13	-	43,43,43,43	0
5	MG	G	1258	1/1	0.94	0.06	-	46,46,46,46	0
5	MG	B	1258	1/1	0.69	0.11	-	54,54,54,54	0
5	MG	B	1259	1/1	0.99	0.04	-	28,28,28,28	0
5	MG	C	1258	1/1	0.97	0.07	-	44,44,44,44	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	MG	E	1258	1/1	0.82	0.13	-	53,53,53,53	0
5	MG	C	1261	1/1	0.85	0.10	-	53,53,53,53	0
5	MG	C	1260	1/1	0.71	0.14	-	51,51,51,51	0
5	MG	D	1258	1/1	0.84	0.13	-	47,47,47,47	0
5	MG	A	1258	1/1	0.94	0.18	-	48,48,48,48	0
2	FE	A	1255	1/1	1.00	0.07	-	19,19,19,19	0
5	MG	A	1259	1/1	0.98	0.04	-	37,37,37,37	0
5	MG	C	1259	1/1	0.94	0.08	-	55,55,55,55	0
5	MG	B	1260	1/1	0.82	0.10	-	52,52,52,52	0
5	MG	H	1258	1/1	0.93	0.08	-	39,39,39,39	0
5	MG	F	1258	1/1	0.95	0.11	-	44,44,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.