



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 12:24 pm GMT

PDB ID : 4BOW  
Title : Crystal structure of LamA\_E269S from *Z. galactanivorans* in complex with laminaritriose and laminaritetraose  
Authors : Labourel, A.; Jeudy, A.; Czjzek, M.; Michel, G.  
Deposited on : 2013-05-22  
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1024 (1.38-1.34)
Clashscore	112137	1063 (1.38-1.34)
Ramachandran outliers	110173	1048 (1.38-1.34)
Sidechain outliers	110143	1048 (1.38-1.34)
RSRZ outliers	101464	1025 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	<div> <div>2%</div> <div> <div></div> <div>81%</div> <div>13%</div> <div>...</div> </div> </div>
1	B	256	<div> <div>%</div> <div> <div></div> <div>82%</div> <div>11%</div> <div>...</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	BGC	B	402[B]	X	-	-	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4754 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENDO-1,3-BETA-GLUCANASE, FAMILY GH16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	248	Total	C	N	O	S	0	11	0
			2068	1320	334	410	4			
1	B	248	Total	C	N	O	S	0	10	0
			2047	1305	330	408	4			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	128	HIS	-	EXPRESSION TAG	UNP G0L5X4
A	129	HIS	-	EXPRESSION TAG	UNP G0L5X4
A	130	HIS	-	EXPRESSION TAG	UNP G0L5X4
A	131	HIS	-	EXPRESSION TAG	UNP G0L5X4
A	132	HIS	-	EXPRESSION TAG	UNP G0L5X4
A	133	HIS	-	EXPRESSION TAG	UNP G0L5X4
A	134	GLY	-	EXPRESSION TAG	UNP G0L5X4
A	135	SER	-	EXPRESSION TAG	UNP G0L5X4
A	269	SER	GLU	ENGINEERED MUTATION	UNP G0L5X4
B	128	HIS	-	EXPRESSION TAG	UNP G0L5X4
B	129	HIS	-	EXPRESSION TAG	UNP G0L5X4
B	130	HIS	-	EXPRESSION TAG	UNP G0L5X4
B	131	HIS	-	EXPRESSION TAG	UNP G0L5X4
B	132	HIS	-	EXPRESSION TAG	UNP G0L5X4
B	133	HIS	-	EXPRESSION TAG	UNP G0L5X4
B	134	GLY	-	EXPRESSION TAG	UNP G0L5X4
B	135	SER	-	EXPRESSION TAG	UNP G0L5X4
B	269	SER	GLU	ENGINEERED MUTATION	UNP G0L5X4

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ca	0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	3	Total	C	O	0	0
			34	18	16		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Na	0	0
			1	1		

- Molecule 5 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	5	Total	C	O	0	2
			57	30	27		


- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	316	Total	O	0	0
			316	316		
6	B	229	Total	O	0	0
			229	229		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ENDO-1,3-BETA-GLUCANASE, FAMILY GH16

Chain A: 



- Molecule 1: ENDO-1,3-BETA-GLUCANASE, FAMILY GH16

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	44.46Å 76.50Å 142.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	67.44 – 1.35 38.44 – 1.35	Depositor EDS
% Data completeness (in resolution range)	99.5 (67.44-1.35) 99.6 (38.44-1.35)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.20 (at 1.35Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.130 , 0.167 0.135 , 0.173	Depositor DCC
$R_{free}$ test set	5366 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	14.4	Xtriage
Anisotropy	0.582	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 44.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	4754	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, NA, CA, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.29	6/2153 (0.3%)	1.24	11/2926 (0.4%)
1	B	1.34	5/2138 (0.2%)	1.21	11/2909 (0.4%)
All	All	1.31	11/4291 (0.3%)	1.23	22/5835 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
5	B	1	0
All	All	1	1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	274	GLU	CD-OE2	8.28	1.34	1.25
1	B	274	GLU	CD-OE2	7.14	1.33	1.25
1	B	182	GLU	CD-OE2	5.81	1.32	1.25
1	A	148	TYR	CB-CG	5.68	1.60	1.51
1	A	316	TYR	CG-CD1	5.66	1.46	1.39
1	B	169	TRP	CD2-CE2	5.65	1.48	1.41
1	A	159	TYR	CD1-CE1	-5.63	1.30	1.39
1	A	299	ASN	CB-CG	-5.17	1.39	1.51
1	A	350	TYR	CG-CD2	5.04	1.45	1.39
1	B	148	TYR	CZ-OH	5.03	1.46	1.37
1	B	220	PHE	CE1-CZ	5.02	1.46	1.37

All (22) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	155	GLU	N-CA-CB	-15.36	82.95	110.60
1	A	274	GLU	OE1-CD-OE2	-9.48	111.92	123.30
1	A	269[A]	SER	N-CA-CB	-7.74	98.89	110.50
1	A	269[B]	SER	N-CA-CB	-7.74	98.89	110.50
1	A	374	PHE	CB-CG-CD1	7.30	125.91	120.80
1	B	155	GLU	N-CA-C	-7.17	91.65	111.00
1	B	364	ASP	CB-CG-OD2	-7.06	111.95	118.30
1	A	374	PHE	CB-CG-CD2	-6.89	115.98	120.80
1	A	271	ASP	CB-CG-OD2	6.78	124.41	118.30
1	B	374	PHE	CB-CG-CD2	-6.78	116.05	120.80
1	A	255	PHE	CB-CG-CD1	5.63	124.74	120.80
1	B	156	LYS	CD-CE-NZ	-5.57	98.89	111.70
1	A	194	ARG	NE-CZ-NH2	-5.53	117.53	120.30
1	B	374	PHE	CB-CG-CD1	5.50	124.65	120.80
1	B	154	PRO	N-CA-C	5.48	126.35	112.10
1	B	154	PRO	C-N-CA	-5.47	108.02	121.70
1	B	271[A]	ASP	CB-CG-OD2	5.39	123.16	118.30
1	B	271[B]	ASP	CB-CG-OD2	5.39	123.16	118.30
1	B	380	ARG	NE-CZ-NH2	-5.38	117.61	120.30
1	A	279	ASP	CB-CG-OD1	5.38	123.14	118.30
1	A	377	ASP	CB-CG-OD2	-5.25	113.58	118.30
1	A	209	TYR	CZ-CE2-CD2	-5.14	115.17	119.80

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	B	402[B]	BGC	C1

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	175	GLN	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2068	0	1932	23	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2047	0	1908	24	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	34	0	30	3	0
4	B	1	0	0	0	0
5	B	57	0	49	3	0
6	A	316	0	0	9	1
6	B	229	0	0	5	2
All	All	4754	0	3919	49	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:403:BGC:O3	6:A:2316:HOH:O	1.55	1.19
1:A:294:SER:HB2	6:A:2228:HOH:O	1.64	0.97
1:B:269[A]:SER:OG	6:B:2128:HOH:O	1.85	0.94
1:B:154:PRO:O	1:B:217:LYS:NZ	2.13	0.78
1:B:276:ASN:HD21	1:B:279:ASP:H	1.36	0.73
1:A:203[A]:GLU:OE1	6:A:2102:HOH:O	2.07	0.71
1:B:276:ASN:ND2	1:B:279:ASP:H	1.90	0.69
1:A:253:ASN:HD21	1:A:256:GLY:H	1.45	0.64
1:B:253:ASN:HD22	1:B:253:ASN:C	2.01	0.64
1:B:271[A]:ASP:OD2	5:B:401[A]:BGC:O1	2.16	0.63
1:B:175:GLN:NE2	1:B:357:MET:H	1.97	0.62
1:A:173:GLU:HB3	1:A:175:GLN:HE22	1.65	0.61
1:A:366:PRO:HG2	6:A:2150:HOH:O	2.01	0.60
1:B:276:ASN:C	1:B:276:ASN:HD22	2.06	0.59
1:B:173:GLU:HB3	1:B:175:GLN:HE22	1.66	0.58
1:B:175:GLN:HE21	1:B:357:MET:H	1.51	0.58
1:A:155:GLU:HG2	6:A:2027:HOH:O	2.05	0.57
1:A:175:GLN:NE2	1:A:357:MET:H	2.03	0.56
1:A:176:HIS:HE1	1:A:205:SER:O	1.88	0.56
1:B:198:GLU:CD	6:B:2069:HOH:O	2.45	0.54
1:B:152:PRO:O	6:B:2017:HOH:O	2.19	0.53
1:A:175:GLN:HE21	1:A:357:MET:H	1.55	0.53
1:A:286:HIS:NE2	1:A:299:ASN:OD1	2.41	0.53
1:B:253:ASN:ND2	1:B:253:ASN:C	2.62	0.53
1:A:151:LYS:HG2	6:A:2032:HOH:O	2.08	0.52

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:253:ASN:HD21	1:B:256:GLY:H	1.57	0.52
6:A:2191:HOH:O	1:B:342[A]:ASN:ND2	2.46	0.49
1:A:238:TRP:CE2	1:A:361:LEU:HB2	2.48	0.48
1:A:170:HIS:CD2	3:A:402:BGC:H4	2.49	0.47
1:A:276:ASN:HD22	1:A:276:ASN:C	2.17	0.47
1:A:288:HIS:HD2	1:A:297:TYR:OH	1.98	0.47
3:A:403:BGC:C3	6:A:2316:HOH:O	2.37	0.47
1:B:339:ASN:HD22	1:B:340:SER:H	1.63	0.46
1:B:238:TRP:CE2	1:B:361:LEU:HB2	2.52	0.45
1:A:151:LYS:HB2	1:A:151:LYS:HE3	1.81	0.45
1:A:339:ASN:HD22	1:A:340:SER:H	1.65	0.44
1:B:187:SER:O	1:B:192:LYS:NZ	2.50	0.44
1:A:179:ASN:HD22	1:A:179:ASN:C	2.21	0.44
1:A:176:HIS:HD2	6:A:2037:HOH:O	1.99	0.43
1:A:253:ASN:ND2	1:A:256:GLY:H	2.13	0.43
1:B:342[A]:ASN:ND2	6:B:2202:HOH:O	2.38	0.43
1:B:271[B]:ASP:OD2	5:B:402[B]:BGC:O5	2.36	0.43
1:A:224:LYS:HA	1:A:318:LEU:O	2.19	0.42
1:A:276:ASN:ND2	1:A:279:ASP:H	2.16	0.42
1:B:243:THR:OG1	1:B:349:HIS:HD2	2.01	0.42
1:B:156:LYS:HE3	6:B:2007:HOH:O	2.19	0.41
1:A:179:ASN:ND2	1:A:179:ASN:C	2.73	0.41
1:B:288:HIS:CE1	5:B:401[A]:BGC:O1	2.74	0.41
1:B:277:GLY:HA3	1:B:362:GLY:O	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:2154:HOH:O	6:B:2195:HOH:O[4_455]	2.17	0.03
6:A:2184:HOH:O	6:B:2144:HOH:O[4_455]	2.19	0.01

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/256 (100%)	249 (97%)	8 (3%)	0	100	100
1	B	256/256 (100%)	245 (96%)	11 (4%)	0	100	100
All	All	513/512 (100%)	494 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/221 (102%)	216 (96%)	9 (4%)	36	5
1	B	224/221 (101%)	217 (97%)	7 (3%)	45	10
All	All	449/442 (102%)	433 (96%)	16 (4%)	39	8

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	151	LYS
1	A	175	GLN
1	A	178	THR
1	A	179	ASN
1	A	199	LYS
1	A	253	ASN
1	A	276	ASN
1	A	299	ASN
1	A	339	ASN
1	B	175	GLN
1	B	179	ASN
1	B	253	ASN
1	B	263	GLU
1	B	276	ASN
1	B	339	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	374	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	175	GLN
1	A	176	HIS
1	A	179	ASN
1	A	253	ASN
1	A	276	ASN
1	A	288	HIS
1	A	339	ASN
1	A	383	GLN
1	B	175	GLN
1	B	179	ASN
1	B	215	ASN
1	B	253	ASN
1	B	276	ASN
1	B	298	GLN
1	B	339	ASN
1	B	349	HIS
1	B	383	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

8 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GLC	A	401	3	12,12,12	1.13	0	17,17,17	1.78	4 (23%)
3	BGC	A	402	3	11,11,12	1.24	1 (9%)	13,15,17	1.44	4 (30%)
3	BGC	A	403	3	11,11,12	0.65	0	13,15,17	1.86	3 (23%)
5	BGC	B	401[A]	5	12,12,12	0.83	0	17,17,17	1.80	3 (17%)
5	BGC	B	402[B]	5	12,12,12	0.75	0	17,17,17	3.04	7 (41%)
5	BGC	B	403	5	11,11,12	0.82	0	13,15,17	1.12	2 (15%)
5	BGC	B	404	5	11,11,12	0.52	0	13,15,17	1.71	3 (23%)
5	BGC	B	405	5	11,11,12	1.28	1 (9%)	13,15,17	1.38	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	A	401	3	-	0/2/22/22	0/1/1/1
3	BGC	A	402	3	-	0/2/19/22	0/1/1/1
3	BGC	A	403	3	-	0/2/19/22	0/1/1/1
5	BGC	B	401[A]	5	-	0/2/22/22	0/1/1/1
5	BGC	B	402[B]	5	1/1/5/5	0/2/22/22	0/1/1/1
5	BGC	B	403	5	-	0/2/19/22	0/1/1/1
5	BGC	B	404	5	-	0/2/19/22	0/1/1/1
5	BGC	B	405	5	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	BGC	C2-C3	-2.06	1.49	1.52
5	B	405	BGC	C2-C3	3.02	1.56	1.52

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	402[B]	BGC	C1-C2-C3	-7.25	97.55	110.65
5	B	402[B]	BGC	C1-O5-C5	-5.96	102.64	113.39
3	A	403	BGC	C1-C2-C3	-4.67	103.73	109.65
5	B	401[A]	BGC	C1-C2-C3	-3.91	103.58	110.65

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	404	BGC	C1-O5-C5	-3.84	106.87	112.17
5	B	401[A]	BGC	C3-C4-C5	-3.81	103.50	110.22
3	A	401	GLC	C4-C3-C2	-3.78	104.18	110.84
3	A	401	GLC	O1-C1-O5	-3.17	100.83	110.20
3	A	403	BGC	O4-C4-C5	-2.98	101.77	109.28
5	B	402[B]	BGC	C4-C3-C2	-2.91	105.70	110.84
3	A	402	BGC	O3-C3-C4	-2.76	104.34	110.36
3	A	403	BGC	O6-C6-C5	-2.70	102.25	111.34
3	A	401	GLC	C3-C4-C5	-2.64	105.57	110.22
5	B	404	BGC	O3-C3-C2	-2.59	105.31	110.02
5	B	401[A]	BGC	O3-C3-C4	-2.55	104.81	110.36
5	B	404	BGC	O4-C4-C3	-2.45	105.02	110.36
5	B	403	BGC	O3-C3-C2	-2.41	105.63	110.02
5	B	405	BGC	O6-C6-C5	-2.41	103.23	111.34
3	A	402	BGC	C1-C2-C3	-2.36	106.66	109.65
5	B	403	BGC	C1-O5-C5	-2.24	109.08	112.17
3	A	402	BGC	O3-C3-C2	-2.21	106.01	110.02
3	A	402	BGC	O4-C4-C5	-2.05	104.11	109.28
3	A	401	GLC	C1-C2-C3	2.23	114.67	110.65
5	B	405	BGC	C2-C3-C4	2.46	115.17	110.88
5	B	402[B]	BGC	C6-C5-C4	2.70	119.31	113.00
5	B	402[B]	BGC	O5-C1-C2	3.02	115.05	110.04
5	B	402[B]	BGC	O1-C1-O5	3.73	121.24	110.20
5	B	402[B]	BGC	O1-C1-C2	4.78	122.92	109.42

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	B	402[B]	BGC	C1

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	BGC	1	0
3	A	403	BGC	2	0
5	B	401[A]	BGC	2	0
5	B	402[B]	BGC	1	0

## 5.6 Ligand geometry

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	248/256 (96%)	-0.27	4 (1%) 72 77	11, 17, 29, 47	0
1	B	248/256 (96%)	-0.32	2 (0%) 86 88	14, 23, 44, 56	0
All	All	496/512 (96%)	-0.29	6 (1%) 79 83	11, 19, 40, 56	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	295[A]	ASP	4.1
1	A	136	ALA	3.1
1	A	293	ASN	2.4
1	B	185	PHE	2.2
1	A	368	ASN	2.1
1	B	155	GLU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	BGC	B	402[B]	12/12	0.97	0.07	0.86	17,20,22,25	12
5	BGC	B	401[A]	12/12	0.97	0.06	0.19	11,14,18,25	12

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GLC	A	401	12/12	0.97	0.06	-0.26	12,14,18,21	0
3	BGC	A	402	11/12	0.99	0.04	-1.46	13,15,16,16	0
5	BGC	B	403	11/12	0.97	0.05	-1.67	16,17,20,20	0
3	BGC	A	403	11/12	0.94	0.12	-	21,25,31,33	0
5	BGC	B	404	11/12	0.94	0.10	-	24,28,32,37	0
5	BGC	B	405	11/12	0.87	0.24	-	41,49,57,59	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CA	A	400	1/1	1.00	0.04	-0.84	17,17,17,17	1
4	NA	B	399	1/1	0.99	0.05	-0.92	35,35,35,35	0
2	CA	B	400	1/1	0.99	0.05	-1.23	34,34,34,34	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.