



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 11:41 pm GMT

PDB ID : 4BPU  
Title : Crystal structure of human primase in heterodimeric form, comprising PriS and truncated PriL lacking the C-terminal Fe-S domain.  
Authors : Kilkenny, M.L.; Perera, R.L.; Pellegrini, L.  
Deposited on : 2013-05-28  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2259 (2.70-2.70)
Clashscore	112137	2590 (2.70-2.70)
Ramachandran outliers	110173	2550 (2.70-2.70)
Sidechain outliers	110143	2550 (2.70-2.70)
RSRZ outliers	101464	2275 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	423	
1	C	423	
2	B	253	
2	D	253	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	1410	-	-	-	X
4	GOL	A	1412	-	-	-	X
4	GOL	A	1413	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20014 atoms, of which 9940 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA PRIMASE SMALL SUBUNIT.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	380	Total	C	H	N	O	S	0	10	0
			6450	2077	3221	563	576	13			
1	C	367	Total	C	H	N	O	S	0	3	0
			6171	1994	3079	536	549	13			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP P49642
A	-1	THR	-	EXPRESSION TAG	UNP P49642
A	0	SER	-	EXPRESSION TAG	UNP P49642
A	72	ALA	LYS	ENGINEERED MUTATION	UNP P49642
A	73	ALA	MET	ENGINEERED MUTATION	UNP P49642
C	-2	GLY	-	EXPRESSION TAG	UNP P49642
C	-1	THR	-	EXPRESSION TAG	UNP P49642
C	0	SER	-	EXPRESSION TAG	UNP P49642
C	72	ALA	LYS	ENGINEERED MUTATION	UNP P49642
C	73	ALA	MET	ENGINEERED MUTATION	UNP P49642

- Molecule 2 is a protein called DNA PRIMASE LARGE SUBUNIT.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	216	Total	C	H	N	O	S	0	0	0
			3616	1148	1835	307	324	2			
2	D	209	Total	C	H	N	O	S	0	0	0
			3505	1116	1773	297	317	2			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	Zn	0	0
			1	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			14	3	8	3		
4	A	1	Total	C	H	O	0	0
			14	3	8	3		
4	A	1	Total	C	H	O	0	0
			14	3	8	3		
4	A	1	Total	C	H	O	0	0
			14	3	8	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	143	Total	O	0	0
			143	143		
5	B	13	Total	O	0	0
			13	13		
5	C	56	Total	O	0	0
			56	56		
5	D	2	Total	O	0	0
			2	2		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: DNA PRIMASE SMALL SUBUNIT




#### • Molecule 1: DNA PRIMASE SMALL SUBUNIT





#### • Molecule 2: DNA PRIMASE LARGE SUBUNIT





● Molecule 2: DNA PRIMASE LARGE SUBUNIT



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	114.55Å 68.64Å 126.76Å 90.00° 104.36° 90.00°	Depositor
Resolution (Å)	29.62 – 2.70 29.90 – 2.70	Depositor EDS
% Data completeness (in resolution range)	98.5 (29.62-2.70) 98.5 (29.90-2.70)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.58 (at 2.68Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.203 , 0.245 0.201 , 0.245	Depositor DCC
$R_{free}$ test set	2654 reflections (5.10%)	DCC
Wilson B-factor (Å <sup>2</sup> )	81.6	Xtriage
Anisotropy	0.254	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 70.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	20014	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	100.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/3360	0.52	0/4535
1	C	0.32	0/3182	0.47	0/4296
2	B	0.33	0/1811	0.52	0/2430
2	D	0.32	0/1763	0.52	0/2371
All	All	0.34	0/10116	0.50	0/13632

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3229	3221	3167	17	0
1	C	3092	3079	3058	18	0
2	B	1781	1835	1832	19	0
2	D	1732	1773	1770	9	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
4	A	24	32	32	0	0
5	A	143	0	0	3	0
5	B	13	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	56	0	0	6	0
5	D	2	0	0	0	0
All	All	10074	9940	9859	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:ASN:OD1	1:A:87:ARG:NH2	2.20	0.74
1:C:403:ASP:OD1	1:C:406:ARG:NH2	2.26	0.69
1:A:403:ASP:OD1	1:A:406:ARG:NH2	2.26	0.68
2:B:193:ASP:OD2	2:B:220:ASN:ND2	2.27	0.66
1:A:79:ASP:OD1	5:A:2030:HOH:O	2.14	0.65
1:C:148:GLU:OE1	5:C:2034:HOH:O	2.14	0.65
2:D:125:MET:HG2	2:D:219:LEU:HD11	1.80	0.63
1:C:331:SER:OG	5:C:2039:HOH:O	2.15	0.61
2:B:245:ARG:NH1	1:C:91:HIS:O	2.34	0.60
1:C:120:ARG:NH1	1:C:232:ASP:O	2.34	0.59
2:B:99:ASP:OD1	2:B:131:ARG:NH2	2.36	0.55
1:A:162:ARG:NH1	5:A:2077:HOH:O	2.35	0.54
2:D:35:GLU:OE2	2:D:97:ARG:NH1	2.39	0.53
1:C:109:ASP:O	5:C:2023:HOH:O	2.18	0.53
1:C:61:ASN:OD1	1:C:87[A]:ARG:NH2	2.42	0.52
1:C:79:ASP:OD1	5:C:2016:HOH:O	2.19	0.52
2:D:99:ASP:OD1	2:D:131:ARG:NH2	2.42	0.52
2:B:119:TRP:O	2:B:123:GLN:NE2	2.42	0.52
2:B:49:ARG:NH2	2:B:124:GLU:OE2	2.38	0.51
2:D:237:LEU:O	2:D:239:ALA:N	2.41	0.51
1:C:268:ARG:NH2	5:C:2049:HOH:O	2.46	0.49
1:C:200:LYS:NZ	1:C:295:GLU:OE1	2.40	0.49
1:C:54:TYR:OH	5:C:2008:HOH:O	2.17	0.49
1:C:121:CYS:SG	1:C:231:GLN:NE2	2.79	0.48
1:A:237:LYS:HA	1:A:240:TRP:CE2	2.49	0.47
1:A:232:ASP:OD1	1:A:265:SER:OG	2.27	0.47
2:B:151:PHE:CE1	2:B:214:ILE:HD11	2.50	0.47
2:B:111:CYS:HB2	2:B:234:ALA:HB2	1.97	0.46
1:A:104:LYS:NZ	5:A:2031:HOH:O	2.48	0.46
1:A:5:ASP:OD2	1:A:7:THR:OG1	2.33	0.45
2:B:149:LEU:HB3	2:B:151:PHE:CE2	2.51	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:68:GLU:O	2:B:72:SER:N	2.49	0.45
2:D:119:TRP:O	2:D:123:GLN:NE2	2.50	0.45
2:B:237:LEU:O	2:B:239:ALA:N	2.46	0.45
2:B:49:ARG:HB2	2:B:102:SER:HB2	1.98	0.45
1:C:107:VAL:HG21	1:C:317:LEU:HD23	2.00	0.44
2:B:81:LEU:O	2:B:98:ARG:NH2	2.51	0.44
1:A:29:TRP:HB2	1:A:399:LEU:HD21	1.99	0.44
1:C:107:VAL:HG21	1:C:317:LEU:CD2	2.48	0.44
2:B:192:LEU:HA	2:B:195:PHE:CE2	2.53	0.44
2:B:247:GLN:N	2:B:248:PRO:HD2	2.32	0.44
1:C:5:ASP:OD2	1:C:7:THR:OG1	2.35	0.44
1:A:177:LEU:HD22	1:A:181:VAL:HG11	2.00	0.44
1:C:389:ALA:N	1:C:390:PRO:HD2	2.32	0.43
2:D:94:GLU:N	2:D:95:PRO:CD	2.81	0.43
2:D:68:GLU:O	2:D:72:SER:N	2.51	0.43
1:A:48:THR:HB	1:A:77:LYS:HB2	1.99	0.43
1:A:233:ILE:HD12	1:A:243:ILE:HD11	2.01	0.42
2:B:216:ALA:HA	2:B:219:LEU:HB3	2.00	0.42
1:A:125:ALA:CB	1:A:163:ARG:HD2	2.49	0.42
2:D:151:PHE:HD2	2:D:214:ILE:HD11	1.84	0.42
1:A:333:PRO:HD2	1:A:391:TYR:HB3	2.02	0.42
1:A:186:VAL:HG21	1:A:310:SER:HB2	2.02	0.42
2:B:237:LEU:HD12	2:B:238:PRO:HD2	2.01	0.42
2:B:237:LEU:HD23	2:B:240:VAL:HG13	2.03	0.41
2:B:237:LEU:HD23	2:B:240:VAL:CG1	2.51	0.41
2:D:151:PHE:CD2	2:D:214:ILE:HD11	2.55	0.41
1:C:46:SER:OG	1:C:79:ASP:HB2	2.21	0.41
1:A:181:VAL:O	1:A:185:ILE:HG13	2.21	0.41
1:C:269:TRP:CD1	1:C:273:LYS:HD2	2.56	0.41
1:A:210:PRO:O	1:A:214:LYS:HG3	2.21	0.40
2:B:246:LEU:O	2:B:250:LEU:HG	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	384/423 (91%)	368 (96%)	16 (4%)	0	100	100
1	C	364/423 (86%)	353 (97%)	11 (3%)	0	100	100
2	B	208/253 (82%)	198 (95%)	9 (4%)	1 (0%)	32	60
2	D	203/253 (80%)	197 (97%)	6 (3%)	0	100	100
All	All	1159/1352 (86%)	1116 (96%)	42 (4%)	1 (0%)	55	82

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	239	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	363/393 (92%)	360 (99%)	3 (1%)	85	95
1	C	344/393 (88%)	343 (100%)	1 (0%)	94	98
2	B	196/230 (85%)	196 (100%)	0	100	100
2	D	192/230 (84%)	192 (100%)	0	100	100
All	All	1095/1246 (88%)	1091 (100%)	4 (0%)	93	98

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	124	SER
1	A	240	TRP
1	A	282	ASN
1	C	240	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	166	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	A	1410	-	5,5,5	0.35	0	5,5,5	0.33	0
4	GOL	A	1411	-	5,5,5	0.31	0	5,5,5	0.17	0
4	GOL	A	1412	-	5,5,5	0.40	0	5,5,5	0.28	0
4	GOL	A	1413	-	5,5,5	0.38	0	5,5,5	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1410	-	-	0/4/4/4	0/0/0/0
4	GOL	A	1411	-	-	0/4/4/4	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1412	-	-	0/4/4/4	0/0/0/0
4	GOL	A	1413	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	380/423 (89%)	0.45	25 (6%) 19 17	43, 64, 97, 125	0
1	C	367/423 (86%)	1.00	71 (19%) 1 1	58, 95, 135, 161	0
2	B	216/253 (85%)	0.84	31 (14%) 3 2	48, 106, 149, 158	0
2	D	209/253 (82%)	0.99	38 (18%) 1 1	75, 110, 151, 167	0
All	All	1172/1352 (86%)	0.79	165 (14%) 3 2	43, 91, 143, 167	0

All (165) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	195	GLY	7.1
1	C	197	ASP	7.0
2	B	148	GLN	6.9
1	C	322	SER	6.9
1	C	198	VAL	6.6
2	B	61	VAL	6.2
1	C	331	SER	6.2
1	C	320	PRO	6.0
1	C	357	LEU	5.9
2	B	68	GLU	5.8
1	C	319	SER	5.8
2	D	36	ASN	5.6
1	C	125	ALA	5.4
2	B	30	LEU	5.3
1	C	167	CYS	5.1
1	C	354	CYS	5.1
2	D	62	SER	5.0
1	C	4	PHE	5.0
1	C	158	VAL	4.9
2	B	64	VAL	4.9
1	C	355	ARG	4.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	D	63	TYR	4.8
2	B	69	GLN	4.8
2	B	62	SER	4.6
2	D	67	THR	4.4
2	D	34	SER	4.4
1	A	320	PRO	4.4
1	C	321	PHE	4.3
2	B	252	HIS	4.2
2	D	68	GLU	4.2
2	D	64	VAL	4.1
1	C	166	HIS	4.1
1	C	124	SER	4.1
2	B	96	ARG	4.0
2	D	173	LEU	3.9
2	D	172	SER	3.9
2	D	124	GLU	3.9
1	A	301	CYS	3.8
1	C	19	ARG	3.8
2	B	79	ARG	3.7
2	D	26	LEU	3.7
1	C	279	TYR	3.7
2	D	157	GLU	3.7
1	C	168	TRP	3.7
1	C	123	SER	3.6
2	B	95	PRO	3.6
2	D	33	PRO	3.6
1	C	108	PHE	3.6
1	C	127	ILE	3.6
2	B	31	GLN	3.6
1	C	159	TYR	3.6
1	A	319	SER	3.6
2	B	238	PRO	3.6
2	B	151	PHE	3.5
1	A	303	PRO	3.5
2	B	178	LEU	3.5
1	A	300	TYR	3.5
1	A	297	MET	3.4
1	A	407	LYS	3.4
2	D	37	ILE	3.4
2	B	70	TYR	3.3
2	D	167	VAL	3.3
2	D	127	LEU	3.3

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	15	LEU	3.3
1	C	50	LYS	3.3
1	A	331	SER	3.2
1	C	404	LYS	3.2
1	C	407	LYS	3.2
2	D	130	PHE	3.2
1	C	206	GLU	3.2
1	C	318	LYS	3.2
1	C	160	SER	3.1
1	C	121	CYS	3.1
2	D	139	LYS	3.1
1	A	302	PHE	3.1
2	B	180	PHE	3.0
1	C	53	ILE	3.0
1	A	167	CYS	3.0
2	D	28	PHE	3.0
1	C	332	VAL	3.0
1	C	278	ARG	3.0
2	B	236	SER	3.0
1	C	307	ILE	2.9
1	A	168	TRP	2.9
1	C	157	TRP	2.9
2	D	137	LYS	2.9
1	A	298	LEU	2.8
1	A	158	VAL	2.8
1	C	80	ILE	2.8
1	C	317	LEU	2.8
2	D	69	GLN	2.8
1	A	138	ALA	2.8
2	B	73	LYS	2.8
1	C	290	PRO	2.7
2	B	147	SER	2.7
2	D	125	MET	2.7
2	B	58	ASN	2.6
1	C	120	ARG	2.6
2	D	160	THR	2.6
2	D	158	GLU	2.6
1	C	297	MET	2.6
2	B	235	ARG	2.6
1	C	109	ASP	2.6
1	C	27	TYR	2.6
1	C	6	PRO	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	166	HIS	2.5
1	C	8	GLU	2.5
1	A	397[A]	HIS	2.5
1	C	253	ASP	2.5
1	C	301	CYS	2.5
2	D	61	VAL	2.5
1	A	321	PHE	2.5
1	A	18	ARG	2.5
2	D	192	LEU	2.5
1	C	165	VAL	2.5
2	D	235	ARG	2.5
2	B	145	LYS	2.5
2	D	71	GLN	2.4
1	C	107	VAL	2.4
2	B	72	SER	2.4
1	C	323	VAL	2.4
2	D	135	LEU	2.4
2	D	232	LEU	2.4
1	A	80	ILE	2.4
1	A	107	VAL	2.4
1	A	282	ASN	2.4
1	C	403	ASP	2.4
1	C	87[A]	ARG	2.4
1	C	300	TYR	2.3
1	A	53	ILE	2.3
1	A	296	ILE	2.3
2	B	228	LYS	2.3
2	D	32	PRO	2.3
2	D	136	PRO	2.3
1	C	196	GLN	2.3
1	C	386	THR	2.2
2	B	103	HIS	2.2
1	A	26	TYR	2.2
2	B	106	LEU	2.2
2	D	31	GLN	2.2
2	B	118	ARG	2.2
2	B	105	ILE	2.2
1	C	333	PRO	2.2
2	D	161	LEU	2.2
1	C	408	GLY	2.2
1	C	12	LEU	2.2
1	C	9	LEU	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	256	GLN	2.1
2	D	151	PHE	2.1
1	C	122	CYS	2.1
1	A	197	ASP	2.1
2	B	144	LEU	2.1
1	C	11	GLU	2.1
2	D	40	ILE	2.1
1	C	18	ARG	2.1
2	D	126	ASP	2.1
1	C	78	ILE	2.1
1	C	130	LYS	2.1
2	D	138	ASP	2.1
1	C	302	PHE	2.0
2	B	173	LEU	2.0
1	C	26	TYR	2.0
1	C	193	LYS	2.0
1	C	5	ASP	2.0
1	C	199	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	A	1413	6/6	0.91	0.47	7.08	76,91,98,101	0
4	GOL	A	1412	6/6	0.85	0.42	3.37	53,75,97,102	0
4	GOL	A	1410	6/6	0.80	0.37	2.32	62,89,107,108	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ZN	A	1409	1/1	1.00	0.16	-0.14	61,61,61,61	0
4	GOL	A	1411	6/6	0.93	0.19	-0.50	71,85,102,106	0
3	ZN	C	1409	1/1	0.98	0.06	-2.32	104,104,104,104	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.