



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 12, 2017 – 07:59 AM EDT

PDB ID : 5BTL  
Title : Crystal structure of a topoisomerase II complex  
Authors : Blower, T.R.; Williamson, B.H.; Kerns, R.J.; Berger, J.M.  
Deposited on : unknown  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20029824  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824




# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3846 (2.50-2.50)
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)
RSRZ outliers	101464	3876 (2.50-2.50)




The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	503	<div> <div>8%</div> <div>93%</div> <div>• •</div> </div>
1	C	503	<div> <div>11%</div> <div>93%</div> <div>• •</div> </div>
2	B	253	<div> <div>8%</div> <div>92%</div> <div>• •</div> </div>
2	D	253	<div> <div>6%</div> <div>91%</div> <div>7% •</div> </div>
3	E	24	<div> <div>21%</div> <div>71%</div> <div>17% 13%</div> </div>

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	H	24	
4	F	24	
4	G	24	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	PTR	A	129	-	-	X	-
1	PTR	C	129	-	-	X	-



## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 25993 atoms, of which 12526 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA gyrase subunit A.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	487	Total	C	H	N	O	P	S	0	1	0
			7691	2388	3853	704	732	1	13			
1	C	487	Total	C	H	N	O	P	S	0	1	0
			7689	2388	3851	704	732	1	13			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	501	ILE	-	expression tag	UNP P9WG47
A	502	GLY	-	expression tag	UNP P9WG47
A	503	SER	-	expression tag	UNP P9WG47
A	504	GLY	-	expression tag	UNP P9WG47
C	501	ILE	-	expression tag	UNP P9WG47
C	502	GLY	-	expression tag	UNP P9WG47
C	503	SER	-	expression tag	UNP P9WG47
C	504	GLY	-	expression tag	UNP P9WG47

- Molecule 2 is a protein called DNA gyrase subunit B.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	245	Total	C	H	N	O	S	0	0	0
			3906	1217	1974	348	360	7			
2	D	247	Total	C	H	N	O	S	0	0	0
			3930	1224	1984	351	364	7			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	423	SER	-	expression tag	UNP P9WG44
B	424	ASN	-	expression tag	UNP P9WG44
B	425	ALA	-	expression tag	UNP P9WG44
D	423	SER	-	expression tag	UNP P9WG44

*Continued on next page...*



*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	424	ASN	-	expression tag	UNP P9WG44
D	425	ALA	-	expression tag	UNP P9WG44

- Molecule 3 is a DNA chain called DNA substrate 24-mer GGTCATGAATGACTATGCAC GTAA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	E	21	Total	C	H	N	O	P	0	21	0
			620	196	210	74	120	20			
3	H	21	Total	C	H	N	O	P	0	21	0
			615	196	205	74	120	20			

- Molecule 4 is a DNA chain called DNA substrate 24-mer TTACGTGCATAGTCATTTCAT GACC.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	F	21	Total	C	H	N	O	P	0	21	0
			612	196	202	74	120	20			
4	G	21	Total	C	H	N	O	P	0	21	0
			611	196	201	74	120	20			

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Mg	0	0
			1	1		
5	D	1	Total	Mg	0	0
			1	1		
5	F	1	Total	Mg	0	0
			1	1		
5	E	1	Total	Mg	0	0
			1	1		

- Molecule 6 is 1-cyclopropyl-6-fluoro-8-methyl-7-[(4aS,7aS)-octahydro-6H-pyrrolo[3,4-b]pyridin-6-yl]-4-oxo-1,4-dihydroquinoline-3-carboxylic acid (three-letter code: 8MX) (formula: C<sub>21</sub>H<sub>24</sub>FN<sub>3</sub>O<sub>3</sub>).





Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
6	E	1	Total	C	F	H	N	O	0	0
			51	21	1	23	3	3		
6	H	1	Total	C	F	H	N	O	0	0
			51	21	1	23	3	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	95	Total	O	0	0
			95	95		
7	B	32	Total	O	0	0
			32	32		
7	C	38	Total	O	0	0
			38	38		
7	D	28	Total	O	0	0
			28	28		
7	E	7	Total	O	0	0
			7	7		
7	F	6	Total	O	0	0
			6	6		
7	G	1	Total	O	0	0
			1	1		
7	H	6	Total	O	0	0
			6	6		




**i**


- Molecule 1: DNA gyrase subunit A








- Molecule 3: DNA substrate 24-mer GGTCATGAATGACTATGCACGTAA




- Molecule 3: DNA substrate 24-mer GGTCATGAATGACTATGCACGTAA



- Molecule 4: DNA substrate 24-mer TTACGTGCATAGTCATTCATGACC



- Molecule 4: DNA substrate 24-mer TTACGTGCATAGTCATTCATGACC





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	107.94Å 83.05Å 128.74Å 90.00° 108.88° 90.00°	Depositor
Resolution (Å)	49.60 – 2.50 49.11 – 2.50	Depositor EDS
% Data completeness (in resolution range)	95.3 (49.60-2.50) 94.5 (49.11-2.50)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.10 (at 2.51Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.230 , 0.256 0.225 , 0.251	Depositor DCC
$R_{free}$ test set	3547 reflections (5.02%)	DCC
Wilson B-factor (Å <sup>2</sup> )	50.4	Xtriage
Anisotropy	0.290	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 41.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	25993	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 8MX, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.21	0/3881	0.39	0/5253
1	C	0.21	0/3881	0.38	0/5253
2	B	0.23	0/1960	0.39	0/2634
2	D	0.22	0/1974	0.39	0/2653
3	E	0.53	0/458	0.95	0/704
3	H	0.54	0/458	0.96	0/704
4	F	0.53	0/458	0.94	0/704
4	G	0.50	0/458	0.94	0/704
All	All	0.28	0/13528	0.51	0/18609

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3838	3853	3853	18	0
1	C	3838	3851	3853	20	0
2	B	1932	1974	1973	8	0
2	D	1946	1984	1984	14	0
3	E	410	210	210	6	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	H	410	205	205	6	0
4	F	410	202	202	5	0
4	G	410	201	201	8	0
5	B	1	0	0	0	0
5	D	1	0	0	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
6	E	28	23	0	1	0
6	H	28	23	0	2	0
7	A	95	0	0	1	0
7	B	32	0	0	1	0
7	C	38	0	0	1	0
7	D	28	0	0	0	0
7	E	7	0	0	0	0
7	F	6	0	0	0	0
7	G	1	0	0	0	0
7	H	6	0	0	0	0
All	All	13467	12526	12481	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:129:PTR:O2P	4:G:11[B]:DA:C5'	2.03	1.06
1:A:129:PTR:O3P	3:H:11[B]:DG:C5'	2.17	0.93
1:C:129:PTR:O2P	3:E:11[A]:DG:H4'	1.74	0.86
1:A:129:PTR:O3P	3:H:11[B]:DG:H4'	1.78	0.83
2:D:653:GLY:O	2:D:659:ARG:NH2	2.11	0.83
1:A:129:PTR:O3P	4:F:11[A]:DA:H4'	1.80	0.81
1:A:129:PTR:O3P	3:H:11[B]:DG:C4'	2.29	0.80
1:A:287:ILE:O	1:A:291:VAL:HG23	1.82	0.80
4:G:18[B]:DC:O2	3:H:7[B]:DG:N2	2.15	0.79
1:A:129:PTR:O3P	4:F:11[A]:DA:C5'	2.31	0.78
1:C:129:PTR:O2P	3:E:11[A]:DG:C5'	2.32	0.76
4:G:7[B]:DG:N2	3:H:18[B]:DC:O2	2.19	0.73
1:C:129:PTR:O2P	3:E:11[A]:DG:C4'	2.35	0.73
1:C:129:PTR:O2P	4:G:11[B]:DA:H4'	1.95	0.66
2:D:429:ARG:NH1	2:D:439:PRO:O	2.30	0.65
2:B:429:ARG:NH1	2:B:439:PRO:O	2.33	0.61

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31:TYR:HA	2:B:535:VAL:CG1	2.31	0.60
1:C:129:PTR:O2P	4:G:11[B]:DA:C4'	2.39	0.60
1:C:287:ILE:O	1:C:291:VAL:HG23	2.03	0.59
2:B:508:ALA:O	2:B:523:ARG:NE	2.35	0.59
1:C:262:ASP:OD1	1:C:263:SER:N	2.37	0.58
2:D:659:ARG:NH1	4:G:19[B]:DA:OP1	2.37	0.57
1:C:389:HIS:O	1:C:431:GLN:NE2	2.37	0.57
1:C:68:ARG:NH1	7:C:601:HOH:O	2.38	0.56
1:A:129:PTR:O3P	4:F:11[A]:DA:C4'	2.40	0.55
2:B:446:ARG:NH1	2:B:473:SER:OG	2.41	0.54
2:D:659:ARG:NH1	3:E:19[A]:DA:OP1	2.41	0.54
6:H:101:8MX:C01	6:H:101:8MX:C09	2.86	0.53
2:B:634:ARG:NH2	7:B:801:HOH:O	2.36	0.51
1:C:428:ASP:OD1	1:C:429:GLU:N	2.42	0.50
1:A:428:ASP:OD1	1:A:429:GLU:N	2.43	0.49
1:A:31:TYR:HD1	2:B:535:VAL:HG12	1.78	0.49
1:A:441:ARG:NE	1:C:437:ASP:OD1	2.43	0.48
1:A:427:ILE:CG2	1:A:431:GLN:HB2	2.43	0.47
1:A:51:VAL:HG11	3:E:8[A]:DA:H5'	1.96	0.47
2:B:426:LEU:HD21	2:B:523:ARG:HD2	1.97	0.47
1:A:290:GLN:O	1:A:295:LYS:HB2	2.15	0.46
2:D:446:ARG:NH2	2:D:473:SER:OG	2.49	0.46
2:D:511:THR:HG21	2:D:517:PHE:CE2	2.52	0.45
2:D:497:LEU:O	2:D:503:GLN:NE2	2.49	0.44
4:F:16[A]:DT:C6	4:F:17[A]:DT:H72	2.53	0.44
1:C:31:TYR:HA	2:D:535:VAL:CG1	2.47	0.44
1:A:275:PRO:HG2	1:A:278:VAL:HG21	2.00	0.44
4:G:5[B]:DG:N2	3:H:20[B]:DC:O2	2.51	0.44
1:C:65:ARG:HB3	1:C:66:PRO:HD2	2.00	0.43
1:C:31:TYR:HA	2:D:535:VAL:HG13	2.00	0.43
1:C:60:PHE:HB2	1:C:142:LEU:HD13	2.01	0.43
1:C:31:TYR:HD1	2:D:535:VAL:HG12	1.84	0.43
2:D:643:ALA:O	2:D:646:GLU:HG2	2.18	0.43
6:H:101:8MX:C23	6:H:101:8MX:C01	2.97	0.43
6:E:101:8MX:C01	6:E:101:8MX:C23	2.97	0.42
3:E:12[A]:DA:N6	4:F:12[A]:DG:O6	2.53	0.42
2:B:553:ARG:N	2:B:554:PRO:CD	2.83	0.42
1:C:275:PRO:HG2	1:C:278:VAL:HG21	2.01	0.42
2:D:553:ARG:N	2:D:554:PRO:CD	2.83	0.41
1:A:52:HIS:NE2	4:G:8[B]:DC:OP1	2.49	0.41
1:C:186:ALA:O	1:C:344:ASN:ND2	2.53	0.41

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:313:ARG:NH1	7:A:606:HOH:O	2.51	0.41
1:A:424:LEU:HG	1:A:425:LEU:HG	2.03	0.41
1:C:34:SER:OG	2:D:535:VAL:HG11	2.20	0.41
2:D:627:ASP:HB3	2:D:630:VAL:HG22	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	485/503 (96%)	469 (97%)	16 (3%)	0	100	100
1	C	485/503 (96%)	468 (96%)	17 (4%)	0	100	100
2	B	241/253 (95%)	233 (97%)	8 (3%)	0	100	100
2	D	243/253 (96%)	235 (97%)	8 (3%)	0	100	100
All	All	1454/1512 (96%)	1405 (97%)	49 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	407/420 (97%)	407 (100%)	0	100	100
1	C	407/420 (97%)	407 (100%)	0	100	100

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	205/211 (97%)	205 (100%)	0	100	100
2	D	207/211 (98%)	207 (100%)	0	100	100
All	All	1226/1262 (97%)	1226 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
1	PTR	A	129	1	15,16,17	1.28	1 (6%)	19,22,24	0.98	1 (5%)
1	PTR	C	129	1	15,16,17	1.04	1 (6%)	19,22,24	1.06	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	129	1	-	0/9/11/13	0/1/1/1
1	PTR	C	129	1	-	0/9/11/13	0/1/1/1

All (2) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	129	PTR	CA-C	-3.03	1.46	1.50
1	C	129	PTR	CA-C	-2.64	1.46	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	129	PTR	CB-CA-C	-2.84	105.94	111.41
1	A	129	PTR	CB-CA-C	-2.54	106.52	111.41

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	129	PTR	6	0
1	C	129	PTR	6	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	8MX	E	101	5	24,32,32	5.82	12 (50%)	24,49,49	2.14	9 (37%)
6	8MX	H	101	5	24,32,32	5.82	10 (41%)	24,49,49	3.00	14 (58%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical



component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	8MX	E	101	5	-	0/8/33/33	1/4/5/5
6	8MX	H	101	5	-	0/8/33/33	0/4/5/5

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	H	101	8MX	C16-C15	-19.29	1.19	1.52
6	E	101	8MX	C16-C15	-16.89	1.23	1.52
6	H	101	8MX	C09-C10	-15.74	1.29	1.53
6	E	101	8MX	C09-C10	-15.06	1.30	1.53
6	H	101	8MX	C04-C03	-3.54	1.38	1.41
6	E	101	8MX	C04-C03	-3.11	1.39	1.41
6	H	101	8MX	C12-C11	-2.67	1.46	1.53
6	E	101	8MX	C12-C11	-2.18	1.47	1.53
6	H	101	8MX	C07-N08	2.04	1.46	1.40
6	H	101	8MX	C02-C03	2.05	1.46	1.43
6	E	101	8MX	C13-N14	2.07	1.52	1.47
6	E	101	8MX	C07-N08	2.29	1.47	1.40
6	E	101	8MX	C05-C06	2.40	1.39	1.35
6	H	101	8MX	C05-C06	2.63	1.39	1.35
6	E	101	8MX	C09-N08	3.13	1.50	1.46
6	H	101	8MX	C12-C13	3.31	1.65	1.49
6	E	101	8MX	C12-C13	3.77	1.68	1.49
6	E	101	8MX	C15-N14	3.94	1.51	1.47
6	H	101	8MX	C10-C15	7.49	1.64	1.53
6	H	101	8MX	C16-N08	9.08	1.58	1.46
6	E	101	8MX	C10-C15	9.82	1.67	1.53
6	E	101	8MX	C16-N08	11.23	1.61	1.46

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	H	101	8MX	C13-N14-C15	-8.88	105.97	111.62
6	H	101	8MX	C09-N08-C07	-3.90	115.62	123.04
6	E	101	8MX	C05-C04-C18	-3.29	116.32	121.69
6	H	101	8MX	C05-C04-C18	-3.28	116.34	121.69
6	E	101	8MX	C21-C20-C18	-3.13	118.53	119.99
6	E	101	8MX	C09-N08-C07	-3.09	117.16	123.04
6	H	101	8MX	C21-C20-C18	-3.02	118.58	119.99
6	E	101	8MX	C13-N14-C15	-2.61	109.96	111.62

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	E	101	8MX	C06-C07-N08	-2.49	115.19	121.68
6	H	101	8MX	F17-C06-C05	-2.46	116.88	120.31
6	H	101	8MX	C16-N08-C07	-2.21	118.83	123.04
6	H	101	8MX	C05-C06-C07	-2.18	120.21	123.37
6	H	101	8MX	C20-C18-C04	-2.07	118.41	122.09
6	E	101	8MX	C05-C06-C07	-2.00	120.47	123.37
6	H	101	8MX	C10-C09-N08	2.14	106.03	103.30
6	H	101	8MX	C12-C11-C10	2.54	116.06	111.95
6	H	101	8MX	F17-C06-C07	3.19	122.91	118.30
6	H	101	8MX	C01-C02-C03	3.67	123.71	119.44
6	E	101	8MX	C23-N22-C03	3.68	125.56	120.87
6	H	101	8MX	C23-N22-C03	3.68	125.57	120.87
6	E	101	8MX	C18-C04-C03	3.91	122.81	119.90
6	E	101	8MX	C01-C02-C03	4.22	124.35	119.44
6	H	101	8MX	C18-C04-C03	4.27	123.08	119.90

There are no chirality outliers.

There are no torsion outliers.

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	E	101	8MX	C10-C11-C12-C13-C15-N14

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	E	101	8MX	1	0
6	H	101	8MX	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	486/503 (96%)	0.82	41 (8%) 12 11	33, 56, 105, 163	0
1	C	486/503 (96%)	0.91	55 (11%) 6 5	37, 58, 105, 149	0
2	B	245/253 (96%)	0.85	19 (7%) 14 14	42, 64, 97, 130	0
2	D	247/253 (97%)	0.64	14 (5%) 24 25	44, 61, 95, 150	0
3	E	21/24 (87%)	1.90	5 (23%) 1 0	64, 105, 173, 203	21 (100%)
3	H	21/24 (87%)	1.50	7 (33%) 0 0	26, 57, 128, 179	21 (100%)
4	F	21/24 (87%)	1.83	8 (38%) 0 0	59, 98, 176, 209	21 (100%)
4	G	21/24 (87%)	1.89	6 (28%) 1 0	32, 56, 157, 183	21 (100%)
All	All	1548/1608 (96%)	0.88	155 (10%) 8 7	26, 61, 112, 209	84 (5%)

All (155) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	E	23[A]	DA	10.6
4	G	23[B]	DC	9.9
1	C	292	ARG	8.4
1	A	264	ARG	8.1
2	B	426	LEU	6.3
1	C	264	ARG	6.2
1	A	292	ARG	6.0
1	A	452	ILE	5.4
1	A	265	GLY	5.3
2	B	603	LYS	5.1
1	A	401	LEU	5.0
1	A	412	GLU	4.9
1	A	408	ILE	4.7
2	B	601	ILE	4.6
1	C	294	GLY	4.5
1	A	387	ARG	4.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
4	F	23[A]	DC	4.5
4	G	8[B]	DC	4.5
1	C	442	ARG	4.3
2	D	601	ILE	4.2
3	E	8[A]	DA	4.1
1	C	259	VAL	4.1
1	A	263	SER	4.0
1	A	266	ARG	3.9
1	A	501	ILE	3.9
1	C	412	GLU	3.8
3	E	11[A]	DG	3.7
1	C	470	ALA	3.7
4	F	3[A]	DA	3.7
1	A	446	LEU	3.6
2	D	426	LEU	3.5
2	B	498	LYS	3.4
4	F	12[A]	DG	3.4
1	C	501	ILE	3.4
1	A	15	ILE	3.3
1	C	450	ARG	3.2
1	C	265	GLY	3.2
1	C	323	VAL	3.2
1	A	323	VAL	3.1
1	C	414	VAL	3.1
2	D	602	ASN	3.1
1	C	318	ILE	3.1
1	C	295	LYS	3.1
3	H	23[B]	DA	3.1
1	C	126	ALA	3.1
3	H	11[B]	DG	3.1
1	C	394	LEU	3.0
1	C	15	ILE	3.0
2	B	427	VAL	3.0
1	A	325	LYS	3.0
4	G	11[B]	DA	3.0
1	C	463	ALA	3.0
1	A	414	VAL	3.0
4	F	22[A]	DA	3.0
4	F	7[A]	DG	3.0
1	C	429	GLU	2.9
2	B	430	LYS	2.9
1	C	296	LEU	2.9

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	436	LEU	2.9
1	C	123	PRO	2.8
1	A	442	ARG	2.8
2	B	481	LEU	2.8
1	C	387	ARG	2.8
2	B	674	ASP	2.8
1	C	293	ASP	2.8
3	H	7[B]	DG	2.8
1	C	130	THR	2.7
2	B	515	ASP	2.7
1	C	114	GLY	2.7
1	A	434	ALA	2.7
3	H	12[B]	DA	2.7
1	A	438	MET	2.7
1	C	92	ILE	2.6
1	C	116	PHE	2.6
1	C	267	THR	2.6
4	G	12[B]	DG	2.6
1	C	331	LEU	2.6
1	C	128[A]	ARG	2.6
1	C	325	LYS	2.6
1	A	259	VAL	2.5
2	D	492	ARG	2.5
1	A	421	LEU	2.5
1	A	248	ARG	2.5
1	A	92	ILE	2.5
1	A	429	GLU	2.5
3	H	3[B]	DT	2.5
1	C	125	ALA	2.5
2	B	450	PRO	2.5
1	A	91	SER	2.5
1	A	443	LEU	2.5
1	C	404	VAL	2.5
2	B	609	ARG	2.5
4	F	8[A]	DC	2.5
1	A	261	GLU	2.5
1	A	126	ALA	2.5
4	G	7[B]	DG	2.5
1	C	451	ILE	2.5
1	C	322	ALA	2.4
2	B	492	ARG	2.4
2	D	438	LEU	2.4

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	427	ILE	2.4
2	D	497	LEU	2.4
1	C	298	GLY	2.4
3	E	7[A]	DG	2.4
1	A	436	LEU	2.4
1	C	419	ALA	2.4
1	C	416	ILE	2.4
1	A	430	ILE	2.3
1	C	438	MET	2.3
2	B	467	ALA	2.3
2	B	462	SER	2.3
1	A	404	VAL	2.3
2	B	475	PHE	2.3
1	A	410	ALA	2.3
2	B	612	GLY	2.3
4	F	11[A]	DA	2.3
1	C	261	GLU	2.3
1	C	423	GLU	2.3
2	D	498	LYS	2.2
2	D	475	PHE	2.2
1	C	266	ARG	2.2
1	C	407	LEU	2.2
1	C	93	TYR	2.2
3	H	8[B]	DA	2.2
1	C	452	ILE	2.2
1	C	430	ILE	2.2
1	A	86	PRO	2.2
2	D	600	LYS	2.2
2	D	603	LYS	2.2
4	F	21[A]	DG	2.2
4	G	22[B]	DA	2.2
1	C	302	ILE	2.2
2	D	675	VAL	2.2
1	C	440	LEU	2.2
1	A	470	ALA	2.1
1	C	326	VAL	2.1
1	C	262	ASP	2.1
1	C	94	ASP	2.1
1	A	51	VAL	2.1
2	D	427	VAL	2.1
1	C	316	ILE	2.1
2	B	497	LEU	2.1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	302	ILE	2.1
2	B	506	ILE	2.1
3	H	22[B]	DT	2.1
3	E	12[A]	DA	2.1
2	D	609	ARG	2.1
1	A	123	PRO	2.1
1	A	48	LEU	2.1
1	A	130	THR	2.0
1	C	315	VAL	2.0
2	B	437	GLY	2.0
1	A	395	VAL	2.0
2	D	511	THR	2.0
1	C	335	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	PTR	A	129	16/17	0.95	0.24	-	45,50,70,72	0
1	PTR	C	129	16/17	0.94	0.31	-	48,52,67,78	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	MG	F	101	1/1	0.98	0.33	1.47	47,47,47,47	0
5	MG	E	102	1/1	0.95	0.33	1.03	65,65,65,65	0
6	8MX	E	101	28/28	0.83	0.29	0.60	41,50,65,66	0
6	8MX	H	101	28/28	0.89	0.25	-0.10	39,47,56,56	0
5	MG	D	701	1/1	0.97	0.18	-1.27	40,40,40,40	0
5	MG	B	701	1/1	0.94	0.20	-1.76	48,48,48,48	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.