



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 07:59 pm GMT

PDB ID : 3BUV  
Title : Crystal structure of human Delta(4)-3-ketosteroid 5-beta-reductase in complex with NADP and HEPES. Resolution: 1.35 Å.  
Authors : Di Costanzo, L.; Drury, J.; Penning, T.M.; Christianson, D.W.  
Deposited on : 2008-01-03  
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1024 (1.38-1.34)
Clashscore	112137	1063 (1.38-1.34)
Ramachandran outliers	110173	1048 (1.38-1.34)
Sidechain outliers	110143	1048 (1.38-1.34)
RSRZ outliers	101464	1025 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	<div> <div>14%</div> <div>85%</div> <div>14%</div> </div>
1	B	326	<div> <div>6%</div> <div>88%</div> <div>11%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EPE	A	3999	-	-	-	X
4	GOL	B	329	-	-	-	X
4	GOL	B	330	-	-	-	X

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 5933 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-oxo-5-beta-steroid 4-dehydrogenase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	325	Total	C	N	O	S	0	0	0
			2627	1680	455	481	11			
1	A	325	Total	C	N	O	S	0	0	0
			2627	1680	455	481	11			

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C<sub>21</sub>H<sub>28</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C<sub>8</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	252	Total 252	O 252	0	0
5	B	304	Total 304	O 304	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3-oxo-5-beta-steroid 4-dehydrogenase



- Molecule 1: 3-oxo-5-beta-steroid 4-dehydrogenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.71Å 109.80Å 128.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.69 – 1.35 33.69 – 1.35	Depositor EDS
% Data completeness (in resolution range)	91.0 (33.69-1.35) 95.5 (33.69-1.35)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.14 (at 1.35Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.218 , 0.232 0.215 , 0.217	Depositor DCC
$R_{free}$ test set	7400 reflections (5.02%)	DCC
Wilson B-factor (Å <sup>2</sup> )	14.1	Xtriage
Anisotropy	0.590	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 43.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5933	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.62% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAP, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.28	0/2692	0.54	0/3649
1	B	0.30	0/2692	0.58	0/3649
All	All	0.29	0/5384	0.56	0/7298

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2627	0	2622	31	0
1	B	2627	0	2622	31	0
2	A	48	0	25	1	0
2	B	48	0	25	2	0
3	A	15	0	17	4	0
4	B	12	0	16	1	0
5	A	252	0	0	3	0
5	B	304	0	0	5	0
All	All	5933	0	5327	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:3999:EPE:N1	3:A:3999:EPE:C2	1.70	1.48
3:A:3999:EPE:C2	3:A:3999:EPE:C6	2.44	0.95
3:A:3999:EPE:C9	3:A:3999:EPE:C2	2.56	0.84
1:B:114:VAL:HG22	1:B:163:VAL:HG22	1.57	0.84
1:B:102:LEU:HD21	1:B:163:VAL:HG23	1.61	0.81
1:A:232:ASN:HD21	1:A:234:SER:HB3	1.50	0.76
1:A:263:ASN:O	1:A:266:ARG:HG2	1.85	0.75
1:B:114:VAL:CG2	1:B:163:VAL:HG22	2.19	0.72
1:B:152:GLU:HG3	1:B:182:LYS:HE3	1.70	0.71
1:A:263:ASN:OD1	1:A:266:ARG:HD3	1.92	0.68
3:A:3999:EPE:C3	3:A:3999:EPE:N1	2.59	0.64
1:A:232:ASN:ND2	1:A:234:SER:HB3	2.12	0.63
1:A:185:LEU:HD21	1:A:188:LYS:HG2	1.80	0.63
1:A:206:LEU:O	1:A:210:GLN:HG3	2.00	0.62
1:B:188:LYS:HE2	5:B:3958:HOH:O	2.03	0.58
1:A:122:PRO:HG3	1:A:321:PRO:HG3	1.86	0.57
1:A:122:PRO:CG	1:A:321:PRO:HG3	2.35	0.56
1:B:119:ILE:HG12	1:B:154:MET:HE2	1.88	0.54
1:B:243:LEU:HD23	1:B:302:LEU:HD21	1.89	0.53
1:B:80:GLU:CD	1:B:80:GLU:H	2.12	0.53
1:B:191:SER:HB2	5:B:4025:HOH:O	2.08	0.53
1:A:266:ARG:HG3	1:A:268:VAL:HG23	1.89	0.52
1:B:119:ILE:HG12	1:B:154:MET:CE	2.39	0.52
1:A:273:LYS:O	2:A:3902:NAP:H8A	2.09	0.52
1:A:307:ARG:HG3	1:A:310:GLU:HG2	1.91	0.51
1:B:71:LYS:HA	1:B:74:GLU:HG2	1.92	0.51
1:A:207:LYS:O	1:A:211:GLN:HG3	2.09	0.51
1:B:273:LYS:O	2:B:3901:NAP:H8A	2.11	0.51
1:B:232:ASN:ND2	1:B:234:SER:OG	2.44	0.50
1:A:185:LEU:CD2	1:A:188:LYS:HG2	2.42	0.50
1:A:273:LYS:HD3	1:A:273:LYS:C	2.32	0.50
1:A:99:ARG:HB3	1:A:100:PRO:HD3	1.93	0.50
1:A:293:GLU:HG3	1:A:297:LYS:HE3	1.94	0.49
1:B:99:ARG:HB3	1:B:100:PRO:HD3	1.95	0.49
1:B:207:LYS:O	1:B:211:GLN:HG3	2.13	0.49
1:A:256:ALA:O	1:A:259:VAL:HG22	2.11	0.49
1:A:118:ILE:HD12	1:A:169:SER:HB2	1.96	0.48
1:A:56:TYR:CD1	1:A:104:ARG:HD3	2.49	0.48
1:B:188:LYS:NZ	1:B:188:LYS:HB3	2.28	0.48
1:A:131:ILE:C	1:A:133:PRO:HD3	2.35	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:88:LEU:HD22	1:B:154:MET:HE2	1.97	0.47
1:B:62:HIS:HE1	5:B:4013:HOH:O	1.97	0.46
1:B:100:PRO:HB2	4:B:330:GOL:H11	1.97	0.46
1:B:203:PRO:HG2	5:B:3986:HOH:O	2.15	0.46
1:A:80:GLU:CD	1:A:80:GLU:H	2.18	0.46
1:B:273:LYS:HD3	1:B:273:LYS:C	2.36	0.46
1:A:292:THR:OG1	1:A:295:GLU:HG3	2.17	0.45
1:A:196:CYS:HB3	1:A:218:ALA:CB	2.47	0.45
1:A:220:SER:N	1:A:221:PRO:CD	2.79	0.45
1:B:281:LYS:O	1:B:285:GLN:HG2	2.17	0.44
1:A:151:TRP:HD1	1:A:182:LYS:NZ	2.16	0.43
1:A:236:PRO:HG2	5:A:4018:HOH:O	2.17	0.43
1:A:236:PRO:HA	1:A:237:PRO:HD3	1.87	0.42
1:B:62:HIS:HD2	5:A:4004:HOH:O	2.01	0.42
1:A:177:GLU:HG2	5:A:4139:HOH:O	2.18	0.42
1:B:232:ASN:HB2	5:B:4070:HOH:O	2.19	0.42
1:A:89:TRP:CG	1:A:90:ALA:N	2.87	0.42
1:B:226:ARG:HA	1:B:231:VAL:HG21	2.02	0.42
1:A:320:TYR:HA	1:A:321:PRO:HD3	1.87	0.41
1:B:220:SER:N	1:B:221:PRO:CD	2.82	0.41
1:B:193:GLN:OE1	2:B:3901:NAP:H2N	2.20	0.41
1:B:166:LEU:O	1:B:190:VAL:HG22	2.20	0.41
1:A:244:LEU:HD13	1:A:259:VAL:CG1	2.51	0.41
1:B:89:TRP:CG	1:B:90:ALA:N	2.88	0.41
1:B:19:PRO:HB2	1:B:51:HIS:HB2	2.03	0.41
1:B:71:LYS:HA	1:B:71:LYS:HD3	1.93	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	323/326 (99%)	317 (98%)	6 (2%)	0	100	100
1	B	323/326 (99%)	317 (98%)	6 (2%)	0	100	100
All	All	646/652 (99%)	634 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	289/290 (100%)	286 (99%)	3 (1%)	80	51
1	B	289/290 (100%)	287 (99%)	2 (1%)	87	65
All	All	578/580 (100%)	573 (99%)	5 (1%)	82	55

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	219	TYR
1	B	232	ASN
1	A	146	ASN
1	A	219	TYR
1	A	323	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	62	HIS
1	B	92	ASN
1	B	232	ASN
1	A	92	ASN
1	A	110	GLN
1	A	146	ASN
1	A	211	GLN
1	A	232	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	323	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAP	A	3902	-	44,52,52	2.31	14 (31%)	51,80,80	2.33	13 (25%)
3	EPE	A	3999	-	15,15,15	3.69	6 (40%)	18,20,20	4.76	8 (44%)
4	GOL	B	329	-	5,5,5	0.48	0	5,5,5	0.63	0
4	GOL	B	330	-	5,5,5	0.49	0	5,5,5	0.66	0
2	NAP	B	3901	-	44,52,52	2.32	15 (34%)	51,80,80	2.35	12 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	3902	-	-	0/27/67/67	0/5/5/5
3	EPE	A	3999	-	-	0/9/19/19	0/1/1/1
4	GOL	B	329	-	-	0/4/4/4	0/0/0/0
4	GOL	B	330	-	-	0/4/4/4	0/0/0/0
2	NAP	B	3901	-	-	0/27/67/67	0/5/5/5

All (35) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	3999	EPE	C10-S	-8.28	1.65	1.77
3	A	3999	EPE	O1S-S	-4.96	1.30	1.45
2	A	3902	NAP	O5B-C5B	-4.54	1.26	1.44
2	B	3901	NAP	O5B-C5B	-4.36	1.27	1.44
3	A	3999	EPE	C7-N4	-4.13	1.37	1.47
2	A	3902	NAP	PN-O5D	-3.08	1.46	1.59
2	B	3901	NAP	PN-O5D	-3.06	1.46	1.59
2	B	3901	NAP	PN-O1N	-2.94	1.39	1.50
2	A	3902	NAP	PN-O1N	-2.92	1.40	1.50
2	B	3901	NAP	PA-O1A	-2.52	1.41	1.50
2	A	3902	NAP	PA-O1A	-2.43	1.41	1.50
3	A	3999	EPE	O2S-S	-2.09	1.39	1.45
2	B	3901	NAP	C3D-C4D	2.01	1.58	1.53
2	B	3901	NAP	C3N-C7N	2.06	1.53	1.50
2	A	3902	NAP	C3N-C7N	2.22	1.54	1.50
2	B	3901	NAP	C2A-N3A	2.30	1.36	1.32
2	A	3902	NAP	C2A-N3A	2.51	1.36	1.32
2	A	3902	NAP	C2A-N1A	2.72	1.39	1.33
2	B	3901	NAP	C2A-N1A	2.75	1.39	1.33
2	A	3902	NAP	O4B-C1B	2.79	1.45	1.41
3	A	3999	EPE	C5-N4	2.92	1.54	1.47
2	B	3901	NAP	O4D-C1D	2.99	1.45	1.41
2	A	3902	NAP	O4D-C1D	3.05	1.45	1.41
2	B	3901	NAP	C2N-C3N	3.30	1.44	1.39
2	B	3901	NAP	O4B-C1B	3.51	1.46	1.41
2	A	3902	NAP	C4N-C3N	3.53	1.45	1.39
2	B	3901	NAP	C4N-C3N	3.63	1.45	1.39
2	A	3902	NAP	C2N-C3N	3.69	1.44	1.39
2	B	3901	NAP	C5N-C4N	3.91	1.46	1.38
2	A	3902	NAP	C5N-C4N	4.00	1.46	1.38
2	A	3902	NAP	C6N-N1N	5.88	1.50	1.35
2	B	3901	NAP	C6N-N1N	5.99	1.50	1.35
2	A	3902	NAP	C4A-N3A	6.82	1.45	1.35
2	B	3901	NAP	C4A-N3A	6.93	1.45	1.35

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	3999	EPE	C2-N1	8.46	1.70	1.47

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3901	NAP	N3A-C2A-N1A	-10.66	119.57	128.86
2	A	3902	NAP	N3A-C2A-N1A	-10.32	119.87	128.86
2	A	3902	NAP	C3N-C7N-N7N	-5.34	111.68	117.77
2	B	3901	NAP	C3N-C7N-N7N	-5.20	111.83	117.77
3	A	3999	EPE	O3S-S-O1S	-4.67	100.66	111.37
3	A	3999	EPE	C6-N1-C2	-3.35	101.29	108.87
3	A	3999	EPE	O2S-S-O1S	-2.55	105.01	113.86
2	A	3902	NAP	O4D-C4D-C3D	-2.55	100.10	105.17
3	A	3999	EPE	O3S-S-O2S	-2.43	105.80	111.37
2	B	3901	NAP	O4D-C4D-C3D	-2.40	100.40	105.17
3	A	3999	EPE	C5-N4-C3	-2.30	103.67	108.87
2	B	3901	NAP	C6N-C5N-C4N	-2.21	116.10	119.44
2	A	3902	NAP	C6N-C5N-C4N	-2.10	116.27	119.44
2	A	3902	NAP	O5B-C5B-C4B	2.02	116.15	109.00
2	A	3902	NAP	C2N-C3N-C4N	2.10	120.65	118.26
2	B	3901	NAP	C2N-C3N-C4N	2.13	120.69	118.26
2	A	3902	NAP	O5D-C5D-C4D	2.53	117.95	109.00
2	B	3901	NAP	O5D-C5D-C4D	2.58	118.15	109.00
2	A	3902	NAP	C4A-C5A-N7A	2.61	111.93	109.41
2	B	3901	NAP	C4A-C5A-N7A	2.70	112.02	109.41
2	B	3901	NAP	C2A-N1A-C6A	2.82	123.70	118.77
2	A	3902	NAP	C2A-N1A-C6A	2.83	123.72	118.77
2	B	3901	NAP	O3D-C3D-C2D	3.17	121.98	111.83
2	A	3902	NAP	O3D-C3D-C2D	3.21	122.11	111.83
2	B	3901	NAP	C4B-O4B-C1B	3.35	113.33	109.77
2	A	3902	NAP	C4B-O4B-C1B	3.51	113.50	109.77
3	A	3999	EPE	O2S-S-C10	3.73	109.99	106.79
2	B	3901	NAP	C5A-C6A-N6A	3.89	128.41	120.47
2	A	3902	NAP	C5A-C6A-N6A	3.96	128.54	120.47
2	A	3902	NAP	O7N-C7N-N7N	4.51	129.00	122.58
2	B	3901	NAP	O7N-C7N-N7N	4.56	129.07	122.58
3	A	3999	EPE	O3S-S-C10	5.03	112.24	106.06
3	A	3999	EPE	O1S-S-C10	17.40	121.73	106.79

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	3902	NAP	1	0
3	A	3999	EPE	4	0
4	B	330	GOL	1	0
2	B	3901	NAP	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	325/326 (99%)	0.82	45 (13%) 3 4	10, 17, 36, 71	0
1	B	325/326 (99%)	0.33	19 (5%) 24 29	10, 14, 26, 41	0
All	All	650/652 (99%)	0.57	64 (9%) 8 12	10, 15, 33, 71	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	30	LYS	7.6
1	B	232	ASN	6.9
1	A	320	TYR	6.5
1	B	234	SER	5.8
1	A	2	ASP	5.7
1	A	178	LEU	5.2
1	B	235	SER	5.2
1	A	136	GLU	5.1
1	A	323	HIS	4.8
1	A	138	GLY	4.8
1	B	233	VAL	4.5
1	A	129	ASP	3.9
1	B	277	LEU	3.7
1	A	140	TRP	3.5
1	A	229	ILE	3.5
1	A	316	ASP	3.4
1	B	144	LYS	3.4
1	B	138	GLY	3.3
1	A	183	PRO	3.3
1	A	142	TYR	3.3
1	A	137	ASN	3.3
1	A	118	ILE	3.1
1	A	147	LEU	3.1
1	A	188	LYS	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	136	GLU	2.9
1	A	139	LYS	2.9
1	A	26	TYR	2.9
1	B	231	VAL	2.9
1	A	233	VAL	2.9
1	A	234	SER	2.8
1	B	52	ILE	2.8
1	A	204	LYS	2.7
1	B	30	LYS	2.7
1	A	293	GLU	2.6
1	B	2	ASP	2.6
1	A	141	LEU	2.5
1	A	134	ARG	2.5
1	B	143	HIS	2.5
1	A	173	ARG	2.5
1	A	145	SER	2.5
1	A	315	ARG	2.4
1	A	126	LYS	2.4
1	A	85	CYS	2.4
1	A	297	LYS	2.4
1	A	194	VAL	2.4
1	A	271	ILE	2.3
1	A	116	LEU	2.3
1	B	139	LYS	2.3
1	B	271	ILE	2.3
1	A	131	ILE	2.3
1	B	85	CYS	2.3
1	B	73	ALA	2.2
1	A	179	ILE	2.2
1	A	177	GLU	2.2
1	A	294	GLU	2.1
1	A	325	GLU	2.1
1	A	133	PRO	2.1
1	B	269	VAL	2.1
1	A	228	PRO	2.1
1	A	112	ASP	2.1
1	A	184	GLY	2.1
1	B	118	ILE	2.1
1	A	31	SER	2.0
1	A	78	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	B	330	6/6	0.76	0.76	14.66	46,49,50,51	6
4	GOL	B	329	6/6	0.67	0.64	13.47	36,37,39,42	6
3	EPE	A	3999	15/15	0.89	0.17	3.00	30,34,35,35	0
2	NAP	B	3901	48/48	0.96	0.10	-0.02	9,12,14,14	0
2	NAP	A	3902	48/48	0.95	0.09	-0.33	11,12,14,15	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.