



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 13, 2017 – 08:05 PM EST

PDB ID : 6BU6
Title : Crystal Structure of the Human vaccinia-related kinase bound to a bis-difluorophenol-aminopyridine inhibitor
Authors : Counago, R.M.; dos Reis, C.V.; de Souza, G.P.; Santiago, A.S.; Azevedo, A.; Guimaraes, C.; Mascarello, A.; Gama, F.; Ferreira, M.; Massirer, K.B.; Arruda, P.; Edwards, A.M.; Elkins, J.M.; Structural Genomics Consortium (SGC)
Deposited on : unknown
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	<div> <div>4%</div> <div>86%</div> <div>12%</div> </div>
1	B	364	<div> <div>10%</div> <div>82%</div> <div>16%</div> </div>
1	C	364	<div> <div>4%</div> <div>84%</div> <div>13%</div> </div>
1	D	364	<div> <div>7%</div> <div>85%</div> <div>13%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	SO4	C	402	-	-	-	X
5	SO4	D	403	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11259 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase VRK1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	322	Total	C	N	O	S	0	6	0
			2571	1643	435	476	17			
1	B	307	Total	C	N	O	S	0	1	0
			2421	1551	418	441	11			
1	C	315	Total	C	N	O	S	0	3	0
			2506	1603	428	461	14			
1	D	318	Total	C	N	O	S	0	1	0
			2480	1585	424	458	13			

There are 52 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP Q99986
A	2	MET	-	expression tag	UNP Q99986
A	34	ALA	LYS	engineered mutation	UNP Q99986
A	35	ALA	LYS	engineered mutation	UNP Q99986
A	36	ALA	GLU	engineered mutation	UNP Q99986
A	212	ALA	GLU	engineered mutation	UNP Q99986
A	214	ALA	LYS	engineered mutation	UNP Q99986
A	215	ALA	GLU	engineered mutation	UNP Q99986
A	292	ALA	GLU	engineered mutation	UNP Q99986
A	293	ALA	LYS	engineered mutation	UNP Q99986
A	295	ALA	LYS	engineered mutation	UNP Q99986
A	359	ALA	LYS	engineered mutation	UNP Q99986
A	360	ALA	LYS	engineered mutation	UNP Q99986
B	1	SER	-	expression tag	UNP Q99986
B	2	MET	-	expression tag	UNP Q99986
B	34	ALA	LYS	engineered mutation	UNP Q99986
B	35	ALA	LYS	engineered mutation	UNP Q99986
B	36	ALA	GLU	engineered mutation	UNP Q99986
B	212	ALA	GLU	engineered mutation	UNP Q99986
B	214	ALA	LYS	engineered mutation	UNP Q99986
B	215	ALA	GLU	engineered mutation	UNP Q99986

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	292	ALA	GLU	engineered mutation	UNP Q99986
B	293	ALA	LYS	engineered mutation	UNP Q99986
B	295	ALA	LYS	engineered mutation	UNP Q99986
B	359	ALA	LYS	engineered mutation	UNP Q99986
B	360	ALA	LYS	engineered mutation	UNP Q99986
C	1	SER	-	expression tag	UNP Q99986
C	2	MET	-	expression tag	UNP Q99986
C	34	ALA	LYS	engineered mutation	UNP Q99986
C	35	ALA	LYS	engineered mutation	UNP Q99986
C	36	ALA	GLU	engineered mutation	UNP Q99986
C	212	ALA	GLU	engineered mutation	UNP Q99986
C	214	ALA	LYS	engineered mutation	UNP Q99986
C	215	ALA	GLU	engineered mutation	UNP Q99986
C	292	ALA	GLU	engineered mutation	UNP Q99986
C	293	ALA	LYS	engineered mutation	UNP Q99986
C	295	ALA	LYS	engineered mutation	UNP Q99986
C	359	ALA	LYS	engineered mutation	UNP Q99986
C	360	ALA	LYS	engineered mutation	UNP Q99986
D	1	SER	-	expression tag	UNP Q99986
D	2	MET	-	expression tag	UNP Q99986
D	34	ALA	LYS	engineered mutation	UNP Q99986
D	35	ALA	LYS	engineered mutation	UNP Q99986
D	36	ALA	GLU	engineered mutation	UNP Q99986
D	212	ALA	GLU	engineered mutation	UNP Q99986
D	214	ALA	LYS	engineered mutation	UNP Q99986
D	215	ALA	GLU	engineered mutation	UNP Q99986
D	292	ALA	GLU	engineered mutation	UNP Q99986
D	293	ALA	LYS	engineered mutation	UNP Q99986
D	295	ALA	LYS	engineered mutation	UNP Q99986
D	359	ALA	LYS	engineered mutation	UNP Q99986
D	360	ALA	LYS	engineered mutation	UNP Q99986

- Molecule 2 is 4,4'-(2-aminopyridine-3,5-diyl)bis(2,6-difluorophenol) (three-letter code: E8V) (formula: C₁₇H₁₀F₄N₂O₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			25	17	4	2	2		
2	B	1	Total	C	F	N	O	0	0
			25	17	4	2	2		
2	D	1	Total	C	F	N	O	0	0
			25	17	4	2	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Cl	0	0
			2	2		
4	A	2	Total	Cl	0	0
			2	2		
4	D	1	Total	Cl	0	0
			1	1		
4	C	1	Total	Cl	0	0
			1	1		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	325	Total 325	O 325	0	0
6	B	244	Total 244	O 244	0	0
6	C	311	Total 311	O 311	0	0
6	D	289	Total 289	O 289	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine/threonine-protein kinase VRK1




- Molecule 1: Serine/threonine-protein kinase VRK1




- Molecule 1: Serine/threonine-protein kinase VRK1



- Molecule 1: Serine/threonine-protein kinase VRK1





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	92.22Å 96.57Å 193.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.98 – 1.80 19.98 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (19.98-1.80) 100.0 (19.98-1.80)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.81 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.184 , 0.212 0.193 , 0.218	Depositor DCC
R_{free} test set	8093 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	26.3	Xtriage
Anisotropy	0.641	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 53.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.015 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11259	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: E8V, GOL, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.63	0/2641	0.78	0/3578
1	B	0.60	0/2481	0.74	0/3355
1	C	0.65	0/2571	0.77	0/3482
1	D	0.67	0/2542	0.75	0/3448
All	All	0.64	0/10235	0.76	0/13863

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2571	0	2497	6	0
1	B	2421	0	2348	5	0
1	C	2506	0	2433	5	0
1	D	2480	0	2375	6	0
2	A	25	0	0	2	0
2	B	25	0	0	1	0
2	D	25	0	0	0	0
3	A	6	0	8	1	0
4	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	2	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	B	5	0	0	0	0
5	C	10	0	0	0	0
5	D	10	0	0	0	0
6	A	325	0	0	1	0
6	B	244	0	0	1	0
6	C	311	0	0	2	0
6	D	289	0	0	1	0
All	All	11259	0	9661	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:120:ASP:HA	1:B:125:SER:HB3	1.76	0.67
1:C:303[A]:MET:SD	6:C:783:HOH:O	2.57	0.57
1:B:103[B]:ARG:NH1	6:B:504:HOH:O	2.40	0.55
1:A:200:LEU:HD13	1:A:221:HIS:CG	2.42	0.54
1:A:48:PHE:HB3	2:A:401:E8V:F2	1.99	0.53
1:A:131[B]:MET:HE1	2:A:401:E8V:F1	2.00	0.52
1:D:48:PHE:HB2	1:D:51:ILE:CD1	2.40	0.51
1:A:134:PHE:CE1	3:A:402:GOL:H31	2.46	0.51
1:C:200:LEU:HD13	1:C:221:HIS:CG	2.47	0.50
1:C:87:TYR:HB3	1:C:113:TYR:HB2	1.94	0.49
1:A:87:TYR:HB3	1:A:113:TYR:HB2	1.95	0.47
1:D:277:ARG:NH1	6:D:509:HOH:O	2.48	0.46
1:B:44:GLY:HA3	2:B:401:E8V:C11	2.46	0.46
1:B:84:LEU:HD23	1:B:129:MET:SD	2.57	0.45
1:C:267:ASP:OD1	1:C:269:LYS:HG2	2.18	0.44
1:D:48:PHE:HB2	1:D:51:ILE:HD11	1.99	0.44
1:C:314:LYS:NZ	6:C:508:HOH:O	2.49	0.43
1:D:184:LEU:HD22	1:D:196:VAL:HG11	2.01	0.43
1:A:303[A]:MET:CE	6:A:535:HOH:O	2.67	0.42
1:B:184:LEU:HD22	1:B:196:VAL:HG11	2.01	0.42
1:D:264:ASN:HB3	1:D:270:TYR:CD2	2.55	0.42
1:D:204:TYR:CE2	1:D:205:CYS:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/364 (90%)	318 (98%)	8 (2%)	0	100	100
1	B	300/364 (82%)	293 (98%)	7 (2%)	0	100	100
1	C	314/364 (86%)	305 (97%)	9 (3%)	0	100	100
1	D	315/364 (86%)	309 (98%)	6 (2%)	0	100	100
All	All	1255/1456 (86%)	1225 (98%)	30 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/304 (88%)	263 (99%)	3 (1%)	78	72
1	B	244/304 (80%)	241 (99%)	3 (1%)	75	69
1	C	257/304 (84%)	254 (99%)	3 (1%)	75	69
1	D	249/304 (82%)	248 (100%)	1 (0%)	93	91
All	All	1016/1216 (84%)	1006 (99%)	10 (1%)	87	75

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	103[A]	ARG
1	A	103[B]	ARG
1	A	133	ARG
1	B	103[A]	ARG
1	B	103[B]	ARG
1	B	133	ARG
1	C	103[A]	ARG
1	C	103[B]	ARG
1	C	133	ARG
1	D	133	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 6 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	E8V	A	401	-	27,27,27	0.56	0	38,40,40	1.43	7 (18%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	402	-	5,5,5	0.53	0	5,5,5	1.18	1 (20%)
2	E8V	B	401	-	27,27,27	0.42	0	38,40,40	1.44	8 (21%)
5	SO4	B	402	-	4,4,4	0.55	0	6,6,6	0.27	0
5	SO4	C	401	-	4,4,4	0.28	0	6,6,6	0.83	0
5	SO4	C	402	-	4,4,4	0.59	0	6,6,6	0.47	0
2	E8V	D	401	-	27,27,27	0.44	0	38,40,40	1.27	5 (13%)
5	SO4	D	402	-	4,4,4	0.60	0	6,6,6	0.59	0
5	SO4	D	403	-	4,4,4	0.43	0	6,6,6	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	E8V	A	401	-	-	0/8/8/8	0/3/3/3
3	GOL	A	402	-	-	0/4/4/4	0/0/0/0
2	E8V	B	401	-	-	0/8/8/8	0/3/3/3
5	SO4	B	402	-	-	0/0/0/0	0/0/0/0
5	SO4	C	401	-	-	0/0/0/0	0/0/0/0
5	SO4	C	402	-	-	0/0/0/0	0/0/0/0
2	E8V	D	401	-	-	0/8/8/8	0/3/3/3
5	SO4	D	402	-	-	0/0/0/0	0/0/0/0
5	SO4	D	403	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	E8V	C8-C9-C10	-3.11	121.41	123.78
2	B	401	E8V	C8-C9-C10	-3.08	121.43	123.78
2	A	401	E8V	C13-C14-C15	-2.95	121.53	123.78
2	A	401	E8V	C17-C16-C15	-2.91	121.56	123.78
2	B	401	E8V	C17-C16-C15	-2.80	121.64	123.78
2	D	401	E8V	C8-C9-C10	-2.73	121.69	123.78
2	D	401	E8V	C13-C14-C15	-2.70	121.72	123.78
2	B	401	E8V	C12-C11-C10	-2.69	121.72	123.78
2	B	401	E8V	C13-C14-C15	-2.66	121.75	123.78
2	A	401	E8V	C12-C11-C10	-2.59	121.80	123.78
2	D	401	E8V	C12-C11-C10	-2.46	121.90	123.78
3	A	402	GOL	O3-C3-C2	2.12	120.76	110.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	E8V	F4-C14-C15	2.34	118.82	116.94
2	B	401	E8V	F3-C16-C15	2.80	119.19	116.94
2	A	401	E8V	F4-C14-C15	2.90	119.26	116.94
2	D	401	E8V	F2-C11-C10	3.03	119.37	116.94
2	A	401	E8V	F3-C16-C15	3.31	119.59	116.94
2	A	401	E8V	F1-C9-C10	3.31	119.59	116.94
2	B	401	E8V	F1-C9-C10	3.34	119.61	116.94
2	D	401	E8V	F1-C9-C10	3.62	119.84	116.94
2	B	401	E8V	F2-C11-C10	3.74	119.93	116.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	E8V	2	0
3	A	402	GOL	1	0
2	B	401	E8V	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	322/364 (88%)	0.02	13 (4%) 39 33	19, 30, 47, 73	0
1	B	307/364 (84%)	0.40	35 (11%) 6 4	18, 34, 74, 98	0
1	C	315/364 (86%)	0.08	14 (4%) 35 30	17, 30, 57, 72	0
1	D	318/364 (87%)	0.25	27 (8%) 11 9	15, 30, 68, 116	0
All	All	1262/1456 (86%)	0.19	89 (7%) 17 13	15, 31, 66, 116	0

All (89) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	220	CYS	7.0
1	D	59	SER	7.0
1	B	126	TYR	6.4
1	D	76	ASP	5.9
1	A	44	GLY	5.8
1	D	122	ASN	5.7
1	D	49	GLY	5.6
1	B	34	ALA	5.5
1	B	45	GLN	5.4
1	A	45	GLN	5.4
1	B	63	GLY	5.1
1	D	123	GLY	4.7
1	B	49	GLY	4.6
1	A	220	CYS	4.6
1	D	22	GLN	4.4
1	C	123	GLY	4.4
1	A	217	PRO	4.3
1	B	25	VAL	4.3
1	B	74	PRO	4.3
1	B	64	SER	4.2
1	C	122	ASN	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	46	GLY	4.1
1	B	24	ALA	4.1
1	D	47	GLY	4.0
1	C	217	PRO	3.9
1	B	57	ASN	3.8
1	C	187	TYR	3.8
1	A	48	PHE	3.7
1	D	64	SER	3.7
1	B	36	ALA	3.5
1	B	26	GLY	3.5
1	B	66	ALA	3.5
1	B	81	PHE	3.5
1	B	23	PHE	3.4
1	A	46	GLY	3.4
1	D	77	ASN	3.2
1	D	294	ASN	3.2
1	A	293	ALA	3.2
1	D	50	CYS	3.2
1	B	294	ASN	3.1
1	D	23	PHE	3.1
1	D	125	SER	3.1
1	C	25	VAL	3.0
1	A	20	ALA	3.0
1	B	41	LEU	3.0
1	B	47	GLY	3.0
1	B	67	PRO	2.9
1	B	35	ALA	2.9
1	C	146	ALA	2.9
1	B	125	SER	2.9
1	B	44	GLY	2.9
1	B	42	PRO	2.8
1	A	65	ASP	2.8
1	D	78	GLY	2.7
1	B	65	ASP	2.7
1	D	65	ASP	2.7
1	B	187	TYR	2.7
1	B	79	PRO	2.7
1	B	293	ALA	2.7
1	D	61	SER	2.7
1	D	44	GLY	2.7
1	D	126	TYR	2.6
1	D	62	VAL	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	59	SER	2.5
1	B	43	ILE	2.5
1	D	58	SER	2.5
1	C	332	GLY	2.5
1	C	50	CYS	2.5
1	D	48	PHE	2.4
1	D	293	ALA	2.4
1	D	56	MET	2.3
1	B	30	THR	2.3
1	C	64	SER	2.3
1	B	50	CYS	2.3
1	A	291	PRO	2.3
1	B	144	ALA	2.3
1	C	120	ASP	2.3
1	A	332	GLY	2.2
1	B	27	GLU	2.2
1	B	37	TRP	2.1
1	C	121	LYS	2.1
1	A	294	ASN	2.1
1	D	57	ASN	2.1
1	A	47	GLY	2.1
1	B	39	VAL	2.1
1	C	63	GLY	2.1
1	D	35	ALA	2.0
1	D	66	ALA	2.0
1	D	28	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	SO4	C	402	5/5	0.90	0.21	7.67	58,67,72,72	0
5	SO4	D	403	5/5	0.96	0.17	2.52	49,49,52,55	0
2	E8V	A	401	25/25	0.79	0.20	0.41	36,49,74,90	0
2	E8V	B	401	25/25	0.76	0.18	0.10	47,56,69,75	0
3	GOL	A	402	6/6	0.93	0.11	0.00	32,43,47,48	0
5	SO4	C	401	5/5	0.97	0.11	-0.08	42,42,51,53	0
2	E8V	D	401	25/25	0.91	0.09	-0.83	27,32,40,43	0
4	CL	B	404	1/1	0.83	0.09	-	59,59,59,59	0
4	CL	A	403	1/1	0.88	0.07	-	59,59,59,59	0
4	CL	D	404	1/1	0.96	0.06	-	70,70,70,70	0
5	SO4	B	402	5/5	0.96	0.24	-	53,54,64,65	0
4	CL	A	404	1/1	0.78	0.05	-	71,71,71,71	0
5	SO4	D	402	5/5	0.97	0.20	-	40,47,48,51	0
4	CL	B	403	1/1	0.94	0.07	-	73,73,73,73	0
4	CL	C	403	1/1	0.88	0.06	-	66,66,66,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.