



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 06:28 am GMT

PDB ID : 3BV0
Title : Crystal Structure of PLP Bound 7,8-Diaminopelargonic Acid Synthase in Mycobacterium Tuberculosis
Authors : Dey, S.; Sacchettini, J.C.
Deposited on : 2008-01-04
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4744 (2.24-2.20)
Clashscore	112137	5509 (2.24-2.20)
Ramachandran outliers	110173	5427 (2.24-2.20)
Sidechain outliers	110143	5428 (2.24-2.20)
RSRZ outliers	101464	4776 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	437	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0;">%</div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 15%, yellow 15%, yellow 77%, green 77%, green 100%);"></div> <div style="position: absolute; bottom: -10px; left: 0;">77% 15% • 6%</div> </div> </div>
1	B	437	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0;">2%</div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, red 0%, red 2%, orange 2%, orange 16%, yellow 16%, yellow 76%, green 76%, green 100%);"></div> <div style="position: absolute; bottom: -10px; left: 0;">76% 16% • 5%</div> </div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6460 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Adenosylmethionine-8-amino-7-oxononanoate aminotransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	410	Total	C	N	O	S	0	0	0
			3079	1962	542	555	20			
1	B	414	Total	C	N	O	S	0	1	0
			3106	1977	549	560	20			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	315	ARG	HIS	SEE REMARK 999	UNP P0A4X6
B	315	ARG	HIS	SEE REMARK 999	UNP P0A4X6

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	108	Total	O	0	0
			108	108		
3	B	137	Total	O	0	0
			137	137		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Adenosylmethionine-8-amino-7-oxononanoate aminotransferase



- Molecule 1: Adenosylmethionine-8-amino-7-oxononanoate aminotransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.05Å 66.47Å 203.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.00 – 2.21 41.72 – 2.21	Depositor EDS
% Data completeness (in resolution range)	92.6 (35.00-2.21) 92.6 (41.72-2.21)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.02 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.187 , 0.238 0.183 , 0.235	Depositor DCC
R_{free} test set	2059 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	25.0	Xtriage
Anisotropy	0.151	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6460	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/3155	0.57	0/4313
1	B	0.43	0/3188	0.58	1/4356 (0.0%)
All	All	0.43	0/6343	0.58	1/8669 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	207	LEU	CA-CB-CG	5.70	128.42	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3079	0	3053	66	0
1	B	3106	0	3080	59	0
2	A	15	0	7	0	0
2	B	15	0	7	0	0
3	A	108	0	0	3	0
3	B	137	0	0	4	0
All	All	6460	0	6147	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (105) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:180:ASP:HB2	1:B:148:ARG:NH1	1.69	1.06
1:B:9:THR:H	1:B:12:GLN:HE21	1.13	0.97
1:A:102:ARG:NE	1:B:6:GLY:HA2	1.79	0.97
1:A:117:THR:HG21	1:B:22:TRP:HE1	1.27	0.96
1:A:180:ASP:HB2	1:B:148:ARG:HH12	1.23	0.95
1:A:117:THR:CG2	1:B:22:TRP:HE1	1.80	0.94
1:A:117:THR:CG2	1:B:22:TRP:NE1	2.32	0.93
1:A:117:THR:CG2	1:B:22:TRP:CD1	2.58	0.87
1:A:117:THR:HG21	1:B:22:TRP:NE1	1.90	0.86
1:A:395:ARG:HG3	1:A:395:ARG:HH21	1.41	0.84
3:A:509:HOH:O	1:B:91:MET:HG2	1.77	0.84
1:A:102:ARG:HE	1:B:6:GLY:HA2	1.37	0.83
1:A:162:PHE:HA	1:A:165:MET:CE	2.15	0.76
1:A:222:VAL:H	1:A:232:HIS:CE1	2.03	0.76
1:B:432:LEU:HA	1:B:435:SER:HB3	1.68	0.74
1:A:395:ARG:CG	1:A:395:ARG:HH21	2.02	0.71
1:B:10:PRO:O	1:B:14:ILE:HG23	1.91	0.71
1:A:117:THR:HG23	1:B:22:TRP:CD1	2.25	0.71
1:A:162:PHE:HA	1:A:165:MET:HE3	1.74	0.70
1:B:357:ASP:OD2	1:B:360:ARG:NH1	2.26	0.69
1:A:286:THR:HG22	1:A:329:SER:OG	1.96	0.66
1:A:66:THR:HB	1:A:288:GLY:HA2	1.78	0.66
1:A:207:LEU:HD12	1:A:207:LEU:O	1.98	0.64
1:B:174:MET:HG2	1:B:177:LEU:HD12	1.81	0.62
1:A:425:ALA:O	1:A:429:VAL:HG12	1.99	0.62
1:A:383:VAL:HG21	1:A:401:PRO:HB2	1.81	0.61
1:A:188:ALA:HA	1:A:206:GLN:HE22	1.65	0.61
1:A:73:HIS:HD2	1:A:75:ALA:H	1.49	0.61
1:A:162:PHE:HA	1:A:165:MET:HE2	1.84	0.59
1:B:110:ILE:HG21	1:B:337:LEU:HD21	1.85	0.59
1:A:140:ARG:HG2	1:A:145:PRO:HB3	1.84	0.58
1:B:342:ARG:NH2	3:B:601:HOH:O	2.29	0.57
1:A:395:ARG:CG	1:A:395:ARG:NH2	2.67	0.56
1:A:20:HIS:HD2	1:B:116:ASP:O	1.89	0.56
1:B:151:MET:HG3	1:B:186:VAL:HG22	1.87	0.55
1:B:188:ALA:HA	1:B:206:GLN:HE22	1.71	0.55
1:A:163:LEU:C	1:A:163:LEU:HD13	2.28	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:158:HIS:HE1	3:B:533:HOH:O	1.89	0.54
1:A:73:HIS:CE1	1:A:335:LEU:HD11	2.42	0.54
1:A:383:VAL:HG22	1:A:403:ARG:O	2.08	0.53
1:B:222:VAL:H	1:B:232:HIS:CE1	2.26	0.53
1:B:65:TRP:HB2	1:B:283:LYS:HD3	1.91	0.53
1:A:364:ALA:HB3	1:A:433:VAL:HG12	1.91	0.52
1:A:207:LEU:HD11	1:A:249:VAL:HG21	1.92	0.52
1:B:9:THR:HG22	1:B:12:GLN:H	1.74	0.52
1:A:244:CYS:HB3	1:A:249:VAL:O	2.09	0.51
1:A:65:TRP:HB2	1:A:283:LYS:HD3	1.92	0.51
1:A:117:THR:HG22	1:B:22:TRP:CD1	2.44	0.50
1:B:45:TRP:CE2	1:B:58:LEU:HD13	2.46	0.50
1:A:383:VAL:HG21	1:A:401:PRO:CB	2.42	0.50
1:A:283:LYS:HG2	1:B:318:THR:HG21	1.94	0.50
1:A:180:ASP:HB2	1:B:148:ARG:HH11	1.69	0.49
1:A:207:LEU:HD12	1:A:207:LEU:C	2.32	0.49
1:B:66:THR:HB	1:B:288:GLY:HA2	1.94	0.49
1:B:9:THR:H	1:B:12:GLN:NE2	1.94	0.49
1:A:73:HIS:CD2	1:A:75:ALA:H	2.31	0.49
1:B:407:TYR:CE2	1:B:409:MET:SD	3.06	0.48
1:A:322:ASN:ND2	1:A:325:ALA:H	2.12	0.48
1:B:50:ARG:O	1:B:51:ASP:OD1	2.32	0.47
1:A:66:THR:HB	1:A:288:GLY:CA	2.43	0.47
1:B:385:LEU:CD2	1:B:401:PRO:HD2	2.44	0.47
1:B:244:CYS:HB3	1:B:249:VAL:O	2.14	0.47
1:A:222:VAL:N	1:A:232:HIS:CE1	2.79	0.47
1:B:50:ARG:NH1	1:B:394:ASP:OD1	2.40	0.47
1:A:207:LEU:HD13	1:A:214:LEU:CD1	2.45	0.46
1:B:429:VAL:O	1:B:433:VAL:HG23	2.15	0.46
1:A:179:THR:O	1:B:148:ARG:NH1	2.49	0.46
1:A:162:PHE:HD2	1:B:129:GLU:OE2	1.98	0.46
1:A:125:SER:OG	1:A:159:GLY:HA3	2.16	0.46
1:A:111:THR:HB	1:A:112:PRO:HD2	1.99	0.45
1:A:291:SER:HB3	1:B:319:PHE:HB2	1.98	0.45
1:A:117:THR:HG22	1:B:22:TRP:NE1	2.26	0.45
1:A:326:CYS:O	1:A:330:VAL:HG23	2.17	0.44
1:B:163:LEU:HD13	1:B:163:LEU:C	2.37	0.44
1:B:34:SER:N	1:B:35:PRO:HD2	2.32	0.44
1:B:102:ARG:HD2	3:B:614:HOH:O	2.17	0.44
1:B:9:THR:HG22	1:B:11:GLU:H	1.82	0.44
1:B:14:ILE:HG22	1:B:37:VAL:CG2	2.48	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:252:ILE:HG12	1:A:278:ILE:HB	1.99	0.44
1:B:303:HIS:HB3	3:B:554:HOH:O	2.18	0.43
1:B:210:HIS:HD2	1:B:213:GLU:OE1	2.00	0.43
1:B:253:PHE:HE1	1:B:274:VAL:HG23	1.83	0.43
1:A:290:LEU:HD22	1:B:290:LEU:HD22	2.00	0.43
1:B:57:VAL:HG12	1:B:396:GLY:HA2	1.99	0.43
1:A:383:VAL:HG23	1:A:383:VAL:O	2.17	0.43
1:A:232:HIS:HD2	3:A:547:HOH:O	2.01	0.43
1:A:89:HIS:HA	1:A:323:PRO:HD2	2.00	0.43
1:B:64:TRP:O	1:B:65:TRP:HB2	2.18	0.43
1:A:103:LEU:HD13	1:A:120:PHE:CE2	2.54	0.43
1:B:9:THR:HG22	1:B:11:GLU:N	2.34	0.42
1:A:364:ALA:CB	1:A:433:VAL:HG12	2.49	0.42
1:A:45:TRP:CE2	1:A:58:LEU:HD13	2.55	0.42
1:A:322:ASN:HD22	1:A:325:ALA:H	1.68	0.41
1:B:81:THR:O	1:B:85:ARG:HG3	2.20	0.41
1:A:357:ASP:O	1:A:360:ARG:HB2	2.21	0.41
1:A:110:ILE:HG21	1:A:337:LEU:HD21	2.01	0.41
1:A:343:THR:O	1:A:347:GLU:HG3	2.21	0.41
1:A:422:ILE:O	1:A:426:MET:HG3	2.20	0.41
1:B:169:ASP:HA	1:B:170:PRO:HD3	1.91	0.41
1:B:322:ASN:ND2	1:B:325:ALA:H	2.19	0.41
1:A:98:GLU:HB2	1:B:13:ILE:HD11	2.03	0.41
1:B:409:MET:N	1:B:409:MET:SD	2.94	0.40
1:B:218:VAL:HG13	1:B:218:VAL:O	2.21	0.40
1:A:122:ASP:HB2	3:A:607:HOH:O	2.21	0.40
1:A:117:THR:O	1:A:296:LEU:HA	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	404/437 (92%)	393 (97%)	10 (2%)	1 (0%)	51	58
1	B	409/437 (94%)	397 (97%)	9 (2%)	3 (1%)	25	24
All	All	813/874 (93%)	790 (97%)	19 (2%)	4 (0%)	32	34

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	283	LYS
1	A	283	LYS
1	B	7	GLY
1	B	228	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	314/329 (95%)	297 (95%)	17 (5%)	26	29
1	B	317/329 (96%)	300 (95%)	17 (5%)	26	29
All	All	631/658 (96%)	597 (95%)	34 (5%)	26	29

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	LEU
1	A	64	TRP
1	A	80	LEU
1	A	103	LEU
1	A	142	ARG
1	A	153	TRP
1	A	177	LEU
1	A	207	LEU
1	A	240	LEU
1	A	335	LEU
1	A	337	LEU
1	A	340	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	358	THR
1	A	360	ARG
1	A	395	ARG
1	A	409	MET
1	A	429	VAL
1	B	9	THR
1	B	14	ILE
1	B	58	LEU
1	B	64	TRP
1	B	140	ARG
1	B	153	TRP
1	B	160	ASP
1	B	174	MET
1	B	175	HIS
1	B	180	ASP
1	B	186	VAL
1	B	207	LEU
1	B	246	ARG
1	B	250	LEU
1	B	335	LEU
1	B	435	SER
1	B	436	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	HIS
1	A	42	HIS
1	A	53	GLN
1	A	73	HIS
1	A	97	HIS
1	A	206	GLN
1	A	209	GLN
1	A	232	HIS
1	A	322	ASN
1	B	12	GLN
1	B	158	HIS
1	B	206	GLN
1	B	210	HIS
1	B	232	HIS
1	B	271	HIS
1	B	322	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	PLP	A	501	1	15,15,16	1.12	2 (13%)	20,22,23	1.32	3 (15%)
2	PLP	B	502	1	15,15,16	0.97	1 (6%)	20,22,23	1.02	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	501	1	-	0/6/6/8	0/1/1/1
2	PLP	B	502	1	-	0/6/6/8	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	PLP	C3-C2	-2.46	1.39	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	502	PLP	C2-N1	2.05	1.38	1.33
2	A	501	PLP	C2-N1	2.18	1.38	1.33

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	PLP	C4A-C4-C3	-2.23	116.70	120.54
2	A	501	PLP	O2P-P-O4P	2.08	112.26	106.73
2	A	501	PLP	O4P-C5A-C5	2.99	115.33	109.32

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	410/437 (93%)	-0.16	5 (1%) 79 77	9, 21, 46, 61	9 (2%)
1	B	414/437 (94%)	-0.36	7 (1%) 70 68	7, 18, 39, 60	6 (1%)
All	All	824/874 (94%)	-0.26	12 (1%) 74 72	7, 20, 43, 61	15 (1%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	6	GLY	6.4
1	B	7	GLY	4.0
1	A	361	ALA	3.7
1	B	435	SER	3.5
1	B	51	ASP	3.1
1	B	179	THR	3.0
1	B	180	ASP	3.0
1	A	307	ALA	2.9
1	B	176	SER	2.6
1	A	179	THR	2.4
1	A	365	VAL	2.4
1	A	433	VAL	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	PLP	A	501	15/16	0.97	0.20	0.62	15,20,29,31	0
2	PLP	B	502	15/16	0.96	0.11	-1.03	8,15,22,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.