



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 9, 2018 – 12:16 AM EST

PDB ID : 4BWZ  
Title : Crystal structure of the sodium proton antiporter, NapA  
Authors : Lee, C.; Drew, D.; Cameron, A.D.  
Deposited on : 2013-07-05  
Resolution : 2.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030736  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030736


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2168 (3.00-2.96)
Clashscore	112137	2535 (3.00-2.96)
Ramachandran outliers	110173	2451 (3.00-2.96)
Sidechain outliers	110143	2454 (3.00-2.96)
RSRZ outliers	101464	2192 (3.00-2.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	394	<div> <div>7%</div> <div>77%</div> <div>19%</div> <div>• •</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2824 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NA(+)/H(+) ANTIPORTER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	384	Total	C	N	O	S	0	0	0
			2822	1876	450	487	9			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	CYS	MET	engineered mutation	UNP Q72IM4
A	166	CYS	VAL	engineered mutation	UNP Q72IM4
A	326	CYS	VAL	engineered mutation	UNP Q72IM4
A	387	GLY	-	expression tag	UNP Q72IM4
A	388	SER	-	expression tag	UNP Q72IM4
A	389	GLU	-	expression tag	UNP Q72IM4
A	390	ASN	-	expression tag	UNP Q72IM4
A	391	LEU	-	expression tag	UNP Q72IM4
A	392	TYR	-	expression tag	UNP Q72IM4
A	393	PHE	-	expression tag	UNP Q72IM4
A	394	GLN	-	expression tag	UNP Q72IM4


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NA(+)/H(+) ANTIPORTER



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.75Å 82.15Å 191.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.95 – 2.98 52.76 – 2.98	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.95-2.98) 99.9 (52.76-2.98)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.79 (at 3.01Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.222 , 0.248 0.229 , 0.267	Depositor DCC
$R_{free}$ test set	584 reflections (4.78%)	DCC
Wilson B-factor (Å <sup>2</sup> )	91.4	Xtriage
Anisotropy	0.269	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 96.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.42$ , $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	2824	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	107.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.38	0/2869	0.54	1/3907 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	68	LEU	CA-CB-CG	-5.35	102.98	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2822	0	3031	45	0
2	A	2	0	0	0	0
All	All	2824	0	3031	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:LEU:HA	1:A:46:LEU:HD12	1.67	0.76
1:A:348:VAL:HG13	1:A:352:GLU:HB2	1.71	0.71
1:A:67:LEU:HA	1:A:265:GLU:HG3	1.75	0.68
1:A:68:LEU:HD13	1:A:238:ILE:HG12	1.77	0.67
1:A:64:ALA:HB1	1:A:238:ILE:HD11	1.81	0.61
1:A:65:VAL:HG13	1:A:241:ALA:HB2	1.83	0.60
1:A:212:VAL:HG11	1:A:218:PHE:HB2	1.84	0.60
1:A:275:MET:O	1:A:279:ARG:HG3	2.08	0.53
1:A:39:GLY:HA3	1:A:274:ALA:O	2.08	0.52
1:A:105:TYR:CZ	1:A:109:ILE:HD11	2.45	0.52
1:A:136:GLN:HG3	1:A:141:LEU:HD11	1.92	0.52
1:A:130:ILE:HG12	1:A:273:PHE:HE1	1.78	0.49
1:A:135:LEU:HD13	1:A:146:SER:HB2	1.94	0.49
1:A:33:ILE:O	1:A:37:LEU:HG	2.14	0.48
1:A:98:PRO:HB2	1:A:123:LEU:HB3	1.95	0.48
1:A:138:LEU:HD12	1:A:375:ILE:HD12	1.97	0.47
1:A:309:GLY:HA3	1:A:325:GLY:HA2	1.96	0.47
1:A:357:LEU:HA	1:A:360:VAL:HG22	1.97	0.47
1:A:41:LEU:O	1:A:46:LEU:HB2	2.16	0.46
1:A:12:TYR:HB3	1:A:42:VAL:HG11	1.97	0.46
1:A:61:GLU:OE1	1:A:237:PRO:HG2	2.17	0.45
1:A:114:LEU:HB3	1:A:115:PRO:HD3	1.98	0.45
1:A:106:GLY:O	1:A:111:PHE:HB2	2.17	0.44
1:A:136:GLN:NE2	1:A:265:GLU:OE2	2.46	0.44
1:A:178:VAL:O	1:A:182:THR:HG23	2.17	0.44
1:A:13:LEU:HG	1:A:50:VAL:HG21	2.01	0.43
1:A:191:PHE:HE1	1:A:229:LEU:HD23	1.83	0.43
1:A:294:GLY:O	1:A:298:THR:OG1	2.31	0.43
1:A:344:LYS:C	1:A:346:GLY:H	2.22	0.43
1:A:105:TYR:CE2	1:A:109:ILE:HD11	2.53	0.42
1:A:32:VAL:HG13	1:A:133:ARG:HG2	2.00	0.42
1:A:24:PHE:CZ	1:A:37:LEU:HD12	2.54	0.42
1:A:79:ASP:HB3	1:A:147:ARG:HD3	2.01	0.42
1:A:190:VAL:O	1:A:194:LEU:HB2	2.19	0.42
1:A:123:LEU:O	1:A:331:ARG:NH1	2.53	0.42
1:A:199:SER:HB3	1:A:247:LEU:HA	2.02	0.41
1:A:73:LEU:HD23	1:A:73:LEU:HA	1.87	0.41
1:A:68:LEU:HD13	1:A:238:ILE:HG23	2.03	0.41
1:A:210:LEU:HA	1:A:211:PRO:HD3	1.85	0.41
1:A:130:ILE:HD12	1:A:364:THR:HG23	2.03	0.41
1:A:15:LEU:HD12	1:A:15:LEU:HA	1.89	0.40
1:A:149:ILE:HG13	1:A:324:VAL:HG22	2.03	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138:LEU:HD11	1:A:372:LYS:HG3	2.04	0.40
1:A:371:LEU:O	1:A:375:ILE:HG13	2.21	0.40
1:A:42:VAL:HG13	1:A:49:LEU:HB2	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	382/394 (97%)	361 (94%)	18 (5%)	3 (1%)	22 62

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	288	PRO
1	A	141	LEU
1	A	223	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	289/298 (97%)	277 (96%)	12 (4%)	34 72

All (12) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	A	5	GLU
1	A	32	VAL
1	A	68	LEU
1	A	73	LEU
1	A	194	LEU
1	A	207	LEU
1	A	233	ILE
1	A	238	ILE
1	A	251	VAL
1	A	253	GLU
1	A	331	ARG
1	A	359	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	384/394 (97%)	0.33	27 (7%) 17 9	74, 105, 139, 168	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	343	LEU	4.2
1	A	348	VAL	4.2
1	A	286	ALA	3.3
1	A	138	LEU	3.2
1	A	105	TYR	3.0
1	A	256	ARG	2.9
1	A	339	ALA	2.9
1	A	139	GLY	2.9
1	A	341	LEU	2.9
1	A	386	GLU	2.8
1	A	384	ALA	2.8
1	A	140	VAL	2.7
1	A	342	GLY	2.7
1	A	285	LEU	2.6
1	A	77	LEU	2.6
1	A	109	ILE	2.5
1	A	89	LEU	2.5
1	A	318	VAL	2.5
1	A	382	ARG	2.5
1	A	340	ALA	2.4
1	A	319	ARG	2.3
1	A	111	PHE	2.3
1	A	112	GLU	2.2
1	A	118	PHE	2.2
1	A	197	PHE	2.2
1	A	263	ALA	2.2
1	A	107	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	A	1387	1/1	0.91	0.19	-	131,131,131,131	0
2	ZN	A	1388	1/1	0.80	0.26	-	128,128,128,128	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.