



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 23, 2018 – 06:27 PM EST

PDB ID : 1C1G
Title : CRYSTAL STRUCTURE OF TROPOMYOSIN AT 7 ANGSTROMS RESOLUTION IN THE SPERMINE-INDUCED CRYSTAL FORM
Authors : Whitby, F.G.; Phillips Jr., G.N.
Deposited on : 1999-07-22
Resolution : 7.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030736
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 7.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	1035 (10.00-3.80)
Ramachandran outliers	110173	1003 (10.00-3.76)
Sidechain outliers	110143	1098 (10.00-3.70)
RSRZ outliers	101464	1003 (10.00-3.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	284	
1	B	284	
1	C	284	
1	D	284	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 9160 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called TROPOMYOSIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	284	Total	C	N	O	S	0	0	0
			2290	1402	387	494	7			
1	B	284	Total	C	N	O	S	0	0	0
			2290	1402	387	494	7			
1	C	284	Total	C	N	O	S	0	0	0
			2290	1402	387	494	7			
1	D	284	Total	C	N	O	S	0	0	0
			2290	1402	387	494	7			


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TROPOMYOSIN




• Molecule 1: TROPOMYOSIN






• Molecule 1: TROPOMYOSIN



• Molecule 1: TROPOMYOSIN



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	259.74Å 55.30Å 136.26Å 90.00° 97.42° 90.00°	Depositor
Resolution (Å)	100.00 – 7.00 99.89 – 7.00	Depositor EDS
% Data completeness (in resolution range)	96.4 (100.00-7.00) 96.4 (99.89-7.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.35 (at 6.73Å)	Xtriage
Refinement program	XTALVIEW, X-PLOR	Depositor
R, R_{free}	0.404 , (Not available) 0.413 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	210.9	Xtriage
Anisotropy	1.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.54 , 457.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.34$, $\langle L^2 \rangle = 0.17$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.67	EDS
Total number of atoms	9160	wwPDB-VP
Average B, all atoms (Å ²)	174.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	3/2299 (0.1%)	1.79	56/3062 (1.8%)
1	B	0.80	2/2299 (0.1%)	1.63	41/3062 (1.3%)
1	C	1.28	11/2299 (0.5%)	1.86	59/3062 (1.9%)
1	D	0.91	8/2299 (0.3%)	1.82	67/3062 (2.2%)
All	All	0.98	24/9196 (0.3%)	1.78	223/12248 (1.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	C	0	5
1	D	0	5
All	All	0	13

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	852	ILE	N-CA	30.96	2.08	1.46
1	C	852	ILE	CA-CB	21.53	2.04	1.54
1	C	851	SER	C-N	15.91	1.70	1.34
1	C	572	ILE	CA-CB	11.53	1.81	1.54
1	D	859	LYS	N-CA	10.74	1.67	1.46

The worst 5 of 223 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	4	ILE	CA-CB-CG2	-21.88	67.14	110.90
1	C	572	ILE	N-CA-CB	-21.77	60.73	110.80
1	B	568	ILE	N-CA-CB	-21.05	62.38	110.80
1	C	852	ILE	CB-CA-C	-18.23	75.14	111.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	223	GLU	CA-C-N	-17.61	78.46	117.20

There are no chirality outliers.

5 of 13 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	223	GLU	Mainchain,Peptide
1	A	224	GLU	Mainchain
1	C	702	ALA	Mainchain
1	C	703	GLN	Mainchain
1	C	705	ASP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2290	0	2295	96	0
1	B	2290	0	2292	87	0
1	C	2290	0	2291	120	3
1	D	2290	0	2292	84	3
All	All	9160	0	9170	330	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 330 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:572:ILE:CA	1:C:572:ILE:CB	1.81	1.58
1:D:859:LYS:CA	1:D:859:LYS:N	1.67	1.51
1:C:705:ASP:C	1:C:705:ASP:CA	1.76	1.50
1:C:572:ILE:CB	1:C:572:ILE:N	1.71	1.46
1:C:851:SER:C	1:C:852:ILE:N	1.70	1.44

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:851:SER:O	1:D:857:LYS:O[3_364]	2.14	0.06
1:C:848:ASP:O	1:D:856:ILE:O[3_364]	2.16	0.04
1:C:852:ILE:N	1:D:860:MET:N[3_364]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/284 (99%)	241 (86%)	32 (11%)	9 (3%)	5	35
1	B	282/284 (99%)	250 (89%)	28 (10%)	4 (1%)	13	54
1	C	282/284 (99%)	246 (87%)	30 (11%)	6 (2%)	8	45
1	D	282/284 (99%)	246 (87%)	28 (10%)	8 (3%)	6	39
All	All	1128/1136 (99%)	983 (87%)	118 (10%)	27 (2%)	7	42

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	570	ASP
1	C	845	ALA
1	D	854	ASP
1	D	880	ASP
1	D	1133	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/245 (100%)	209 (85%)	36 (15%)	3	20
1	B	245/245 (100%)	220 (90%)	25 (10%)	8	33
1	C	245/245 (100%)	217 (89%)	28 (11%)	7	28
1	D	245/245 (100%)	212 (86%)	33 (14%)	4	24
All	All	980/980 (100%)	858 (88%)	122 (12%)	5	26

5 of 122 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	552	LYS
1	C	634	ASP
1	D	1038	SER
1	B	556	GLU
1	C	572	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	203	ASN
1	B	431	GLN
1	D	1055	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	C	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	851:SER	C	852:ILE	N	1.70

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	284/284 (100%)	3.25	200 (70%) 0 3	174, 174, 174, 174	0
1	B	284/284 (100%)	3.70	237 (83%) 0 2	174, 174, 174, 174	0
1	C	284/284 (100%)	3.41	192 (67%) 0 3	174, 174, 174, 174	0
1	D	284/284 (100%)	2.96	185 (65%) 0 3	174, 174, 174, 174	0
All	All	1136/1136 (100%)	3.33	814 (71%) 0 3	174, 174, 174, 174	0

The worst 5 of 814 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	657	ASN	14.7
1	A	2	ASP	14.6
1	C	596	ASP	12.6
1	A	36	SER	11.5
1	C	658	ARG	11.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.